

ENLARGED MIXED SHIMURA VARIETIES, BI-ALGEBRAIC SYSTEM AND SOME AX TYPE TRANSCENDENTAL RESULTS

ZIYANG GAO

ABSTRACT. We develop a theory of enlarged mixed Shimura varieties, putting the universal vectorial bi-extension defined by Coleman into this framework to study some functional transcendental results of Ax type. We study their bi-algebraic systems, formulate the Ax-Schanuel conjecture and explain its relation with the Ax logarithmique theorem and the Ax-Lindemann theorem which we shall prove. We also prove the whole Ax-Schanuel conjecture for the unipotent part. All these bi-algebraic and transcendental results generalize their counterparts for mixed Shimura varieties. In the end we briefly discuss about the André-Oort and Zilber-Pink type problems for enlarged mixed Shimura varieties.

CONTENTS

1. Introduction	1
2. Pink's work on equivariant families of mixed Hodge structures	8
3. Enlarged mixed Shimura data	12
4. Enlarged mixed Shimura varieties	20
5. Bi-algebraic setting for enlarged mixed Shimura varieties	24
6. The Ax logarithmique theorem	30
7. The Ax-Lindemann theorem	32
8. The Ax-Schanuel conjecture	33
9. Proof of Ax-Schanuel when restricted to a fiber of $S^{\natural} \rightarrow S_G$	35
10. Special subvarieties	48
11. Appendix: Proof of Ax-Lindemann (Theorem 7.1)	50
References	57

1. INTRODUCTION

1.1. Motivations: From Manin-Mumford and André-Oort to bi-algebraicity. We start with two famous conjectures in arithmetic geometry: the Manin-Mumford and the André-Oort conjectures.

	Manin-Mumford	André-Oort
object	Abelian variety A and a subvariety Y $\Sigma := \text{set of torsion points of } A$	Shimura variety S and a subvariety Y $\Sigma := \text{set of special points of } S$
hypothesis	$(Y \cap \Sigma)^{\text{Zar}} = Y$	$(Y \cap \Sigma)^{\text{Zar}} = Y$
conclusion	Y is a torsion coset	Y is a Shimura subvariety

By standard specialization argument it suffices to prove Manin-Mumford for abelian varieties over $\overline{\mathbb{Q}}$. These two conjectures are independent, although similar. The mixed André-Oort conjecture, replacing Shimura varieties by mixed Shimura varieties, partly unifies the previous two conjectures because it implies Manin-Mumford for CM abelian varieties. Now in the spirit of [40] we make the following conjecture in terms of the bi-algebraic systems associated with abelian varieties and Shimura varieties, which unifies Manin-Mumford and André-Oort completely.

Conjecture 1.1. *Let S be an algebraic variety over $\overline{\mathbb{Q}}$ such that its uniformization $\mathcal{X}$ has a structure of algebraic variety over $\overline{\mathbb{Q}}$. Suppose Y is a subvariety of S containing a Zariski dense subset of arithmetically bi-algebraic points, then Y is geometrically bi-algebraic.*

Let us explain this conjecture for $\mathcal{A}_g$. Its uniformization is $\text{unif}: \mathcal{H}_g^+ \rightarrow \mathcal{A}_g$ where $\mathcal{H}_g^+ = \{Z = X + \sqrt{-1}Y \in M_{2g \times 2g}(\mathbb{C}) \mid Z = Z^t, Y > 0\}$. We say that $s \in \mathcal{A}_g(\overline{\mathbb{Q}})$ is **arithmetically bi-algebraic** if $\text{unif}^{-1}(s) \subset \mathcal{H}_g^+ \cap M_{2g \times 2g}(\overline{\mathbb{Q}})$. A theorem of Cohen [11] and Shiga-Wolfart [36] asserts that s is arithmetically bi-algebraic iff s is a special point, i.e. the abelian variety parametrized by s is CM. Hence the set Σ in the André-Oort conjecture for $\mathcal{A}_g$ equals the set of arithmetically bi-algebraic points. On the other hand $\mathcal{H}_g^+$ is a semi-algebraic subset of $\mathbb{C}^{g(g+1)/2}$, open in the usual topology. We say that a subvariety Y of $\mathcal{A}_g$ is **geometrically bi-algebraic** if one (and hence all) complex analytic irreducible component of $\text{unif}^{-1}(Y)$ is algebraizable, i.e. it has the same dimension as its Zariski closure in $\mathbb{C}^{g(g+1)/2}$. By Ullmo-Yafaev [41] the geometrically bi-algebraic subvarieties of $\mathcal{A}_g$ containing special points are precisely the Shimura subvarieties of $\mathcal{A}_g$. Hence Conjecture 1.1 is equivalent to the André-Oort conjecture when $S = \mathcal{A}_g$.

The situation for abelian varieties A over $\overline{\mathbb{Q}}$ is more complicated. We wish to endow a $\overline{\mathbb{Q}}$ -structure on $\text{Lie}A_{\mathbb{C}}$ so that torsion points of A are precisely arithmetically bi-algebraic points. However using the Schneider-Lang and Wüstholz' analytic subgroup theorems, Ullmo [40, Proposition 2.6] proved: any torsion point of A , except the origin, becomes transcendental in $\text{Lie}A_{\mathbb{C}}$. A solution to this problem is proposed by Bost (see [40, §2.2.2]): instead of A we study its universal vector extension $A^{\natural}$. We briefly recall some basic facts about $A^{\natural}$.

By a *vector extension* of A , we mean an algebraic group E such that there exist a vector group W and an exact sequence $0 \rightarrow W \rightarrow E \rightarrow A \rightarrow 0$. There exists a universal vector extension $A^{\natural}$ of A such that any vector extension E of A is obtained as:

$$\begin{array}{ccccccc} 0 & \longrightarrow & W_A & \longrightarrow & A^{\natural} & \longrightarrow & A \longrightarrow 0 \\ & & \vdots & & \downarrow & & \downarrow = \\ 0 & \longrightarrow & W & \longrightarrow & E & \longrightarrow & A \longrightarrow 0 \end{array}$$

In fact $A^{\natural}$ is constructed as follows: Let $\Gamma := H_1(A(\mathbb{C}), \mathbb{Z}) \subset H_1(A(\mathbb{C}), \mathbb{C})$ be the period lattice of A . For the Hodge decomposition $H_1(A(\mathbb{C}), \mathbb{C}) = H^{0,-1}(A_{\mathbb{C}}) \oplus H^{-1,0}(A_{\mathbb{C}})$, the holomorphic part $H^{-1,0}(A_{\mathbb{C}})$ equals the tangent space of A at 0, and the anti-holomorphic part $H^{0,-1}(A_{\mathbb{C}})$ equals $\Omega_{A_{\mathbb{C}}}^1$. We have

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^{0,-1}(A_{\mathbb{C}}) & \longrightarrow & H_1(A(\mathbb{C}), \mathbb{C}) & \longrightarrow & H^{-1,0}(A_{\mathbb{C}}) \longrightarrow 0 \\ & & \downarrow = & & \downarrow \text{unif}^{\natural} & & \downarrow \text{unif} \\ 0 & \longrightarrow & \Omega_{A_{\mathbb{C}}}^1 & \longrightarrow & A^{\natural}(\mathbb{C}) \simeq \Gamma \backslash H_1(A(\mathbb{C}), \mathbb{C}) & \longrightarrow & A(\mathbb{C}) \simeq \Gamma \backslash H^{-1,0}(A_{\mathbb{C}}) \longrightarrow 0 \end{array}$$

Take the $\overline{\mathbb{Q}}$ -structure $H_1(A(\mathbb{C}), \mathbb{Z}) \otimes \overline{\mathbb{Q}}$ on $H_1(A(\mathbb{C}), \mathbb{C})$. As an application of Wüstholz' analytic subgroup theorem [45, Theorem 1], Ullmo [40, Théorème 2.10] proved

$$z \in H_1(A(\mathbb{C}), \mathbb{Z}) \otimes \overline{\mathbb{Q}} \text{ such that } \text{unif}^{\natural}(z) \in A^{\natural}(\overline{\mathbb{Q}}) \Leftrightarrow \text{unif}^{\natural}(z) \text{ is a torsion point of } A^{\natural}.$$

Thus we get an arithmetic bi-algebraic description for the torsion points of A because the projection $A^{\natural} \rightarrow A$ induces a bijection between their torsion points. This suggests that in view of Conjecture 1.1, the abelian varieties are not the good objects to study. Instead, one should study their universal vector extensions. We will characterize geometrically bi-algebraic subvarieties of $A^{\natural}$ in the paper, from which we can see that Conjecture 1.1 for $A^{\natural}$ implies the Manin-Mumford conjecture without much effort.

1.2. Motivation: Vector extensions, from Grothendieck to Laumon. The study of $A^{\natural}$ goes back to Grothendieck ($\natural$ -extensions) and he studied the Gauß-Manin connection on $\text{Lie}(\mathfrak{A}^{\natural}/B)$ for any abelian scheme $\mathfrak{A}/B$. Later on Buium [8, 10] studied arithmetic differential equations on $\mathfrak{A}^{\natural}$, making $\mathfrak{A}^{\natural}$ a differential algebraic group (algebraic D -group), although the idea was already used by Manin for his work on algebraic curves over function fields. These are extra tools which do not exist for $\mathfrak{A}$, and thus it is often convenient or even necessary to work with $\mathfrak{A}^{\natural}$ in order to prove properties for $\mathfrak{A}$. Examples of this fact include, but are not limited to: Manin's proof of the geometric Mordell conjecture [23, 24], Buium's effective bound for the geometric Lang conjecture [9], Bertrand-Pillay's relative Lindemann-Weierstrass theorem of [6], Bertrand-Masser-Pillay-Zannier's proof of the relative Manin-Mumford conjecture for semi-abelian surfaces [5], results on Galois groups and Manin maps of Bertrand-Pillay [7], etc.

A 1-motive over $\mathbb{C}$, after Deligne, consists of a complex semi-abelian variety G and a group homomorphism $\mathbb{Z}^n \rightarrow G$. However Deligne's 1-motives do not allow vector extensions. To fix this, Laumon [22] defined the *Laumon 1-motives* by replacing $\mathbb{Z}^n$ by a formal group F . The identity component F° of F gives rise to vector extensions. We refer to [4] for more details on (realizations of) Laumon 1-motives.

The moduli space of Deligne 1-motives is a mixed Shimura variety. Based on works of Deligne, Milne, Brylinski and others, Pink [33] finished the framework for the study of mixed Shimura varieties. However mixed Shimura varieties do NOT allow any vector extension. In particular, let $\mathfrak{A}_g$ be the universal family over $\mathcal{A}_g$, where $\mathcal{A}_g$ is the moduli space of principally polarized abelian varieties with level-4-structure, and let $\mathfrak{A}_g^{\natural}$ be the universal vector extension of the abelian scheme $\mathfrak{A}_g/\mathcal{A}_g$. Then $\mathfrak{A}_g^{\natural}$ is NOT a mixed Shimura variety. This suggests that in order to parametrize Laumon 1-motives we must extend this notion. Note that $\mathfrak{A}_g^{\natural}$ is still not large enough: it cannot give any information on the weight -2 part of Laumon 1-motives, i.e. the toric part of G .

1.3. Goals of this paper. In §1.4-§1.6 of the Introduction, we briefly explain the three main goals of the paper, each one occupying a subsection.

1.4. Moduli space in the Deligne-Pink language: enlarged mixed Shimura varieties. The *first main goal* of this paper is to develop the theory of enlarged mixed Shimura varieties, in order to allow vector extensions and parametrize Laumon 1-motives. This is done in §3 and §4. We define a pair, called an enlarged mixed Shimura datum, consisting of a $\mathbb{Q}$ -group P and a simply-connected complex analytic space $\mathcal{X}^{\natural}$ satisfying some properties (see Definition 3.1), and define an enlarged mixed Shimura variety to be the quotient of $\mathcal{X}^{\natural}$ by a congruence subgroup of

$P(\mathbb{Q})$. Then we prove that any such defined enlarged mixed Shimura variety admits a canonical structure of algebraic variety over a number field canonically associated to the enlarged mixed Shimura datum. This is exactly what Deligne and Pink did for (mixed) Shimura varieties, so we call this language of enlarged mixed Shimura data and enlarged mixed Shimura varieties the **Deligne-Pink language**. The importance of the Deligne-Pink language for the study of mixed Shimura varieties is revealed by almost any literature on (mixed) Shimura varieties. As for enlarged mixed Shimura varieties, let me just point out that every result we prove in this paper (characterization of geometrically bi-algebraic subvarieties, the Ax logarithmique theorem, the Ax-Lindemann theorem and the Ax-Schanuel conjecture for the unipotent part) relies heavily on this language.

The $\mathfrak{A}_g^{\natural}$ defined at the end of last subsection is an example of enlarged mixed Shimura varieties which are not mixed Shimura varieties. Another such example is the universal vectorial bi-extension $\mathfrak{P}_g^{\natural}$ studied by Coleman [12] (see Example 4.5(2)), which is defined as follows in geometric terms: Let $\mathfrak{P}_g$ be the universal Poincaré biextension, i.e. the $\mathbb{G}_m$ -torsor over $\mathfrak{A}_g \times \mathfrak{A}_g^{\vee}$ whose fiber $(\mathfrak{P}_g)_a$ for any point $a \in \mathcal{A}_g$ is the Poincaré biextension over $(\mathfrak{A}_g)_a \times (\mathfrak{A}_g^{\vee})_a$. Let $\mathfrak{A}_g^{\natural}$ be the universal vector extension of the abelian scheme $\mathfrak{A}_g/\mathcal{A}_g$. Then $\mathfrak{P}_g^{\natural}$ is the pullback of $\mathfrak{P}_g$ by $\mathfrak{A}_g^{\natural} \times (\mathfrak{A}_g^{\vee})^{\natural} \rightarrow \mathfrak{A}_g \times \mathfrak{A}_g^{\vee}$. The typical example of enlarged mixed Shimura variety to keep in mind is $(\mathfrak{P}_g^{\natural})^{[n]}$, i.e. the n -fiber product of $\mathfrak{P}_g^{\natural}$ over $\mathfrak{A}_g^{\natural} \times (\mathfrak{A}_g^{\vee})^{\natural}$. The advantage of $(\mathfrak{P}_g^{\natural})^{[n]}$ to $\mathfrak{A}_g^{\natural}$ is that it also reflects information on the weight -2 part of Laumon 1-motives.

1.5. Functional transcendental results: Ax-Schanuel. Let $S^{\natural}$ be a connected enlarged mixed Shimura variety and let $\text{unif}^{\natural}: \mathcal{X}^{\natural+} \rightarrow S^{\natural}$ be its uniformization. A typical example to keep in mind is $(\mathfrak{P}_g^{\natural})^{[n]}$ or, simpler, $\mathfrak{A}_g^{\natural}$. As we shall see in §5.1, there exists a complex algebraic variety $\mathcal{X}^{\natural,\vee}$ such that $\mathcal{X}^{\natural+}$ can be embedded as a semi-algebraic open subset (in the usual topology) of $\mathcal{X}^{\natural,\vee}$. The **second main goal** of this paper is to study some functional transcendental results. We start with (Theorem 6.3 and §7):

Theorem 1.2 (Ax logarithmique). *Let $Z^{\natural}$ be an irreducible subvariety of $S^{\natural}$ and let $\tilde{Z}^{\natural}$ be a complex analytic irreducible component of $(\text{unif}^{\natural})^{-1}(Z^{\natural})$. Then the image of $(\tilde{Z}^{\natural})^{\text{Zar}}$, the Zariski closure of $\tilde{Z}^{\natural}$ in $\mathcal{X}^{\natural+}$,^[1] under $\text{unif}^{\natural}$ is quasi-linear.*

Theorem 1.3 (Ax-Lindemann). *Let $\tilde{Z}^{\natural}$ be a semi-algebraic subset of $\mathcal{X}^{\natural+}$. Then any irreducible component of $\text{unif}^{\natural}(\tilde{Z}^{\natural})^{\text{Zar}}$ is quasi-linear.*

Quasi-linear subvarieties of $S^{\natural}$ will be defined in the next subsection of the Introduction (Definition 1.7). Theorem 1.2 is proven for pure Shimura varieties by Moonen [25, 3.6, 3.7] and for mixed Shimura varieties by the author [16, Theorem 8.1]. Theorem 1.3 plays an essential role in the proof of the André-Oort conjecture. It is proven for $Y(1)^n$ by Pila [29], for projective pure Shimura varieties by Ullmo-Yafaev [43], for $\mathcal{A}_g$ by Pila-Tsimerman [31], for any pure Shimura variety by Klingler-Ullmo-Yafaev [21], and for any mixed Shimura variety by the author [16, Theorem 1.2].

We make some comparison of the proofs of Theorem 1.2 and Theorem 1.3 with the author's previous work on mixed Shimura varieties [16]. In both situations it is important to study the

^[1]It means that $(\tilde{Z}^{\natural})^{\text{Zar}}$ is the complex analytic irreducible component of $\mathcal{X}^{\natural+} \cap (\text{Zariski closure of } \tilde{Z}^{\natural} \text{ in } \mathcal{X}^{\natural,\vee})$ which contains $\tilde{Z}^{\natural}$.

geometric bi-algebraic systems associated with the ambient varieties (we will elaborate this in the next subsection). We need the characterization of geometrically bi-algebraic subvarieties in both situations. For mixed Shimura varieties, Ax logarithmique follows from André's result on the algebraic monodromy groups without much effort and the characterization comes as a byproduct. But for enlarged mixed Shimura varieties, Hodge theory itself is not enough. We should study the geometry of $S^{\natural}$ more carefully and it is better to separate the proofs into two parts: first prove that geometrically bi-algebraic subvarieties of $S^{\natural}$ are precisely quasi-linear subvarieties, then prove that the target objects in the conclusions are geometrically bi-algebraic. The first part is new compared to [16], but the second part is only a slight modification of [16].

A common generalization of Theorem 1.2 and Theorem 1.3 is the following conjecture.

Conjecture 1.4 (Ax-Schanuel). *Let $\Delta^{\natural} \subset \mathcal{X}^{\natural+} \times S^{\natural}$ be the graph of $\text{unif}^{\natural}$. Let $\mathcal{Z}^{\natural} = \text{graph}(\tilde{Z}^{\natural} \xrightarrow{\text{unif}^{\natural}} Z^{\natural})$ be a complex analytic irreducible subvariety of $\Delta^{\natural}$. Let $F^{\natural}$ be the smallest quasi-linear subvariety of $S^{\natural}$ which contains $Z^{\natural}$. Let $\tilde{F}^{\natural}$ be the complex analytic irreducible component of $(\text{unif}^{\natural})^{-1}(F^{\natural})$ which contains $\tilde{Z}^{\natural}$. Then*

- (1) $\dim(\tilde{Z}^{\natural})^{\text{Zar}} + \dim(Z^{\natural})^{\text{Zar}} - \dim \tilde{Z}^{\natural} \geq \dim F^{\natural}$.
- (2) Let $\mathcal{B}^{\natural} := (\mathcal{Z}^{\natural})^{\text{Zar}} \subset \mathcal{X}^{\natural+} \times S^{\natural}$. Then $\dim \mathbf{pr}_{F^{\natural}}^{\text{lin}}(\mathcal{B}^{\natural}) - \dim \mathbf{pr}_{F^{\natural}}^{\text{lin}}(\mathcal{Z}^{\natural}) \geq \dim(F^{\natural})^{\text{lin}}$.

We will explain part (2) in the next subsection of the Introduction (below Definition 1.7). For the moment let me just point out why both parts of Conjecture 1.4 are needed:

- Conjecture 1.4 is a natural generalization of the Ax-Schanuel theorem [2] (see [37, Introduction]). Ax' theorem, concerning $\mathbb{C}^n \rightarrow (\mathbb{C}^*)^n$, implies both Ax logarithmique and Ax-Lindemann for this bi-algebraic system, and describes the Zariski closure of complex analytic subvarieties of the graph. This is still expected to hold for mixed Shimura varieties (Conjecture 8.1). If we want to make an Ax-Schanuel conjecture for enlarged mixed Shimura varieties, it should then reflect both aspects. Note that for mixed Shimura varieties, part (2) already implies part (1).
- Part (1) of Conjecture 1.4 implies both Ax logarithmique (Theorem 1.2) and Ax-Lindemann (Theorem 1.3), while part (2) does not.
- Part (2) of Conjecture 1.4 describes the Zariski closure of $\mathcal{Z}^{\natural}$ in the ambient space $\mathcal{X}^{\natural+} \times S^{\natural}$, while part (1) does not.

We will prove the following cases for Conjecture 1.4.

Theorem 1.5. *Conjecture 1.4 holds in the following cases:*

- (1) *When $\tilde{Z}^{\natural}$ is algebraic (equivalent to Ax-Lindemann by Theorem 8.5).*
- (2) *When $Z^{\natural}$ is algebraic (equivalent to Ax logarithmique by Theorem 8.4).*
- (3) *When $Z^{\natural}$ is contained in a fiber of $S^{\natural} \rightarrow S_G$ under the notation of (1.1) (Theorem 9.1).*

Little is known for Conjecture 1.4 beyond Ax-Lindemann and Ax logarithmique: Ax [3] proved part (2) for the universal vector extension of any semi-abelian variety^[2], and Pila-Tsimerman [32] proved the conjecture for $Y(1)^n$ (one crucial point is that each simple factor of the group attached to $Y(1)^n$, which is $\text{SL}_2(\mathbb{Q})$, is small). Theorem 1.5(3) completely solves the problem

^[2]Ax' theorem was about any complex algebraic group with its Lie algebra, but this is easily reduced to the case of abelian groups with their Lie algebras, and hence is equivalent to the statement for the universal vector extension of any semi-abelian variety.

for the unipotent part, and is the first result for this conjecture when the ambient variety does not admit a natural group structure (take for example the fiber of $\mathfrak{P}_g^\natural \rightarrow \mathcal{A}_g$) and that the algebraic group attached to its uniformization is large. It is also the first result for part (1) of Conjecture 1.4 when part (1) is not implied by part (2).

1.6. Quasi-linear subvarieties and geometric bi-algebraicity. Notation: for any abelian scheme $\mathfrak{A} \rightarrow B$ with unit section ε , denote by $\omega_{\mathfrak{A}/B} := \varepsilon^* \Omega_{\mathfrak{A}/B}^1$.

A crucial ingredient for Conjecture 1.4 is to understand the geometric bi-algebraic system associated with $S^\natural$. This is the **third main goal** of this paper. We briefly explain this in this subsection. We say that an irreducible subvariety $Y^\natural$ of $S^\natural$ is **geometrically bi-algebraic** if one (and hence all) complex analytic irreducible component of $(\text{unif}^\natural)^{-1}(Y^\natural)$ is algebraizable, i.e. its dimension equals the dimension of its Zariski closure in $\mathcal{X}^{\natural, \vee}$. As we pointed out, the first step to prove any Ax type theorem is to establish the characterization of geometrically bi-algebraic subvarieties of $S^\natural$. The result cannot be as elegant as for mixed Shimura varieties. Take for example $S^\natural = \mathfrak{A}_g^\natural$. Any algebraic subvariety $Y^\natural$ contained in a fiber of $\mathfrak{A}_g^\natural \rightarrow \mathcal{A}_g$ is geometrically bi-algebraic. However we prove that this is the only problem: for the exact sequence $0 \rightarrow \omega_{\mathfrak{A}_g/\mathcal{A}_g} \rightarrow \mathfrak{A}_g^\natural \rightarrow \mathfrak{A}_g \rightarrow 0$ of groups over $\mathcal{A}_g$, the “non-linear” part of any geometrically bi-algebraic subvariety $Y^\natural$ of $\mathfrak{A}_g^\natural$ can only lie in the TRIVIAL subbundle of the vector bundle part, i.e. of $\omega_{\mathfrak{A}_g/\mathcal{A}_g}|_{Y_G}$ where Y_G is the image of $Y^\natural$ in $\mathcal{A}_g$. More precisely we prove (Theorem 5.8):

Theorem 1.6 (Characterization of geometrically bi-algebraic subvarieties of enlarged mixed Shimura varieties). *An irreducible subvariety $Y^\natural$ of $S^\natural$ is geometrically bi-algebraic if and only if it is quasi-linear.*

We define quasi-linear subvarieties of $S^\natural$. We shall see in §3.4 that there is a commutative diagram for any connected enlarged mixed Shimura variety (on the right for $(\mathfrak{P}_g^\natural)^{[n]}$):

$$(1.1) \quad \begin{array}{ccc} S^\natural & \xrightarrow{[\pi^\natural]} & S \\ \downarrow [\pi_{P/U}^\natural] \lrcorner & & \downarrow [\pi_{P/U}] \\ S_{P/U}^\natural & \xrightarrow{[\pi_{P/U}^\natural]} & S_{P/U} \xrightarrow{[\pi_{P/U}]} S_G \end{array} \quad \begin{array}{ccc} (\mathfrak{P}_g^\natural)^{[n]} & \xrightarrow{\quad} & \mathfrak{P}_g^{[n]} \\ \downarrow \lrcorner & & \downarrow \\ \mathfrak{A}_g^\natural \times (\mathfrak{A}_g^\vee)^\natural & \xrightarrow{\quad} & \mathfrak{A}_g \times \mathfrak{A}_g^\vee \xrightarrow{\quad} \mathcal{A}_g \end{array}$$

where

- all maps in the diagram are projections defined in some natural way;
- S is a connected mixed Shimura variety, and $S_{P/U}$ is the quotient of S by its weight -2 part (and hence $S_{P/U}$ is an abelian scheme over S_G);
- $S_{P/U}^\natural$ is the universal vector extension of the abelian scheme $S_{P/U} \rightarrow S_G$.

To define quasi-linear subvarieties of $S^\natural$ we need some preparation. Let Y_G be a subvariety of S_G . Let $Y_{P/U} \subset S_{P/U}|_{Y_G} := [\pi_{P/U}]^{-1}(Y_G)$ be the translate of an abelian subscheme of $S_{P/U}|_{Y_G} \rightarrow Y_G$ by a torsion section and then by a constant section of its isotrivial part. Denote by $Y_{P/U}^{\text{univ}}$ the universal vector extension of the abelian scheme $Y_{P/U} \rightarrow Y_G$. Then by the rigidity of the universal vector extension (see Remark 3.5) there is a unique embedding

$Y_{P/U}^{\text{univ}} \subset S_{P/U}^{\natural}|_{Y_{P/U}} := [\pi_{P/U}^{\natural}]^{-1}(Y_{P/U})$ compatible with the embedding $Y_{P/U} \subset S_{P/U}|_{Y_G}$ mentioned above. More concretely there is a unique embedding i (left vertical arrow) of vector groups over Y_G inducing the following push-out:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \omega_{Y_{P/U}^{\vee}/Y_G} & \longrightarrow & Y_{P/U}^{\text{univ}} & \longrightarrow & Y_{P/U} \longrightarrow 0 \\ & & \downarrow i & & \downarrow & & \downarrow = \\ 0 & \longrightarrow & \omega_{[\pi_G]^{-1}(Y_G)^{\vee}/Y_G} & \longrightarrow & S_{P/U}^{\natural}|_{Y_{P/U}} & \longrightarrow & Y_{P/U} \longrightarrow 0 \end{array}$$

Then we have

$$S_{P/U}^{\natural}|_{Y_{P/U}} = Y_{P/U}^{\text{univ}} \times_{Y_G} \frac{\omega_{[\pi_G]^{-1}(Y_G)^{\vee}/Y_G}}{\omega_{Y_{P/U}^{\vee}/Y_G}}$$

Denote by $\mathbf{V}^{(0)}|_{Y_G}$ the largest trivial subbundle of $\frac{\omega_{[\pi_G]^{-1}(Y_G)^{\vee}/Y_G}}{\omega_{Y_{P/U}^{\vee}/Y_G}}$. We are ready to define

Definition 1.7. An irreducible subvariety $Y^{\natural}$ of $S^{\natural}$ is called **quasi-linear** if the followings hold: under the following notations for $Y^{\natural}$ compatible with (1.1)

$$\begin{array}{ccccc} Y^{\natural} & \xrightarrow{\quad} & Y & & \\ \downarrow & & \downarrow & \searrow & \\ Y_{P/U}^{\natural} & \xrightarrow{\quad} & Y_{P/U} & \xrightarrow{\quad} & Y_G \end{array}$$

- (1) Y is a weakly special subvariety of S . In particular $Y_{P/U}$ is the translate of an abelian subscheme of $S_{P/U}|_{Y_G} \rightarrow Y_G$ by a torsion section and then by a constant section of its isotrivial part.
- (2) Under the notations above the theorem, $Y_{P/U}^{\natural} = Y_{P/U}^{\text{univ}} \times_{Y_G} (L^{\natural} \times_{Y_G}) \times_{Y_G} \mathbf{V}^{\dagger}|_{Y_G}$, where $L^{\natural}$ is an irreducible algebraic subvariety of any fiber of $\mathbf{V}^{(0)}|_{Y_G} \rightarrow Y_G$, and $\mathbf{V}^{\dagger}|_{Y_G}$ is a subbundle of $\frac{\omega_{[\pi_G]^{-1}(Y_G)^{\vee}/Y_G}}{\omega_{Y_{P/U}^{\vee}/Y_G}}$ whose intersection with $\mathbf{V}^{(0)}|_{Y_G}$ is trivial.
- (3) $Y^{\natural} = Y \times_{Y_{P/U}} Y_{P/U}^{\natural}$ for the cartesian diagram in (1.1).

Now we are ready to explain the terminology in part (2) of Conjecture 1.4. Apply Theorem 1.6 to the bi-algebraic subvariety $F^{\natural}$ of $S^{\natural}$ (hence we change every letter “ Y ” by “ F ”), then we define

$$(F^{\natural})^{\text{lin}} := F \times_{F_{P/U}} (F_{P/U}^{\text{univ}} \times_{F_G} \mathbf{V}^{\dagger}|_{F_G})$$

and $[pr]_{F^{\natural}}^{\text{lin}}$ the natural projection $F^{\natural} \rightarrow (F^{\natural})^{\text{lin}}$. Let $pr_{\tilde{F}^{\natural}}^{\text{lin}}$ be the natural projection from $\tilde{F}^{\natural}$ to $(\tilde{F}^{\natural})^{\text{lin}}$, the uniformization of $(F^{\natural})^{\text{lin}}$. Then we define

$$\mathbf{pr}_{F^{\natural}}^{\text{lin}} := (pr_{\tilde{F}^{\natural}}^{\text{lin}}, [pr]_{F^{\natural}}^{\text{lin}}): \tilde{F}^{\natural} \times F^{\natural} \rightarrow (\tilde{F}^{\natural})^{\text{lin}} \times (F^{\natural})^{\text{lin}}.$$

Remark 1.8. For $Y^{\natural}$ quasi-linear, if we let X_G be the smallest pure Shimura subvariety of S_G which contains Y_G , then the abelian scheme $Y_{P/U} \rightarrow Y_G$ extends to an abelian scheme $X_{P/U} \rightarrow X_G$. The vector bundle $\mathbf{V}^{\text{extr}}|_{Y_G}$ is then the restriction to Y_G of the automorphic bundle $\frac{\omega_{[\pi_G]^{-1}(X_G)^{\vee}/X_G}}{\omega_{X_{P/U}^{\vee}/X_G}}$ over X_G , and $\mathbf{V}^{(0)}|_{Y_G}$ is the restriction of the largest trivial automorphic

subbundle $\mathbf{V}^{(0)}$ of $\frac{\omega_{[\pi_G]^{-1}(Y_G)^\vee/Y_G}}{\omega_{Y_P^\vee/U/Y_G}}$ to Y_G . The vector bundle $\mathbf{V}^\dagger|_{Y_G}$ is the restriction to Y_G of an automorphic subbundle $\mathbf{V}^\dagger$ over X_G whose intersection with $\mathbf{V}^{(0)}$ is trivial.

1.7. André-Oort and Zilber-Pink type problems for enlarged mixed Shimura varieties. In §10, we briefly discuss about special points and special subvarieties of enlarged mixed Shimura varieties. We will discuss Conjecture 1.1.

Structure of the paper. We review Pink’s work on equivariant families of mixed Hodge structures in §2. This is the foundation of the Deligne-Pink language for enlarged mixed Shimura varieties. Then we define and prove basic properties of enlarged mixed Shimura varieties in §3 (for enlarged mixed Shimura data) and §4 (for enlarged mixed Shimura varieties). In particular we study their relationship with the mixed Shimura varieties, explaining both the categorical comparison and the geometrical comparison. We also prove that any enlarged mixed Shimura variety, a priori defined as a complex analytic space, is algebraic and can be canonically descent to its reflex field. §5 discusses about the geometric bi-algebraicity and proves the characterization of geometrically bi-algebraic subvarieties of enlarged mixed Shimura varieties. Then we study Ax type transcendental results for enlarged mixed Shimura varieties in the next sections. We prove Ax logarithmique in §6 and give the equivalent statements for Ax-Lindemann in §7. The proof of Ax-Lindemann for enlarged mixed Shimura varieties, which is similar to the author’s proof of Ax-Lindemann for mixed Shimura varieties in [16], is executed in the Appendix. In §8 we formulate the Ax-Schanuel conjecture, explain its meaning and prove its relation with Ax logarithmique and with Ax-Lindemann. We prove the Ax-Schanuel conjecture for the unipotent part, i.e. the fiber from any enlarged mixed Shimura variety to its pure part, in §9. Finally we make a small discussion about André-Oort and Zilber-Pink type problems in §10.

Acknowledgements. The paper originated from a discussion with Daniel Bertrand. I would like to thank him for the inspiration. I would also like to thank him for teaching me the theory of universal vector extensions and his suggestions to improve the writing. I would like to thank Gisbert Wüstholz for our discussion on the arithmetic bi-algebraicity. I have benefited from discussions with Richard Pink on the first part of the paper, especially his suggestion to simplify the proof of Theorem 4.4. I would like to thank him for the interest in the paper and his help. I would like to thank Jacob Tsimerman for his comments and suggestions to improve the presentation of the introduction. Part of the work was done when the author was at the IAS (NJ, USA) and the author would like to thank its hospitality. The author was supported by NSF grant DMS-1128155.

Convention. For any abelian scheme $\mathfrak{A} \rightarrow B$ with unit section ε , denote by $\omega_{\mathfrak{A}/B} := \varepsilon^* \Omega_{\mathfrak{A}/B}^1$.

Since we will only talk about geometric bi-algebraicity in the main body of the paper, we abbreviate “geometrically bi-algebraic” to “bi-algebraic”.

When we say “definable”, we mean definable in the o-minimal theory $\mathbb{R}_{\text{an}, \text{exp}}$.

2. PINK’S WORK ON EQUIVARIANT FAMILIES OF MIXED HODGE STRUCTURES

In this section we review Pink’s work on equivariant families of mixed Hodge structures. The reference of the whole section is [33, Chapter 1].

2.1. Mixed Hodge structure. In this subsection we recall some background knowledge about rational mixed Hodge structures. In this subsection, the ring $R = \mathbb{Z}$ or $\mathbb{Q}$.

2.1.1. Basic facts about mixed Hodge structures. We start by collecting some basic notions about Hodge structures.

Let M be a free R -module of finite rank. A **pure Hodge structure of weight** $n \in \mathbb{Z}$ on M is a decomposition $M_{\mathbb{C}} = \bigoplus_{p+q=n} M^{p,q}$ into $\mathbb{C}$ -vector spaces such that for all $p, q \in \mathbb{Z}$ with $p+q = n$ one has $\overline{M^{q,p}} = M^{p,q}$. The associated **Hodge filtration** on $M_{\mathbb{C}}$ is defined by $F^p M_{\mathbb{C}} := \bigoplus_{p' \geq p} M^{p',q}$. It determines the Hodge structure uniquely, because $M^{p,q} = F^p M_{\mathbb{C}} \cap \overline{F^q M_{\mathbb{C}}}$.

A **mixed R -Hodge structure** on M is a triple $(M, \{W_n M\}_{n \in \mathbb{Z}}, \{F^p M_{\mathbb{C}}\}_{p \in \mathbb{Z}})$ consisting of an ascending exhausting separated filtration $\{W_n M\}_{n \in \mathbb{Z}}$ of M by R -modules of finite rank with each $M/W_n M$ free, called **weight filtration**, together with a descending exhausting separated filtration $\{F^p M_{\mathbb{C}}\}_{p \in \mathbb{Z}}$ of $M_{\mathbb{C}}$, called **Hodge filtration**, such that the Hodge filtration induces a pure Hodge structure of weight n on $\text{Gr}_n^W M := W_n M / W_{n-1} M$ for all $n \in \mathbb{Z}$. A pure Hodge structure of weight n is then a special case of a mixed Hodge structure by defining the weight filtration as $W_{n'} M = M$ for $n' \geq n$ and $W_{n'} M = 0$ for $n' < n$. The notions of **weight** $\leq n$ and **weight** $\geq n$ are defined in the obvious way.

The **Hodge numbers** are $h^{p,q} := \dim_{\mathbb{C}}(\text{Gr}_{p+q}^W M)^{p,q}$. They satisfy $h^{q,p} = h^{p,q}$, almost all $h^{p,q}$ are zero, and $\sum h^{p,q} = \dim M$. If $A \subset \mathbb{Z} \oplus \mathbb{Z}$ is an arbitrary subset, then we say that the Hodge structure $(M, \{W_n M\}_{n \in \mathbb{Z}}, \{F^p M_{\mathbb{C}}\}_{p \in \mathbb{Z}})$ is **of type** A if $h^{p,q} = 0$ for all $(p, q) \notin A$.

A **morphism of mixed R -Hodge structures** is a homomorphism $f: M \rightarrow M'$ such that $f(W_n M) \subset W_n M'$ and $f(F^p M_{\mathbb{C}}) \subset F^p M'_{\mathbb{C}}$ for all $n, p \in \mathbb{Z}$. The rational mixed Hodge structures form an abelian category with these morphisms. Given mixed R -Hodge structures on M_1 and M_2 , there are canonical rational mixed Hodge structures on $M_1 \oplus M_2$, on the dual $M_1^{\vee}$ and on $\text{Hom}(M_1, M_2)$.

A mixed Hodge structure on M is said to **split over** $\mathbb{R}$ if there exists a decomposition $M_{\mathbb{C}} = \bigoplus_{p,q} M^{p,q}$ such that $W_n M_{\mathbb{C}} = \bigoplus_{p+q \leq n} M^{p,q}$, $F^p M_{\mathbb{C}} = \bigoplus_{p' \geq p} M^{p',q}$ and $\overline{M^{q,p}} = M^{p,q}$. This decomposition is then uniquely determined by these properties. Every pure Hodge structure splits over $\mathbb{R}$, but not every mixed Hodge structure does. However we still have (see [33, 1.2])

Proposition 2.1 (Deligne). *Fix a mixed R -Hodge structure on M .*

- (1) *There exists a decomposition $M_{\mathbb{C}} = \bigoplus_{p,q} M^{p,q}$ such that $W_n M_{\mathbb{C}} = \bigoplus_{p+q \leq n} M^{p,q}$ and $F^p M_{\mathbb{C}} = \bigoplus_{p' \geq p} M^{p',q}$.*
- (2) *The Hodge structure is uniquely determined by any such decomposition.*
- (3) *There exists a unique decomposition as in (1) which also satisfies*

$$\overline{M^{q,p}} \equiv M^{p,q} \pmod{\bigoplus_{p' < p, q' < q} M^{p',q'}}.$$

2.1.2. Deligne torus. Let $\mathbb{S} := \text{Res}_{\mathbb{C}/\mathbb{R}} \mathbb{G}_{m,\mathbb{C}}$. The torus $\mathbb{S}$ is called the **Deligne-torus**. Over $\mathbb{C}$ it is canonically isomorphic to $\mathbb{G}_{m,\mathbb{C}} \times \mathbb{G}_{m,\mathbb{C}}$, but the action of complex conjugation is twisted by the automorphism c that interchanges the two factors. In particular $\mathbb{S}(\mathbb{R}) = \mathbb{C}^* \subset \mathbb{S}(\mathbb{C}) = \mathbb{C}^* \times \mathbb{C}^*$ consists of the points $(z, \bar{z})$ with $z \in \mathbb{C}^*$. While the character group of $\mathbb{G}_{m,\mathbb{C}}$ is $\mathbb{Z}$ in the standard way, we identify the character group of $\mathbb{S}$ with $\mathbb{Z} \oplus \mathbb{Z}$ such that the character (p, q) maps $z \in \mathbb{S}(\mathbb{R}) = \mathbb{C}^*$ to $z^{-p} \bar{z}^{-q} \in \mathbb{C}^*$. Under this identification the complex conjugation operates on $\mathbb{Z} \oplus \mathbb{Z}$ by interchanging the two factors. The following homomorphisms are important:

- the weight $\omega: \mathbb{G}_{m,\mathbb{R}} \hookrightarrow \mathbb{S}$ induced by $\mathbb{R}^* \subset \mathbb{C}^*$;

- $\mu: \mathbb{G}_{m,\mathbb{C}} \rightarrow \mathbb{S}_{\mathbb{C}}$ sending $z \in \mathbb{C}^* \mapsto (z, 1) \in \mathbb{C}^* \times \mathbb{C}^* = \mathbb{S}(\mathbb{C})$;
- the norm $N: \mathbb{S} \rightarrow \mathbb{G}_{m,\mathbb{R}}$ sending $z \in \mathbb{S}(\mathbb{R}) = \mathbb{C}^* \mapsto z\bar{z} \in \mathbb{R}^*$. The kernel $\mathbb{S}^1$ of N is anisotropic over $\mathbb{R}$, and we have a short exact sequence $1 \rightarrow \mathbb{S}^1 \rightarrow \mathbb{S} \rightarrow \mathbb{G}_{m,\mathbb{R}} \rightarrow 1$.

Let M be a free R -module of finite rank. The choice of a representation $k: \mathbb{S}_{\mathbb{C}} \rightarrow \mathrm{GL}(M_{\mathbb{C}})$ is equivalent to the choice of a decomposition $M_{\mathbb{C}} = \bigoplus_{p,q} M^{p,q}$, where $M^{p,q}$ is the eigenspace in $M_{\mathbb{C}}$ to the character (p, q) . Define $W_n M_{\mathbb{C}} = \bigoplus_{p+q \leq n} M^{p,q}$ and $F^p M_{\mathbb{C}} = \bigoplus_{p' \geq p} M^{p',q}$. We want to understand when the triple $(M, \{W_n M_{\mathbb{C}}\}, \{F^p M_{\mathbb{C}}\})$ is a mixed R -Hodge structure on M (so in particular $W_n M_{\mathbb{C}}$ is defined over R for all n). The following two propositions of Pink will tell us under which condition on k this is the case for $R = \mathbb{Q}$.

Proposition 2.2. ([33, 1.4]) *Let P be a connected $\mathbb{Q}$ -linear algebraic group. Let $W := \mathcal{R}_u(P)$ be its unipotent radical, let $G := P/W$ and let $\pi: P \rightarrow G$ be the quotient map. Let $h: \mathbb{S}_{\mathbb{C}} \rightarrow P_{\mathbb{C}}$ be a homomorphism such that the following conditions hold:*

- $\pi \circ h: \mathbb{S}_{\mathbb{C}} \rightarrow G_{\mathbb{C}}$ is defined over $\mathbb{R}$;
- $\pi \circ h \circ \omega: \mathbb{G}_{m,\mathbb{R}} \rightarrow G_{\mathbb{R}}$ is a cocharacter of the center of G defined over $\mathbb{Q}$;
- Under the weight filtration on $(\mathrm{Lie}P)_{\mathbb{C}}$ defined by $\mathrm{Ad}_P \circ h$ we have $W_{-1}(\mathrm{Lie}P) = \mathrm{Lie}W$.

Then

- (1) For every $(\mathbb{Q})$ -representation $\rho: P \rightarrow \mathrm{GL}(M)$, the homomorphism $\rho \circ h: \mathbb{S}_{\mathbb{C}} \rightarrow \mathrm{GL}(M_{\mathbb{C}})$ induces a rational mixed Hodge structure on M .
- (2) The weight filtration on M is stable under P .
- (3) For any $p \in P(\mathbb{R})W(\mathbb{C})$, the assertions (1) and (2) also hold for $\mathrm{int}(p) \circ h$ in place of h . The weight filtration and the Hodge numbers in any representation are the same for $\mathrm{int}(p) \circ h$ and for h .

Proposition 2.3. *Let M be a finite dimensional $\mathbb{Q}$ -vector space. A representation $k: \mathbb{S}_{\mathbb{C}} \rightarrow \mathrm{GL}(M_{\mathbb{C}})$ defines a rational mixed Hodge structure on M iff there exist a connected $\mathbb{Q}$ -linear algebraic group P , a representation $\rho: P \rightarrow \mathrm{GL}(M)$ and a homomorphism $h: \mathbb{S}_{\mathbb{C}} \rightarrow P_{\mathbb{C}}$ such that $k = \rho \circ h$ and the conditions in Proposition 2.2 are satisfied. Moreover, every rational mixed Hodge structure on M is obtained by a unique representation $k: \mathbb{S}_{\mathbb{C}} \rightarrow \mathrm{GL}(M_{\mathbb{C}})$ with the property above.*

Proof. This is [33, 1.5] except the “Moreover” part, where the existence of k has been explained in the paragraph before Proposition 2.2 and the uniqueness of k follows from Proposition 2.1(3). $\square$

2.1.3. Mumford-Tate group and polarizations. Let M be a free R -module of finite rank equipped with a mixed R -Hodge structure $(M, \{W_n M\}_{n \in \mathbb{Z}}, \{F^p M_{\mathbb{C}}\}_{p \in \mathbb{Z}})$. By Proposition 2.3, the corresponding rational mixed Hodge structure on $M_{\mathbb{Q}}$ gives rise to a representation $k: \mathbb{S}_{\mathbb{C}} \rightarrow \mathrm{GL}(M_{\mathbb{C}})$.

Definition 2.4. *The **Mumford-Tate group** of this mixed R -Hodge structure is defined to be the smallest $\mathbb{Q}$ -subgroup P of $\mathrm{GL}(M_{\mathbb{Q}})$ such that $k(\mathbb{S}_{\mathbb{C}}) \subset P_{\mathbb{C}}$.*

Before defining the polarizations of pure Hodge structures, we introduce the **Tate Hodge structure**, which is defined to be the free R -module of rank 1 $R(1) := 2\pi\sqrt{-1}R$ with the pure R -Hodge structure of type $(-1, -1)$. For every $n \in \mathbb{Z}$, we get a pure R -Hodge structure of type $(-n, -n)$ on $R(n) := R(1)^{\otimes n}$.

Definition 2.5. Suppose that the R -Hodge structure on M is pure of weight n . A **polarization** of this Hodge structure is a homomorphism of Hodge structures

$$Q: M \otimes M \rightarrow R(-n)$$

which is $(-1)^n$ -symmetric and such that the real-valued symmetric bilinear form $Q'(u, v) := (2\pi\sqrt{-1})^n Q(Cu, v)$ is positive-definite on $M_{\mathbb{R}}$, where C acts on $M^{p,q}$ by $C|_{M^{p,q}} = (\sqrt{-1})^{p-q}$.

2.1.4. Variation of mixed Hodge structures.

Definition 2.6. ([27, Definition 14.44]) Let S be a complex manifold. A **variation of mixed R -Hodge structures over S** is a triple $(\mathbb{V}, W, \mathcal{F})$ with

- (1) a local system $\mathbb{V}$ of free R -modules of finite rank on S ;
- (2) a finite increasing filtration $\{W_m\}$ of the local system $\mathbb{V}$ by local sub-systems with torsion free $\text{Gr}_n^W \mathbb{V}$ for each n (this is called the weight filtration);
- (3) a finite decreasing filtration $\{\mathcal{F}^p\}$ of the holomorphic vector bundle $\mathcal{V} := \mathbb{V} \otimes_{R_S} \mathcal{O}_S$, where R_S is the constant sheaf over S , by holomorphic subbundles (this is called the Hodge filtration).

such that

- (1) for each $s \in S$, the filtrations $\{\mathcal{F}^p(s)\}$ and $\{W_m\}$ of $\mathbb{V}(s) \simeq \mathbb{V}_s \otimes_R \mathbb{C}$ define a mixed Hodge structure on the R -module of finite rank $\mathbb{V}_s$;
- (2) the connection $\nabla: \mathcal{V} \rightarrow \mathcal{V} \otimes_{\mathcal{O}_S} \Omega_S^1$ whose sheaf of horizontal sections is $\mathbb{V}_{\mathbb{C}}$ satisfies the Griffiths' transversality condition

$$\nabla(\mathcal{F}^p) \subset \mathcal{F}^{p-1} \otimes \Omega_S^1.$$

Definition 2.7. A variation of mixed Hodge structures over S is said to be **graded-polarizable** if the induced variations of pure Hodge structure $\text{Gr}_n^W \mathbb{V}$ are all polarizable, i.e. for each n , there exists a flat morphism of variations

$$Q_n: \text{Gr}_n^W \mathbb{V} \otimes \text{Gr}_n^W \mathbb{V} \rightarrow R(-n)_S$$

which induces on each fibre a polarization of the corresponding Hodge structure of weight n .

2.2. Equivariant families of Hodge structures. Now we are ready to discuss equivariant families of Hodge structures, or more precisely homogeneous spaces parametrizing certain rational mixed Hodge structures.

Proposition 2.8 (Pink [33, 1.7]). Let P be a connected $\mathbb{Q}$ -linear group and let $W := \mathcal{R}_u(P)$ be its unipotent radical. Let $\mathcal{D}^{\natural}$ be a $P(\mathbb{R})W(\mathbb{C})$ -conjugacy class in $\text{Hom}(\mathbb{S}_{\mathbb{C}}, P_{\mathbb{C}})$. Assume that for one (and hence for all by Proposition 2.2(3)) $h \in \mathcal{D}^{\natural}$, the conditions in Proposition 2.2 holds. Let M be any faithful representation of P and let φ be the obvious map

$$\varphi: \mathcal{D}^{\natural} \rightarrow \{\text{rational mixed Hodge structures on } M\}$$

given by Proposition 2.2(1). Then:

- (1) There exists a unique structure on $\varphi(\mathcal{D}^{\natural})$ as a complex manifold such that the Hodge filtration on $M_{\mathbb{C}}$ depends analytically on $\varphi(h) \in \varphi(\mathcal{D}^{\natural})$. This structure is $P(\mathbb{R})W(\mathbb{C})$ -invariant and $W(\mathbb{C})$ acts analytically on $\varphi(\mathcal{D}^{\natural})$.

(2) For any other representation M' of P the analogous map

$$\varphi': \mathcal{D}^\natural \rightarrow \{\text{rational mixed Hodge structures on } M'\}$$

factors through $\varphi(\mathcal{D}^\natural)$. The Hodge filtration on M' varies analytically with $\varphi(h) \in \varphi(\mathcal{D}^\natural)$.

(3) If in addition M' is faithful, then $\varphi(\mathcal{D}^\natural)$ and $\varphi'(\mathcal{D}^\natural)$ are canonically isomorphic and the isomorphism is compatible with the complex structure.

Remark 2.9. By the proof of [33, 1.7], the map φ factors through

$$\mathcal{D}^\natural \simeq P(\mathbb{R})W(\mathbb{C})/\text{Cent}_{P(\mathbb{R})W(\mathbb{C})}(h) \rightarrow P(\mathbb{C})/F_h^0 P_{\mathbb{C}} \hookrightarrow \text{Grass}(M)(\mathbb{C}),$$

where the last map is a closed embedding.

The following lemma will be useful:

Lemma 2.10 (Pink [33, 1.8]). *Let P , $\mathcal{D}^\natural$, M and φ be as in Proposition 2.8. Then for any $h \in \mathcal{D}^\natural$, the projection $\pi: P \rightarrow G := P/\mathcal{R}_u(P)$ induces an isomorphism*

$$\text{Cent}_{P(\mathbb{R})W(\mathbb{C})}(h) \xrightarrow{\sim} \text{Cent}_{G(\mathbb{R})}(\pi \circ h).$$

Now two natural questions about this equivariant family of Hodge structures arise: under which condition we get a variation of rational Hodge structure on M over $\varphi(\mathcal{D}^\natural)$, and is there a subset $\mathcal{D}$ of $\mathcal{D}^\natural$ having the same image under φ such that $\varphi|_{\mathcal{D}}$ is finite. Both questions are answered by Pink. In the following two propositions, we let P , $\mathcal{D}^\natural$, M and φ be as in Proposition 2.8.

Proposition 2.11 (Pink [33, 1.10]). *We have a variation of rational mixed Hodge structures on M over $\varphi(\mathcal{D}^\natural)$ iff for one (and hence for all) $h \in \mathcal{D}^\natural$ the Hodge structure on $\text{Lie}P$ is of type*

$$\{(-1, 1), (0, 0), (1, -1), (-1, 0), (0, -1), (-1, -1)\}.$$

Proposition 2.12 (Pink [33, 1.16]). *Let $U < W$ be the unique connected subgroup such that $\text{Lie}U = W_{-2}(\text{Lie}W)$ (by Proposition 2.2(3), it does not depend on $h \in \mathcal{D}^\natural$). Let π' be the quotient $P \rightarrow P/U$. Let*

$$\mathcal{D} := \{h \in \mathcal{D}^\natural \mid \pi' \circ h: \mathbb{S}_{\mathbb{C}} \rightarrow (P/U)_{\mathbb{C}} \text{ is defined over } \mathbb{R}\}.$$

Then

- (1) $\mathcal{D}$ is a non-empty $P(\mathbb{R})U(\mathbb{C})$ -orbit in $\text{Hom}(\mathbb{S}_{\mathbb{C}}, P_{\mathbb{C}})$;
- (2) $\varphi(\mathcal{D}) = \varphi(\mathcal{D}^\natural)$;
- (3) If $F^0(\text{Lie}U)_{\mathbb{C}} = 0$, then $\varphi(\mathcal{D}) \simeq \mathcal{D}$.

Before proceeding to the next section, we make the following remark: in the study of mixed Shimura varieties one often identifies $\varphi(\mathcal{D})$ and $\mathcal{D}$ by Proposition 2.12(3), however to study enlarged mixed Shimura varieties it is important to distinguish $\varphi(\mathcal{D})$ and $\mathcal{D}$. We will be careful about this in the whole paper.

3. ENLARGED MIXED SHIMURA DATA

Definition 3.1. *An enlarged mixed Shimura datum $(P, \mathcal{X}^\natural, h)$ is a triple where*

- P is a connected linear algebraic group over $\mathbb{Q}$ with unipotent radical W and with another algebraic subgroup $U \subset W$ which is normal in P and uniquely determined by $\mathcal{X}^\natural$ using condition (3) below;

- $\mathcal{X}^\natural$ is a left homogeneous space under the subgroup $P(\mathbb{R})W(\mathbb{C}) \subset P(\mathbb{C})$, and $\mathcal{X}^\natural \xrightarrow{h} \text{Hom}(\mathbb{S}_{\mathbb{C}}, P_{\mathbb{C}})$ is a $P(\mathbb{R})W(\mathbb{C})$ -equivariant map^[3] such that every fibre of h consists of at most finitely many points,

such that for some (equivalently for all) $x \in \mathcal{X}^\natural$,

- (1) the composite homomorphism $\mathbb{S}_{\mathbb{C}} \xrightarrow{h_x} P_{\mathbb{C}} \rightarrow (P/W)_{\mathbb{C}}$ is defined over $\mathbb{R}$,
- (2) the adjoint representation induces on $\text{Lie}P$ a rational mixed Hodge structure of type

$$\{(-1, 1), (0, 0), (1, -1)\} \cup \{(-1, 0), (0, -1)\} \cup \{(-1, -1)\},$$

- (3) the weight filtration on $\text{Lie}P$ is given by

$$W_n(\text{Lie}P) = \begin{cases} 0 & \text{if } n < -2 \\ \text{Lie}U & \text{if } n = -2 \\ \text{Lie}W & \text{if } n = -1 \\ \text{Lie}P & \text{if } n \geq 0 \end{cases},$$

- (4) the conjugation by $h_x(\sqrt{-1})$ induces a Cartan involution on $G_{\mathbb{R}}^{\text{ad}}$ where $G := P/W$, and G^{ad} possesses no $\mathbb{Q}$ -factor H such that $H(\mathbb{R})$ is compact,
- (5) $P/P^{\text{der}} = Z(G)$ acts on U and on $V := W/U$ through a torus which is an almost direct product of a $\mathbb{Q}$ -split torus with a torus of compact type over $\mathbb{Q}$.

If in addition U is trivial, then $(P, \mathcal{X}^\natural, h)$ is said to be **of Kuga type**. For simplicity we mostly write $(P, \mathcal{X}^\natural)$ since we always consider exactly one map h for every pair $(P, \mathcal{X}^\natural)$.

Remark 3.2. (1) Conditions (2) and (3) together imply that the composite homomorphism $\mathbb{G}_{m, \mathbb{C}} \xrightarrow{\omega} \mathbb{S}_{\mathbb{C}} \xrightarrow{h_x} P_{\mathbb{C}} \rightarrow (P/W)_{\mathbb{C}}$ is a cocharacter of the center of P/W defined over $\mathbb{R}$. This map is called the weight.

- (2) We shall compare Definition 3.1 with that of mixed Shimura data in [33, 2.1] in §3.2 (categorical comparison) and §3.1 (geometric comparison). According to Pink [33, 2.1], a mixed Shimura datum is a triple $(P, \mathcal{X}, h)$ where P is as in Definition 3.1, $\mathcal{X}$ is a left homogeneous space under $P(\mathbb{R})U(\mathbb{C})$ and $h: \mathcal{X} \rightarrow \text{Hom}(\mathbb{S}_{\mathbb{C}}, P_{\mathbb{C}})$ is $P(\mathbb{R})U(\mathbb{C})$ -equivariant such that all the conditions in Definition 3.1 are satisfied for any $x \in \mathcal{X}$, with P/W replaced by P/U in condition (1). For those who are familiar with mixed Shimura data, the action of $P(\mathbb{R})U(\mathbb{C})$ on $\mathcal{X}$ can be extended to an action of $P(\mathbb{R})W(\mathbb{C})$.^[4] But the map $h: \mathcal{X} \rightarrow \text{Hom}(\mathbb{S}_{\mathbb{C}}, P_{\mathbb{C}})$ is NOT $P(\mathbb{R})W(\mathbb{C})$ -equivariant because $\text{Cent}_{P(\mathbb{R})W(\mathbb{C})}(h_x)$ is reductive by Lemma 2.10.
- (3) Any pure Shimura datum or any mixed Shimura datum with trivial weight -1 part is by definition an enlarged mixed Shimura datum.

In view of Remark 3.2(2) and Proposition 2.12, enlarged mixed Shimura data and mixed Shimura data encrypt the same information from the Hodge theory. Therefore in order to define morphisms between enlarged mixed Shimura data, other aspects should be taken into consideration. This is the geometric part which we consider in §3.1. For the moment we define:

^[3]See Remark 3.2(2) for some discussion.

^[4]Let M and φ be as in Proposition 2.8, then the action of $P(\mathbb{R})U(\mathbb{C})$ on $\varphi \circ h(\mathcal{X})$ extends to an action of $P(\mathbb{R})W(\mathbb{C})$ by Proposition 2.12. Hence the action of $P(\mathbb{R})U(\mathbb{C})$ on $\mathcal{X}$ extends to an action of $P(\mathbb{R})W(\mathbb{C})$ because $\varphi(h(\mathcal{X})) \simeq h(\mathcal{X})$ and every fiber of h is at most finite.

Definition 3.3. A *coarse morphism* $(P_1, \mathcal{X}_1^\natural, h_1) \rightarrow (P_2, \mathcal{X}_2^\natural, h_2)$ between enlarged mixed Shimura data consists of a homomorphism $f: P_1 \rightarrow P_2$ and a $P_1(\mathbb{R})W_1(\mathbb{C})$ -equivariant map $f_s: \mathcal{X}_1^\natural \rightarrow \mathcal{X}_2^\natural$ such that the following diagram commutes:

$$\begin{array}{ccc} \mathcal{X}_1^\natural & \xrightarrow{f_s} & \mathcal{X}_2^\natural \\ \downarrow h_1 & & \downarrow h_2 \\ \mathrm{Hom}(\mathbb{S}_{\mathbb{C}}, P_{1,\mathbb{C}}) & \xrightarrow{h \mapsto f \circ h} & \mathrm{Hom}(\mathbb{S}_{\mathbb{C}}, P_{2,\mathbb{C}}) \end{array}$$

3.1. Structure of the underlying space and the category of mixed Shimura data.

In this subsection we study the underlying space $\mathcal{X}^\natural$ of an enlarged mixed Shimura datum. In particular we will make a comparison between enlarged mixed Shimura data and mixed Shimura data. This comparison is geometric. In the end we define morphisms between enlarged mixed Shimura data and the category of enlarged mixed data (in contrast with the coarse one before).

Given an enlarged mixed Shimura datum $(P, \mathcal{X}^\natural, h)$, let $\mathcal{D}^\natural := h(\mathcal{X}^\natural)$ and let $\mathcal{D}$ be the subset of $\mathcal{D}^\natural$ defined in Proposition 2.12. Let M and φ be as in Proposition 2.8. Then $\varphi(\mathcal{D}^\natural) = \varphi(\mathcal{D}) \simeq \mathcal{D}$ by Proposition 2.12. Hence the triple $(P, \varphi(\mathcal{D}), \varphi^{-1})$ is a mixed Shimura datum. Consider the following projection map, which is the **geometric comparison** we will use:

$$(3.1) \quad \pi^\sharp: (P, \mathcal{X}^\natural, h) \rightarrow (P, \varphi(\mathcal{D}), \varphi^{-1}), \quad (p, x) \mapsto (p, \varphi(h_x)).$$

In other words, the map $\pi^\sharp$ is defined by the commutative diagram

$$\begin{array}{ccc} \mathcal{X}^\natural & \xrightarrow{\pi^\sharp} & \varphi(\mathcal{D}) \\ & \searrow h & \swarrow \varphi^{-1} \\ & \mathrm{Hom}(\mathbb{S}_{\mathbb{C}}, P_{\mathbb{C}}) & \end{array}$$

We write $(P, \mathcal{X}^\natural)$ and $(P, \varphi(\mathcal{D}))$ for simplicity. Let $\pi^\natural$ be the composition

$$(P, \mathcal{X}^\natural) \xrightarrow{\pi^\sharp} (P, \varphi(\mathcal{D})) \xrightarrow{\pi} (P, \varphi(\mathcal{D}))/W =: (G, \varphi(\mathcal{D})_G).$$

We have $\mathcal{D}^\natural = h(\mathcal{X}^\natural) \simeq P(\mathbb{R})W(\mathbb{C})/\mathrm{Cent}(h_x)$ for any $h_x \in \mathcal{D}^\natural$, and hence by Lemma 2.10 $\mathcal{D}^\natural$ is a $W(\mathbb{C})$ -torsor over the complex manifold $\varphi(\mathcal{D})_G$. Any Levi-decomposition $P = W \rtimes G$ gives rise to a section of this $W(\mathbb{C})$ -torsor, and hence $\mathcal{D}^\natural \simeq W(\mathbb{C}) \times \varphi(\mathcal{D})_G$ is a trivial $W(\mathbb{C})$ -torsor, which we can then see as a group over $\varphi(\mathcal{D})_G$. This endows $\mathcal{D}^\natural$ with a complex structure, invariant under $P(\mathbb{R})W(\mathbb{C})$. Since $h: \mathcal{X}^\natural \rightarrow \mathcal{D}^\natural$ is a local diffeomorphism, we have a complex structure on $\mathcal{X}^\natural$, invariant under $P(\mathbb{R})W(\mathbb{C})$. With this complex structure, the underlying map of every coarse morphism $(P_1, \mathcal{X}_1^\natural, h_1) \rightarrow (P_2, \mathcal{X}_2^\natural, h_2)$ on the underlying spaces $\mathcal{X}_1^\natural \rightarrow \mathcal{X}_2^\natural$ is holomorphic. Note that $\pi^\sharp: \mathcal{X}^\natural \rightarrow \varphi(\mathcal{D})$ is holomorphic and $P(\mathbb{R})W(\mathbb{C})$ -equivariant by Proposition 2.8(1).

Next we define a left $P(\mathbb{R})V(\mathbb{C})$ -homogeneous space $\mathcal{D}_{P/U}^\natural \subset \mathrm{Hom}(\mathbb{S}_{\mathbb{C}}, (P/U)_{\mathbb{C}})$ to be the image of $\mathcal{D}^\natural$ under the obvious morphism $\mathrm{Hom}(\mathbb{S}_{\mathbb{C}}, P_{\mathbb{C}}) \rightarrow \mathrm{Hom}(\mathbb{S}_{\mathbb{C}}, (P/U)_{\mathbb{C}})$. Then $(P/U, \mathcal{D}_{P/U}^\natural, \mathrm{id})$ is an enlarged mixed Shimura datum. Apply the geometric comparison discussed above to this enlarged mixed Shimura datum we get

$$\pi_{P/U}^\sharp: (P/U, \mathcal{D}_{P/U}^\natural) \rightarrow (P/U, \varphi(\mathcal{D})_{P/U}).$$

We give a better description of $\pi_{P/U}^\sharp$. The homologous description for $\pi^\sharp$ is given in §3.4. Now $V = \text{Lie}V$ is commutative, and hence gives rise to a $\mathbb{Q}$ -representation of G and furthermore a variation of rational Hodge structures over $\varphi(\mathcal{D})_G$ of weight -1 and type $\{(-1, 0), (0, -1)\}$. Let $\mathcal{V} := V(\mathbb{C}) \times \varphi(\mathcal{D})_G$ be a holomorphic bundle over $\varphi(\mathcal{D})_G$ and let $\mathcal{F}^0\mathcal{V}$ be the holomorphic subbundle of $\mathcal{V}$ which induces the Hodge filtration in each fiber. Then we have

$$(3.2) \quad \pi_{P/U}^\sharp: \mathcal{D}_{P/U}^\natural \rightarrow \varphi(\mathcal{D})_{P/U} \text{ is the natural projection } \mathcal{V} \rightarrow \mathcal{V}/\mathcal{F}^0\mathcal{V}.$$

Now we are ready to define morphisms of enlarged mixed Shimura data:

Definition 3.4. (1) A **morphism** $f: (P_1, \mathcal{X}_1^\natural, h_1) \rightarrow (P_2, \mathcal{X}_2^\natural, h_2)$ between enlarged mixed Shimura data is a coarse morphism such that the induced map $f_{P/U}: \mathcal{D}_{1,P/U}^\natural \rightarrow \mathcal{D}_{2,P/U}^\natural$ maps the zero section of $\mathcal{F}^0\mathcal{V}_1 \rightarrow \varphi(\mathcal{D}_1)_G$ into the zero section of $\mathcal{F}^0\mathcal{V}_2 \rightarrow \varphi(\mathcal{D}_2)_G$, where

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{F}^0\mathcal{V}_1 & \longrightarrow & \mathcal{D}_{1,P/U}^\natural = \mathcal{V}_1 & \longrightarrow & \varphi(\mathcal{D}_{1,P/U}) = \mathcal{V}_1/\mathcal{F}^0\mathcal{V}_1 \longrightarrow 0 & / \varphi(\mathcal{D}_1)_G \\ & & \downarrow & & \downarrow f_{P/U} & & \downarrow & \\ 0 & \longrightarrow & \mathcal{F}^0\mathcal{V}_2 & \longrightarrow & \mathcal{D}_{2,P/U}^\natural = \mathcal{V}_2 & \longrightarrow & \varphi(\mathcal{D}_{2,P/U}) = \mathcal{V}_2/\mathcal{F}^0\mathcal{V}_2 \longrightarrow 0 & / \varphi(\mathcal{D}_2)_G \end{array}$$

(2) The **category of enlarged mixed Shimura data** $\underline{\mathcal{EMSD}}$ is defined as follows: its objects are the collection of enlarged mixed Shimura data, and its morphisms are the collection of morphisms between enlarged mixed Shimura data.

Remark 3.5. The condition on the zero section for the definition of morphisms between enlarged mixed Shimura data is to respect the following rigidity of universal vector extensions of abelian schemes: let $\mathfrak{A}$ be an abelian scheme over B and let $i: \mathfrak{A}' \hookrightarrow \mathfrak{A}$ be an abelian subscheme, then there exists a unique embedding $i^\natural: (\mathfrak{A}')^\natural \hookrightarrow \mathfrak{A}^\natural$ respecting i , where $\mathfrak{A}^\natural$ (resp. $(\mathfrak{A}')^\natural$) is the universal vector extension of $\mathfrak{A}/B$ (resp. of $\mathfrak{A}'/B$). The uniqueness of $i^\natural$ is justified by

$$\begin{array}{ccccccc} 0 & \longrightarrow & \omega_{(\mathfrak{A}')^\vee/B} & \longrightarrow & (\mathfrak{A}')^\natural & \longrightarrow & \mathfrak{A}' \longrightarrow 0 \\ & & \downarrow & & \downarrow \text{ } \ulcorner & & \downarrow = \\ 0 & \longrightarrow & \omega_{\mathfrak{A}^\vee/B} & \longrightarrow & \mathfrak{A}^\natural|_{\mathfrak{A}'} & \longrightarrow & \mathfrak{A}' \longrightarrow 0 \end{array}$$

3.2. Some constructions. We start with a comparison lemma between enlarged mixed Shimura data and mixed Shimura data. This is a categorical comparison.

Lemma 3.6 (Categorical Comparison Lemma between $\underline{\mathcal{EMSD}}$ and MSD). *There exists an equivalence between $\underline{\mathcal{EMSD}}$ and the category of mixed Shimura data.*

Proof. Given an enlarged mixed Shimura datum $(P, \mathcal{X}^\natural, h)$, let $\mathcal{D}^\natural := h(\mathcal{X}^\natural)$ and let $\mathcal{D}$ be the subset of $\mathcal{D}^\natural$ defined in Proposition 2.12. Let $\mathcal{X} := h^{-1}(\mathcal{D}) \subset \mathcal{X}^\natural$. Then it is easy to check that $(P, \mathcal{X}, h)$ is a mixed Shimura datum. We say that $(P, \mathcal{X}, h)$ is the **mixed Shimura datum associated with** $(P, \mathcal{X}^\natural, h)$.

Conversely given a mixed Shimura datum $(P, \mathcal{X}, h)$, let $\mathcal{D} := h(\mathcal{X})$ and let $\mathcal{D}^\natural$ be the $P(\mathbb{R})W(\mathbb{C})$ -conjugacy class in $\text{Hom}(\mathbb{S}_{\mathbb{C}}, P_{\mathbb{C}})$ of any element of $\mathcal{D}$. Let M and φ be as in Proposition 2.8. Then we have $\varphi(\mathcal{D}^\natural) = \varphi(\mathcal{D}) \simeq \mathcal{D}$ by Proposition 2.12. Now take the fiber product

$\mathcal{X}^\natural := \mathcal{X} \times_{\varphi(\mathcal{D}^\natural)} \mathcal{D}^\natural$ and let $h^\natural$ be the projection of $\mathcal{X}^\natural$ to the second factor in the fiber product. Recall that φ is $P(\mathbb{R})W(\mathbb{C})$ -equivariant. Then it is not hard to check that $(P, \mathcal{X}^\natural, h^\natural)$ is an enlarged mixed Shimura datum. We say that $(P, \mathcal{X}^\natural, h^\natural)$ is the **enlarged mixed Shimura datum associated with** $(P, \mathcal{X}, h)$.

It is not hard to check that the constructions above are categorical, fully faithful and are inverse to each other. Hence we are done. $\square$

With this lemma and known results of mixed Shimura data, we can do some construction for enlarged mixed Shimura data.

Proposition 3.7 (Quotient). *Let $(P, \mathcal{X}^\natural)$ be an enlarged mixed Shimura datum and let P_0 be a normal subgroup of P . Then there exists a quotient enlarged mixed Shimura datum $(P, \mathcal{X}^\natural)/P_0$ and a morphism $(P, \mathcal{X}^\natural) \rightarrow (P, \mathcal{X}^\natural)/P_0$, unique up to isomorphism, such that every morphism $(P, \mathcal{X}^\natural) \rightarrow (Q, \mathcal{Y}^\natural)$, where the homomorphism $P \rightarrow Q$ factors through P/P_0 , factors in a unique way through $(P, \mathcal{X}^\natural)/P_0$. Moreover let $(P, \mathcal{X})$ be the mixed Shimura datum associated with $(P, \mathcal{X}^\natural)$, then $(P, \mathcal{X}^\natural)/P_0$ is the enlarged mixed Shimura datum associated with $(P, \mathcal{X})/P_0$.*

Proof. Pink [33, 2.9] proved the existence of the quotient mixed Shimura datum $(P, \mathcal{X})/P_0$. Then the proposition follows from the categorical comparison lemma (Lemma 3.6) above. $\square$

Proposition 3.8 (Unipotent extension). *Let $(P, \mathcal{X}^\natural)$ be an enlarged mixed Shimura datum and let $1 \rightarrow W_0 \rightarrow Q \rightarrow P \rightarrow 1$ be an extension of P by a unipotent group W_0 . Assume*

- *the adjoint action of Q on every abelian subquotient of W_0 factors through P ;*
- *every irreducible subquotient of $\text{Lie}W_0$ is of type $\{(-1, 0), (0, -1), (-1, -1)\}$ as a representation of $G := P/\mathcal{R}_u(P)$;*
- *the center of G acts on every irreducible subquotient of $\text{Lie}W_0$ through a torus that is an almost direct product of a $\mathbb{Q}$ -split torus with a torus of compact type over $\mathbb{Q}$.*

Then we have

- (1) *There exists an enlarged mixed Shimura datum $(Q, \mathcal{Y}^\natural)$ and a morphism $(Q, \mathcal{Y}^\natural) \rightarrow (P, \mathcal{X}^\natural)$, both unique up to isomorphism, extending the homomorphism $Q \rightarrow P$ such that $(Q, \mathcal{Y}^\natural)/W_0 \simeq (P, \mathcal{X}^\natural)$.*
- (2) *For every morphism $(P_1, \mathcal{X}_1^\natural) \rightarrow (P, \mathcal{X}^\natural)$ and every factorization $P_1 \rightarrow Q \rightarrow P$, there exists a unique extension $(P_1, \mathcal{X}_1^\natural) \rightarrow (Q, \mathcal{Y}^\natural) \rightarrow (P, \mathcal{X}^\natural)$.*
- (3) *Let $(P, \mathcal{X})$ be the mixed Shimura datum associated with $(P, \mathcal{X}^\natural)$ and let $(Q, \mathcal{Y})$ be the unipotent extension of $(P, \mathcal{X})$ by W_0 defined by Pink [33, 2.17]. Then $(Q, \mathcal{Y}^\natural)$ is the enlarged mixed Shimura datum associated with $(Q, \mathcal{Y})$.*

Proof. In fact Conclusion (3) gives the construction and the categorical comparison lemma (Lemma 3.6) proves that this is what we desire. $\square$

3.3. Structure of the underlying group. One direct corollary of (the proof of) Lemma 3.6 is that the underlying group remains the same for an enlarged mixed Shimura datum and its associated mixed Shimura datum. Therefore for any enlarged mixed Shimura datum, to study its underlying group it suffices to look at its associated mixed Shimura datum. Now by Pink [33, 2.15], we have the following result:

Given an enlarged mixed Shimura datum $(P, \mathcal{X}^\natural)$, we can associate to P a 4-tuple (G, V, U, Ψ) which is defined as follows:

- $G := P/\mathcal{R}_u(P)$ is the reductive part of P ;
- U is the normal subgroup of P as in Definition 3.1 (hence the weight -2 part) and $V := \mathcal{R}_u(P)/U$ (hence the weight -1 part). Both of them are vector groups with an action of G induced by conjugation in P (which factors through G for reason of weight);
- The commutator on $W := \mathcal{R}_u(P)$ induces a G -equivariant alternating form $\Psi: V \times V \rightarrow U$ by reason of weight. Moreover, Ψ is given by a polynomial with coefficients in $\mathbb{Q}$.

On the other hand, P is uniquely determined up to isomorphism by this 4-tuple:

- let W be the central extension of V by U defined by Ψ . More concretely, $W = U \times V$ as a $\mathbb{Q}$ -variety and the group law on W is $(u, v)(u', v') = (u + u' + \frac{1}{2}\Psi(v, v'), v + v')$;
- define the action of G on W by $g((u, v)) := (gu, gv)$;
- define $P := W \rtimes G$.

3.4. Notations for EMSD. We fix some notations for the rest of the paper. Let $(P, \mathcal{X}^\natural, h)$ be an enlarged mixed Shimura datum, which for simplicity is often denoted by $(P, \mathcal{X}^\natural)$.

notation	meaning
$(P, \mathcal{X})$	the associated mixed Shimura datum under Lemma 3.6
$(P/U, \mathcal{X}_{P/U})$	the quotient mixed Shimura datum $(P, \mathcal{X})/U$
$(P/U, \mathcal{X}_{P/U}^\natural)$	$(P, \mathcal{X}^\natural)/U$ = the enlarged mixed Shimura datum associated with $(P/U, \mathcal{X}_{P/U})$
G, W, U, V	as in §3.3
$\mathcal{D}^\natural$	$h(\mathcal{X}^\natural) \subset \text{Hom}(\mathbb{S}_{\mathbb{C}}, P_{\mathbb{C}})$
$\mathcal{D}$	the subset of $\mathcal{D}^\natural$ defined in Proposition 2.12
φ	the map $\mathcal{D}^\natural \rightarrow \{\text{rational mixed Hodge structures on } M\}$ for a faithful representation M of P as in Proposition 2.8
$(P/U, \varphi(\mathcal{D})_{P/U})$	the mixed Shimura datum $(P, \varphi(\mathcal{D}))/U$
$(G, \varphi(\mathcal{D})_G)$	the mixed Shimura datum $(P, \varphi(\mathcal{D}))/W$

The notations we introduced in §3.1 is summarized in the diagram

$$(3.3) \quad \begin{array}{ccc} (P, \mathcal{X}^\natural) & \xrightarrow{\pi^\sharp} & (P, \varphi(\mathcal{D})) \\ \pi_{P/U}^\natural \downarrow \lrcorner & & \downarrow \pi_{P/U} \\ (P/U, \mathcal{X}_{P/U}^\natural) & \xrightarrow{\pi_{P/U}^\sharp} & (P/U, \varphi(\mathcal{D})_{P/U}) \\ \pi_G^\natural \downarrow & & \downarrow \pi_G \\ (G, \varphi(\mathcal{D})_G) & \xrightarrow{\text{id}} & (G, \varphi(\mathcal{D})_G) \end{array} \quad \begin{array}{c} \pi \\ \pi \end{array}$$

We explain why the first square is a pullback. Pink proved in [33, 2.18, 2.19] that $\varphi(\mathcal{D})$ can be identified as a holomorphic complex vector bundle over $\varphi(\mathcal{D})_G$. In fact let $LW := \text{Lie}W_{\mathbb{C}} \times \varphi(\mathcal{D})_G$ and let $\mathcal{F}^0 LW$ be the holomorphic subbundle of LW whose fiber over x_G is $F_{x_G}^0 \text{Lie}W_{\mathbb{C}}$, then $\varphi(\mathcal{D}) = LW/\mathcal{F}^0 LW$. But $F_{x_G}^0 V_{\mathbb{C}} = F_{x_G}^0 \text{Lie}W_{\mathbb{C}} \simeq \exp(F_{x_G}^0 \text{Lie}W_{\mathbb{C}})$ as algebraic varieties and $\exp(F_{x_G}^0 \text{Lie}W_{\mathbb{C}})$ is a subgroup of $W(\mathbb{C})$ by reason of weight and type for every $x_G \in \varphi(\mathcal{D})_G$. So $\mathcal{F}^0 LW$ is a subgroup of $W(\mathbb{C}) \times \varphi(\mathcal{D})_G$ over $\varphi(\mathcal{D})_G$ and we denote by $\mathcal{F}^0 \mathcal{V} := \mathcal{F}^0 LW$. Therefore

$$(3.4) \quad \mathcal{F}^0 \mathcal{V} \subset \mathcal{D}^\natural = W(\mathbb{C}) \times \varphi(\mathcal{D})_G \xrightarrow{\pi^\sharp} \varphi(\mathcal{D}).$$

So the first square of the diagram above is a pull by (3.2).

Before ending this subsection, we remark that it is often enough to consider the case where $\mathcal{X}^\natural = \mathcal{D}^\natural$ (hence $\mathcal{X} = \mathcal{D}$) because of the following lemma:

Lemma 3.9. *For any enlarged mixed Shimura datum $(P, \mathcal{X}^\natural, h)$, the canonical morphism*

$$(P, \mathcal{X}^\natural) \rightarrow (P, \mathcal{X}^\natural)/P^{\text{der}} \times (P, h(\mathcal{X}^\natural))$$

is an embedding.

Proof. This follows from the categorical comparison Lemma 3.6 and the parallel result for mixed Shimura data [33, 2.11]. $\square$

3.5. Enlarged mixed Shimura data of Siegel type. In this subsection we focus on some important examples. Let $g \in \mathbb{N}_{>0}$. Let V_{2g} be a $\mathbb{Q}$ -vector space of dimension $2g$ and let

$$(3.5) \quad \Psi: V_{2g} \times V_{2g} \rightarrow U_{2g} := \mathbb{G}_{a, \mathbb{Q}}$$

be a non-degenerate alternating form. Define

$$\text{GSp}_{2g} := \{g \in \text{GL}(V_{2g}) \mid \Psi(gv, gv') = \nu(g)\Psi(v, v') \text{ for some } \nu(g) \in \mathbb{G}_m\},$$

and $\mathcal{H}_g \subset \text{Hom}(\mathbb{S}, \text{GSp}_{2g, \mathbb{R}})$ to be the set of all homomorphisms inducing a pure Hodge structure of type $\{(-1, 0), (0, -1)\}$ on V_{2g} and for which Ψ or $-\Psi$ defines a polarization. It is well-known that GSp_{2g} is a reductive group and $\text{GSp}_{2g}(\mathbb{R})$ acts transitively on $\mathcal{H}_g$.

3.5.1. We start with the enlarged mixed Shimura datum $(P_{2g, a}, \mathcal{X}_{2g, a}^\natural)^{[5]}$ which corresponds to the universal vector extension of the universal abelian variety (over a fine moduli space). See §4.2.1 for further details.

Let

$$\mathcal{X}_{2g, a} := V_{2g}(\mathbb{R}) \rtimes \mathcal{H}_g \subset \text{Hom}(\mathbb{S}, V_{2g, \mathbb{R}} \rtimes \text{GSp}_{2g, \mathbb{R}})$$

denote the conjugacy class under $V_{2g}(\mathbb{R}) \rtimes \text{GSp}_{2g}(\mathbb{R})$ generated by any element of $\mathcal{H}_g$. Then let

$$\mathcal{X}_{2g, a}^\natural := V_{2g}(\mathbb{C}) \rtimes \mathcal{H}_g \subset \text{Hom}(\mathbb{S}_{\mathbb{C}}, V_{2g, \mathbb{C}} \rtimes \text{GSp}_{2g, \mathbb{C}})$$

be the conjugacy class under $V_{2g}(\mathbb{C}) \rtimes \text{GSp}_{2g}(\mathbb{R})$ generated by any element of $\mathcal{X}_{2g, a}$. The notion $V_{2g}(R) \rtimes \mathcal{H}_g$ ($R = \mathbb{R}, \mathbb{C}$) is justified by the natural bijection

$$(3.6) \quad V_{2g}(R) \times \mathcal{H}_g \xrightarrow{\sim} V_{2g}(R) \rtimes \mathcal{H}_g, \quad (v', x) \mapsto \text{int}(v') \circ x.$$

Under this bijection the action of $(v, t) \in V_{2g}(\mathbb{C}) \rtimes \text{GSp}_{2g}(\mathbb{R})$ on $V_{2g}(\mathbb{C}) \rtimes \mathcal{H}_g$ is given by $(v, t) \cdot (v', x) := (v + tv', tx)$.

Denote by $(P_{2g, a}, \mathcal{X}_{2g, a}^\natural) := (V_{2g} \rtimes \text{GSp}_{2g}, V_{2g}(\mathbb{C}) \rtimes \mathcal{H}_g)$. Then $(P_{2g, a}, \mathcal{X}_{2g, a})$ is a mixed Shimura datum ([33, 2.25]) and $(P_{2g, a}, \mathcal{X}_{2g, a}^\natural)$ is the enlarged mixed Shimura datum associated with $(P_{2g, a}, \mathcal{X}_{2g, a})$.

Now let us turn to the geometric comparison considered in §3.1. Let M and φ be as in Proposition 2.8. In fact we can take M to be a $\mathbb{Q}$ -vector space of dimension $2g + 1$ because

$$P_{2g, a} = \begin{pmatrix} \text{GSp}_{2g} & V_{2g} \\ 0 & 1 \end{pmatrix} < \text{GL}_{2g+1}$$

[5]The index “a” is short for “abelian”.

and then φ factors through $\{\text{rational mixed Hodge structures on } M \text{ of type } \{(-1, 0), (0, -1), (0, 0)\}\}$. We shall work with the mixed Shimura datum $(P_{2g,a}, \varphi(\mathcal{X}_{2g,a}))$ instead of $(P_{2g,a}, \mathcal{X}_{2g,a})$. Now denote by $\mathcal{V}_{2g} := V_{2g}(\mathbb{C}) \times \mathcal{H}_g$ and let $\mathcal{F}^0 \mathcal{V}_{2g}$ be the holomorphic subbundle of $\mathcal{V}_{2g}$ such that its fiber over $x \in \mathcal{H}_g$ is $(V_{2g})_x^{0,-1}$. Then by (3.2), we have that on the underlying spaces

$$(3.7) \quad (P_{2g,a}, \mathcal{X}_{2g,a}^\natural) \xrightarrow{\pi^\natural} (P_{2g,a}, \varphi(\mathcal{X}_{2g,a})) \text{ is the natural projection } \mathcal{V}_{2g} \rightarrow \mathcal{V}_{2g}/\mathcal{F}^0 \mathcal{V}_{2g}.$$

3.5.2. Next we define the enlarged mixed Shimura datum $(P_{2g}, \mathcal{X}_{2g}^\natural)$, which corresponds to the pullback of the canonical relatively ample $\mathbb{G}_m$ -torsor over the universal abelian variety (over a fine moduli space) to its universal vector extension. See Example 4.5(1) for further details.

Let W_{2g} be the central extension of V_{2g} by U_{2g} defined by Ψ and let P_{2g} be the group associated to the 4-tuple $(\mathrm{GSp}_{2g}, V_{2g}, U_{2g}, \Psi)$ as in §3.3. Then $P_{2g} = W_{2g} \rtimes \mathrm{GSp}_{2g}$. The action of GSp_{2g} on W_{2g} induces a Hodge structure of type $\{(-1, 0), (0, -1), (-1, -1)\}$ on $\mathrm{Lie} W_{2g, \mathbb{C}}$. Let

$$\mathcal{X}_{2g} \subset \mathrm{Hom}(\mathbb{S}_{\mathbb{C}}, P_{2g, \mathbb{C}})$$

be the conjugacy class under $P_{2g}(\mathbb{R})U_{2g}(\mathbb{C})$ generated by any element of $\mathcal{H}_g$. Then $(P_{2g}, \mathcal{X}_{2g})$ defines a mixed Shimura datum (see [33, 2.25]). Pink [33, 10.10] proved that the mixed Shimura datum $(P_{2g}, \mathcal{X}_{2g})$ corresponds to the canonical ample $\mathbb{G}_m$ -torsor over the universal abelian variety (over a fine moduli space). Finally we define $(P_{2g}, \mathcal{X}_{2g}^\natural)$ be the enlarged mixed Shimura datum associated with $(P_{2g}, \mathcal{X}_{2g})$ (see Lemma 3.6).

3.5.3. Now define the enlarged mixed Shimura datum $(P_{2g,b}, \mathcal{X}_{2g,b}^\natural)^{[6]}$ which corresponds to the universal vectorial bi-extension defined by Coleman [12]. See Example 4.5(2) for further details.

Define the group $P_{2g,b}$ to be the unipotent extension of $P_{2g,a} = V_{2g} \rtimes \mathrm{GSp}_{2g}$ by $V_{2g} \oplus U_{2g}$ be the action $P_{2g,a} \curvearrowright V_{2g} \oplus U_{2g}$ defined by $(v, g)(v', u) := (gv', gu + \Psi(v, v'))$. Let $(P_{2g,b}, \mathcal{X}_{2g,b}^\natural)$ be the unipotent of $(P_{2g,a}, \mathcal{X}_{2g,a}^\natural)$ by $V_{2g} \oplus U_{2g}$ given by Proposition 3.8. Then its associated mixed Shimura datum $(P_{2g,b}, \mathcal{X}_{2g,b})$ is the unipotent extension of $(P_{2g,a}, \mathcal{X}_{2g,a})$ by $V_{2g} \oplus U_{2g}$ in the sense of Pink [33, 2.17]. Moreover by [34, Remark 2.13], the mixed Shimura datum $(P_{2g,b}, \mathcal{X}_{2g,b})$ corresponds to the Poincaré biextension over the product of the universal abelian variety with its dual (over a fine moduli space). We denote by $W_{2g,b} := \mathcal{R}_u(P_{2g,b})$.

3.6. Reduction Theorem for enlarged Shimura data.

Definition 3.10. An enlarged mixed Shimura datum $(P, \mathcal{X}^\natural, h)$ (resp. mixed Shimura datum $(P, \mathcal{X}, h)$) is said to **have generic Mumford-Tate group** if P possesses no proper normal subgroup Q such that for one (equivalently all) $x \in \mathcal{X}^\natural$ (resp. $x \in \mathcal{X}$), h_x factors through $Q_{\mathbb{C}} \subset P_{\mathbb{C}}$. We shall denote this case by $P = \mathrm{MT}(\mathcal{X}^\natural)$ (resp. $P = \mathrm{MT}(\mathcal{X})$).

Remark 3.11. (1) Pink [33, 2.13] used the term “irreducible” for this definition in the case of mixed Shimura data.

(2) Let M and φ be as in Proposition 2.8. Then an enlarged mixed Shimura datum $(P, \mathcal{X}^\natural, h)$ has generic Mumford-Tate group iff $P = \mathrm{MT}(\varphi(h(\mathcal{X}^\natural)))$ (resp. a mixed Shimura datum $(P, \mathcal{X}, h)$ has generic Mumford-Tate group iff $P = \mathrm{MT}(\varphi(h(\mathcal{X})))$). But $\varphi(h(\mathcal{X}^\natural)) = \varphi(h(\mathcal{X}))$ by Proposition 2.12, so an enlarged mixed Shimura datum has generic Mumford-Tate group iff its associated mixed Shimura datum has generic Mumford-Tate group.

[6]The index “b” is short for “biextension”.

Proposition 3.12. *Let $(P, \mathcal{X}^\natural)$ be a connected mixed Shimura datum, then*

- (1) *there exists an enlarged mixed Shimura datum $(Q, \mathcal{Y}^\natural)$ having generic Mumford-Tate group such that $(Q, \mathcal{Y}^\natural) \hookrightarrow (P, \mathcal{X}^\natural)$ and $\mathcal{Y}^{\natural+} \simeq \mathcal{X}^{\natural+}$ under this embedding;*
- (2) *if $(P, \mathcal{X}^\natural)$ has generic Mumford-Tate group, then P acts on U via a scalar. In particular, any subgroup of U is normal in P .*

Proof. This is true for mixed Shimura data by Pink [33, 2.13, 2.14]. Then it suffices to apply Lemma 3.6. $\square$

We close this section by the following Reduction Theorem for enlarged mixed Shimura data.

Theorem 3.13 (Reduction Theorem). *Let $(P, \mathcal{X}^\natural)$ be an enlarged mixed Shimura datum with generic Mumford-Tate group. Assume $\dim V = 2g > 0$ (otherwise $V = 0$ and $(P, \mathcal{X}^\natural)$ is a mixed Shimura datum). Then there exist pure Shimura data $(T, \mathcal{Y})$ and $(G_0, \mathcal{D}_0)$, where T is a torus and $\mathcal{D}_0 \subset \text{Hom}(\mathbb{S}, G_{0, \mathbb{R}})$ such that the follows hold: there exists an enlarged mixed Shimura datum $(P_*, \mathcal{X}_*^\natural)$ with morphisms*

$$(P_*, \mathcal{X}_*^\natural) \twoheadrightarrow (P, \mathcal{X}^\natural) \text{ which is a } ((P_0, \mathcal{X}_0) \rightarrow (\mathbb{G}_m, \mathcal{H}_0))\text{-torsor}$$

$$\text{and } (P_*, \mathcal{X}_*^\natural) \xrightarrow{\lambda} (T, \mathcal{Y}) \times (G_0, \mathcal{D}_0) \times \prod_{i=1}^r (P_{2g}, \mathcal{X}_{2g}^\natural)$$

such that $\lambda|_V: V \simeq V_{2g} \rightarrow \oplus_{i=1}^r V_{2g}$ is the diagonal map, $\lambda|_{U_*}: U_* \simeq \oplus_{i=1}^r U_{2g}$ and $G \xrightarrow{\bar{\lambda}} T \times G_0 \times \prod_{i=1}^r \text{GSp}_{2g} \rightarrow \text{GSp}_{2g}$ is non-trivial for each projection.

Proof. The first part is [33, 2.26(a)]. For the second part, let $(P, \mathcal{X})$ be mixed Shimura datum associated with $(P, \mathcal{X}^\natural)$. Then by [33, 2.26(b)] and [16, Lemma 2.12], there exists a mixed Shimura datum $(P_*, \mathcal{X}_*)$ with Shimura morphisms

$$(P_*, \mathcal{X}_*) \twoheadrightarrow (P, \mathcal{X}) \text{ and } (P_*, \mathcal{X}_*) \hookrightarrow (T, \mathcal{Y}) \times (G_0, \mathcal{D}_0) \times \prod_{i=1}^r (P_{2g}, \mathcal{X}_{2g})$$

with the desired properties. Now it suffices to take $(P_*, \mathcal{X}_*^\natural)$ to be the enlarged mixed Shimura datum associated with $(P_*, \mathcal{X}_*)$. $\square$

4. ENLARGED MIXED SHIMURA VARIETIES

4.1. Basic definition and complex space structure.

Definition 4.1. (1) *Let $(P, \mathcal{X}^\natural)$ be an enlarged mixed Shimura datum and let K be an open compact subgroup of $P(\mathbb{A}_f)$. Define the corresponding **enlarged mixed Shimura variety** as*

$$\text{EM}_K(P, \mathcal{X}^\natural)_{\mathbb{C}} := P(\mathbb{Q}) \backslash \mathcal{X}^\natural \times P(\mathbb{A}_f) / K,$$

*where $P(\mathbb{Q})$ acts diagonally on both factors from the left. This enlarged mixed Shimura variety is called **of Kuga type** if $(P, \mathcal{X}^\natural)$ is of Kuga type. We shall see that $\text{EM}_K(P, \mathcal{X}^\natural)_{\mathbb{C}}$ is a complex analytic variety (Remark 4.2).*

(2) Under the notation of (1) and §3.4, we say that the mixed Shimura variety

$$M_K(P, \mathcal{X})(\mathbb{C}) = P(\mathbb{Q}) \backslash \mathcal{X} \times P(\mathbb{A}_f) / K$$

is the **mixed Shimura variety associated with** $EM_K(P, \mathcal{X}^\natural)_{\mathbb{C}}$ and $EM_K(P, \mathcal{X}^\natural)_{\mathbb{C}}$ is the **enlarged mixed Shimura variety associated with** $M_K(P, \mathcal{X})(\mathbb{C})$. Here $M_K(P, \mathcal{X})(\mathbb{C})$ means the $\mathbb{C}$ -points of the algebraic variety $M_K(P, \mathcal{X})$ over the number field $E(P, \mathcal{X})$ on which it is defined (this is proven by Pink).

Remark 4.2. Let $\mathcal{X}^{\natural+}$ be a connected component of $\mathcal{X}^\natural$, and thus $\mathcal{X}^{\natural+}$ is a $P(\mathbb{R})^+W(\mathbb{C})$ -homogeneous space. Then as for mixed Shimura varieties, we have

$$EM_K(P, \mathcal{X}^\natural)_{\mathbb{C}} = \bigcup_{[p] \in P(\mathbb{Q})_+ \backslash P(\mathbb{A}_f) / K} \Gamma(p) \backslash \mathcal{X}^{\natural+}$$

where $\Gamma(p) := P(\mathbb{Q})_+ \cap pKp^{-1}$.

Proposition 4.3. The action of $\Gamma(p)$ on $\mathcal{X}^{\natural+}$ factors through a quotient $\overline{\Gamma(p)}$ which acts properly discontinuously on $\mathcal{X}^{\natural+}$. There is a canonical structure of a normal complex space on $EM_K(P, \mathcal{X}^\natural)_{\mathbb{C}}$, whose singularities are quotient singularities by finite groups. Moreover $EM_K(P, \mathcal{X}^\natural)_{\mathbb{C}}$ is a complex manifold if K is neat.

Proof. (Compare with [33, 3.3]) The image of $\Gamma(p)$ in $(P/P^{\text{der}}Z(P))(\mathbb{Q})$ is finite by Definition 3.1(5), so it suffices to prove the assertion for $\Gamma := \Gamma(p) \cap (P^{\text{der}}Z(P))(\mathbb{Q})$. Note that $\Gamma = \Gamma(p)$ when K is neat. Now $Z(P)(\mathbb{R})$ acts trivially on $\mathcal{X}^{\natural+}$, so we may replace $(P, \mathcal{X}^\natural)$ by $(P, \mathcal{X}^\natural)/Z(P)$. Then Γ is an arithmetic subgroup of $P^{\text{der}}(\mathbb{Q})$, torsion free if K is neat, and therefore is a discrete subgroup of $P^{\text{der}}(\mathbb{R})$. So Γ acts properly discontinuously on $P_\infty := P^{\text{der}}(\mathbb{R})^+W(\mathbb{C})$. But $\mathcal{X}^{\natural+} \simeq P_\infty / \text{Stab}_{P_\infty}(x)$ for any $x \in \mathcal{X}^{\natural+}$ and $\text{Stab}_{P_\infty}(x)$ is compact by Lemma 2.10 and Definition 3.1(4). So Γ acts properly discontinuously on $\mathcal{X}^{\natural+}$. Note that $\Gamma \cap \text{Stab}_{P_\infty}(x)$ is finite, and hence trivial if K is neat. The complex structure of $\Gamma(p) \backslash \mathcal{X}^{\natural+}$ comes from the $P(\mathbb{R})W(\mathbb{C})$ -invariant complex structure on $\mathcal{X}^\natural$, and $\Gamma(p) \backslash \mathcal{X}^{\natural+}$ is a complex manifold when K is neat. $\square$

4.2. Algebraic structure and canonical model. The goal of this section is to prove:

Theorem 4.4. Every enlarged mixed Shimura variety $EM_K(P, \mathcal{X}^\natural)_{\mathbb{C}}$ is an algebraic variety. It has a model $EM_K(P, \mathcal{X}^\natural)$ over $E(P, \mathcal{X}) := \text{the reflex field of } (P, \mathcal{X})^{[7]}$, called the **canonical model**, such that the map induced by the geometric comparison (3.1)

$$[\pi^\natural]: EM_K(P, \mathcal{X}^\natural)_{\mathbb{C}} \rightarrow M_K(P, \varphi(\mathcal{D}))(\mathbb{C})$$

descends to a map over the field $E(P, \mathcal{X}) = E(P, \varphi(\mathcal{D}))$

$$EM_K(P, \mathcal{X}^\natural) \rightarrow M_K(P, \varphi(\mathcal{D}))$$

where $M_K(P, \varphi(\mathcal{D}))$ is the canonical model of $M_K(P, \varphi(\mathcal{D}))_{\mathbb{C}}$.

An important tool to prove this theorem is the geometric comparison considered in §3.1.

^[7]See [33, 11.1] for definition.

4.2.1. *Universal vector extension of the universal abelian variety.* Let $(P_{2g,a}, \mathcal{X}_{2g,a}^\natural)$ and $(P_{2g,a}, \mathcal{X}_{2g,a})$ as in §3.5.1. Let $N \geq 3$ be an integer. Define

$$K_{\mathrm{GSp}}(N) := \{h \in \mathrm{GSp}_{2g}(\mathbb{Z}) \mid h \equiv 1 \pmod{N}\} \text{ and } K_{2g,a}(N) := V_{2g}(\mathbb{Z}) \rtimes K_{\mathrm{GSp}}(N).$$

Consider $\mathrm{EM}_{K_{2g,a}(N)}(P_{2g,a}, \mathcal{X}_{2g,a}^\natural)_{\mathbb{C}}$ and the geometric comparison (3.7). Denote by $\mathcal{A}_g(N) := \mathrm{Sh}_{K_{\mathrm{GSp}}(N)}(\mathrm{GSp}_{2g}, \mathcal{H}_g)$ and $\mathfrak{A}_g(N) := (P_{2g,a}, \mathcal{X}_{2g,a})$.

Pink proved [33, 10.10] that $\mathrm{M}_{K_{2g,a}(N)}(P_{2g,a}, \varphi(\mathcal{X}_{2g,a})) \simeq \mathfrak{A}_g(N)$ is the universal abelian variety over the fine moduli space $\mathrm{Sh}_{K_{\mathrm{GSp}}(N)}(\mathrm{GSp}_{2g}, \varphi(\mathcal{H}_g)) \simeq \mathcal{A}_g(N)$ and therefore is an algebraic variety. To better illustrate the moduli interpretation of $\mathrm{EM}_{K_{2g,a}(N)}(P_{2g,a}, \mathcal{X}_{2g,a}^\natural)_{\mathbb{C}}$, we consider its connected components given by Remark 4.2.

Denote by $\mathcal{V}_{2g}^+ := V_{2g}(\mathbb{C}) \times \mathcal{H}_g^+$. Let $\Gamma_{\mathrm{GSp}}(N) := \mathrm{Sp}_{2g}(\mathbb{Z}) \cap K_{\mathrm{GSp}}(N)$ and

$$\Gamma_{2g,a}(N) := P_{2g,a}^{\mathrm{der}}(\mathbb{Z}) \cap K_{2g,a}(N) = V_{2g}(\mathbb{Z}) \rtimes \Gamma_{\mathrm{GSp}}(N).$$

Denote by $\mathfrak{A}_g(N)^+$ (resp. $\mathcal{A}_g(N)^+$) the connected component $\Gamma_{2g,a}(N) \backslash \varphi(\mathcal{X}_{2g,a}^+)$ (resp. $\Gamma_{\mathrm{GSp}}(N) \backslash \varphi(\mathcal{H}_g^+)$) of $\mathfrak{A}_g(N)$ (resp. $\mathcal{A}_g(N)$). From the geometric comparison (3.7) we have

$$\begin{aligned} 0 \rightarrow \mathcal{F}^0 \mathcal{V}_{2g}^+ \rightarrow \mathcal{X}_{2g,a}^{\natural+} = \mathcal{V}_{2g}^+ \rightarrow \varphi(\mathcal{X}_{2g,a}^+) = \mathcal{V}_{2g}^+ / \mathcal{F}^0 \mathcal{V}_{2g}^+ \rightarrow 0 \text{ as vector bundles over } \mathcal{H}_g^+ \\ \Rightarrow 0 \rightarrow \Gamma_{\mathrm{GSp}}(N) \backslash \mathcal{F}^0 \mathcal{V}_{2g}^+ \rightarrow \Gamma_{\mathrm{GSp}}(N) \backslash \mathcal{V}_{2g}^+ \rightarrow \Gamma_{\mathrm{GSp}}(N) \backslash \varphi(\mathcal{X}_{2g,a}^+) \rightarrow 0 \text{ as vector bundles}^{[8]} \text{ over } \mathcal{A}_g(N)^+ \\ \Rightarrow 0 \rightarrow \Gamma_{\mathrm{GSp}}(N) \backslash \mathcal{F}^0 \mathcal{V}_{2g}^+ \rightarrow V_{2g}(\mathbb{Z}) \backslash \left(\Gamma_{\mathrm{GSp}}(N) \backslash \mathcal{V}_{2g}^+ \right) \rightarrow V_{2g}(\mathbb{Z}) \backslash \left(\Gamma_{\mathrm{GSp}}(N) \backslash \varphi(\mathcal{X}_{2g,a}^+) \right) \rightarrow 0, \\ \text{i.e. } 0 \rightarrow \Gamma_{\mathrm{GSp}}(N) \backslash \mathcal{F}^0 \mathcal{V}_{2g}^+ \rightarrow \Gamma_{2g,a}(N) \backslash \mathcal{V}_{2g}^+ \rightarrow \mathfrak{A}_g(N)^+ \text{ as Lie groups over } \mathcal{A}_g(N)^+ \\ \Rightarrow 0 \rightarrow \omega_{\mathfrak{A}_g(N)^+ / \mathcal{A}_g(N)^+} \rightarrow \Gamma_{2g,a}(N) \backslash \mathcal{V}_{2g}^+ \rightarrow \mathfrak{A}_g(N)^+ \rightarrow 0 \text{ as Lie groups over } \mathcal{A}_g(N)^+ \end{aligned}$$

Here the last implication is the canonical isomorphism $\omega_{\mathfrak{A}_g(N)^+ / \mathcal{A}_g(N)^+} \simeq \Gamma_{\mathrm{GSp}}(N) \backslash \mathcal{F}^0 \mathcal{V}_{2g}^+$ implied by the definition of $\mathcal{F}^0 \mathcal{V}_{2g}^+$, and $\mathfrak{A}_g(N)^{\natural+} := \Gamma(N) \backslash \mathcal{V}_{2g}^+$ is the universal vector extension of the universal abelian variety $\mathfrak{A}_g(N)^+$ over $\mathcal{A}_g(N)^+$. In particular $\mathfrak{A}_g(N)^{\natural+}$ is an algebraic variety and the maps in the last line above are algebraic.

The above discussion holds for any $pK_{2g,a}(N)p^{-1} \cap P_{2g,a}(\mathbb{Q})_+$. So by Remark 4.2 $\mathfrak{A}_g(N)^{\natural} := \mathrm{EM}_{K_{2g,a}(N)}(P_{2g,a}, \mathcal{X}_{2g,a}^\natural)_{\mathbb{C}}$ is the universal vector extension of the universal abelian variety $\mathfrak{A}_g(N)$ over the fine moduli space $\mathcal{A}_g(N)$, and in particular is an algebraic variety. But $\mathfrak{A}_g(N)$ has a canonical model over $\mathbb{Q}$, so $\mathfrak{A}_g(N)^{\natural}$ has a model over $\mathbb{Q}$ and the projection $\mathfrak{A}_g(N)^{\natural} \rightarrow \mathfrak{A}_g(N)$ is defined over $\mathbb{Q}$.

4.2.2. *General Case.* It suffices to consider the case $\dim V = 2g > 0$ since otherwise $\mathrm{EM}_K(P, \mathcal{X}^\natural)_{\mathbb{C}}$ is a mixed Shimura variety and the result is known by Pink. By Proposition 3.12(1) and Remark 4.2, we may assume that $(P, \mathcal{X}^\natural)$ has generic Mumford-Tate group. By Lemma 3.9 we may assume that $\mathcal{X}^\natural \subset \mathrm{Hom}(\mathbb{S}_{\mathbb{C}}, P_{\mathbb{C}})$. Denote by $K_{P/U} := \pi_{P/U}(K) < P/U$, $K_W := K \cap W(\mathbb{A}_f)$ and $K_V := K_W / (K \cap U(\mathbb{A}_f))$. Up to replacing K be a subgroup of finite index we may assume $K = K_W \rtimes K_G$ and $K_V \subset K_{2g,a}(N)$ for some $N \geq 4$ even.

[8] In fact these vector bundles are automorphism bundles over $\mathcal{A}_g(N)^+$. This is because the variation of pure Hodge structures $(\mathcal{V}_{2g}^+, \mathcal{F}^0)$ over $\mathcal{H}_g^+$ extends naturally to $\mathcal{H}_g^\vee$: the fibers over $\mathcal{H}_g^+$ define polarizable Hodge structures of weight -1 , whereas a general fiber over $\mathcal{H}_g^\vee$ defines Hodge structure of weight -1 not necessarily polarizable.

We start with $\mathrm{EM}_{K_{P/U}}(P/U, \mathcal{X}_{P/U}^\natural)_{\mathbb{C}}$. Use the notation of §3.4 (note that we are in the case $\mathcal{X}^\natural = \mathcal{D}^\natural$). Apply Theorem 3.13(2) to $(P/U, \mathcal{X}_{P/U}^\natural)$ and we get an inclusion

$$(P/U, \mathcal{X}_{P/U}^\natural) \hookrightarrow (T, \mathcal{Y}) \times (G_0, \mathcal{D}_0) \times (P_{2g,a}, \mathcal{X}_{2g,a}^\natural),$$

which induces a Cartesian diagram in the category of complex varieties, where all objects and morphisms are algebraic except those concerning $\mathrm{EM}_{K_{P/U}}(P/U, \mathcal{X}_{P/U}^\natural)_{\mathbb{C}}$:

$$\begin{array}{ccc} \mathrm{EM}_{K_{P/U}}(P/U, \mathcal{X}_{P/U}^\natural)_{\mathbb{C}} & \longrightarrow & \mathrm{Sh}_{K_T}(T, \mathcal{Y})(\mathbb{C}) \times \mathrm{Sh}_{K_{G_0}}(G_0, \mathcal{D}_0)(\mathbb{C}) \times \mathfrak{A}_g(N)^\natural . \\ \downarrow [\pi_{P/U}^\natural] \lrcorner & & \downarrow \\ \mathrm{M}_{K_{P/U}}(P/U, \varphi(\mathcal{X})_{P/U})(\mathbb{C}) & \longrightarrow & \mathrm{Sh}_{K_T}(T, \mathcal{Y})(\mathbb{C}) \times \mathrm{Sh}_{K_{G_0}}(G_0, \mathcal{D}_0)(\mathbb{C}) \times \mathfrak{A}_g(N) \end{array}$$

But $\mathfrak{A}_g(N)^\natural$ is the universal vector extension of the universal abelian scheme $\mathfrak{A}_g(N)$ over $\mathcal{A}_g(N)$, so we have that $\mathrm{EM}_{K_{P/U}}(P/U, \mathcal{X}_{P/U}^\natural)_{\mathbb{C}}$ is the universal vector extension of the abelian scheme $\mathrm{M}_{K_{P/U}}(P/U, \varphi(\mathcal{X})_{P/U})(\mathbb{C}) \rightarrow \mathrm{Sh}_{K_G}(G, \varphi(\mathcal{X})_G)(\mathbb{C})$. So Theorem 4.4 holds for $\mathrm{EM}_{K_{P/U}}(P/U, \mathcal{X}_{P/U}^\natural)_{\mathbb{C}}$.

Now we turn to $\mathrm{EM}_K(P, \mathcal{X}^\natural)_{\mathbb{C}}$. The Cartesian diagram in (3.3) induces a Cartesian diagram in the category of complex varieties, where all objects and morphism are algebraic over $E(P, \mathcal{X})$ except those concerning $\mathrm{EM}_K(P, \mathcal{X}^\natural)_{\mathbb{C}}$:

$$(4.1) \quad \begin{array}{ccc} \mathrm{EM}_K(P, \mathcal{X}^\natural)_{\mathbb{C}} & \longrightarrow & \mathrm{EM}_{K_{P/U}}(P/U, \mathcal{X}_{P/U}^\natural)_{\mathbb{C}} \\ \downarrow \lrcorner & & \downarrow \\ \mathrm{M}_K(P, \varphi(\mathcal{X}))(\mathbb{C}) & \longrightarrow & \mathrm{M}_{K_{P/U}}(P/U, \varphi(\mathcal{X})_{P/U})(\mathbb{C}) \end{array}$$

But $\mathrm{M}_K(P, \varphi(\mathcal{X})) \rightarrow \mathrm{M}_{K_{P/U}}(P/U, \varphi(\mathcal{X})_{P/U})$ is a T -torsor for some algebraic group T over $E(P, \mathcal{X})$ by Pink [33, 3.12, 3.18]. So Theorem 4.4 holds for $\mathrm{EM}_K(P, \mathcal{X}^\natural)_{\mathbb{C}}$.

Example 4.5. (1) Let $(P_{2g}, \mathcal{X}_{2g}^\natural)$ be as in §3.5.2. Keep the notation of §4.2.1. Let $N \geq 4$ be an even integer. Define $K_{2g}(N) := W_{2g}(\mathbb{Z}) \rtimes K_{\mathrm{GSp}}(N)$. Denote by $\mathfrak{L}_g(N) := \mathrm{M}_{K_{2g}(N)}(P_{2g}, \mathcal{X}_{2g})$. Pink [33, 10.10] proved that $\mathfrak{L}_g(N)$ is (the total space of) the canonical symmetric relatively ample $\mathbb{G}_m$ -torsor over the universal abelian variety $\mathfrak{A}_g(N) \rightarrow \mathcal{A}_g(N)$. The proof of Theorem 4.4 shows that $\mathrm{EM}_K(P_{2g}, \mathcal{X}_{2g})$, which we denote by $\mathfrak{L}_g^\natural(N)$, is defined over $\mathbb{Q}$ and is the pullback of the $\mathbb{G}_m$ -torsor $\mathfrak{L}_g(N)$ over the universal abelian variety $\mathfrak{A}_g(N) \rightarrow \mathcal{A}_g(N)$ to the universal vector extension $\mathfrak{A}_g(N)^\natural$ of $\mathfrak{A}_g(N)$.

(2) Let $(P_{2g,b}, \mathcal{X}_{2g,b}^\natural)$ be as in §3.5.3. Keep the notation of §4.2.1. Let $N \geq 4$ be an even integer. Define $K_{2g,b}(N) := W_{2g,b}(\mathbb{Z}) \rtimes K_{\mathrm{GSp}}(N)$. Pink proved in [34, Remark 2.13] that the mixed Shimura variety $\mathfrak{P}_g(N) := \mathrm{M}_{K_{2g,b}(N)}(P_{2g,b}, \mathcal{X}_{2g,b})$ is (the total space of) the Poincaré biextension over $\mathfrak{A}_g(N) \times_{\mathcal{A}_g(N)} \mathfrak{A}_g(N)^\vee$. The proof of Theorem 4.4 shows that $\mathrm{EM}_{K_{2g,b}(N)}(P_{2g,b}, \mathcal{X}_{2g,b}^\natural)$, which we denote by $\mathfrak{P}_g^\natural(N)$, is defined over $\mathbb{Q}$ and is the pullback the Poincaré biextension $\mathfrak{P}_g(N)$ to $\mathfrak{A}_g(N)^\natural \times_{\mathcal{A}_g(N)} \mathfrak{A}_g(N)^\vee$. This is the “universal vectorial bi-extension” considered by Coleman in [12].

4.3. Morphisms between enlarged mixed Shimura varieties.

Proposition 4.6. *Let $f: (P_1, \mathcal{X}_1^\natural) \rightarrow (P_2, \mathcal{X}_2^\natural)$ be a morphism of enlarged mixed Shimura data, and let $K_1 \subset P_1(\mathbb{A}_f)$ and $K_2 \subset P_2(\mathbb{A}_f)$ be open compact subgroups such that $f(K_1) \subset K_2$. Then we have:*

(1) *The canonical map induces by f*

$$[f]: \text{EM}_{K_1}(P_1, \mathcal{X}_1^\natural)_{\mathbb{C}} \rightarrow \text{EM}_{K_2}(P_2, \mathcal{X}_2^\natural)_{\mathbb{C}}, \quad [(x, p)] \mapsto [(f(x), f(p))]$$

is algebraic, closed and descends to $E(P_1, \mathcal{X}_1)$.

(2) *If furthermore f is an embedding such that $K_1 = f^{-1}(K_2)$, then $[f]$ is finite.*

(3) *If f is an embedding, then for every K_1 there exists a K_2 such that $[f]$ is a closed embedding.*

Proof. This proposition can be proven using the geometric interpretations in §4.2.2. By Proposition 3.12(1) and Remark 4.2, we may assume that $(P_i, \mathcal{X}_i^\natural)$ has generic Mumford-Tate group for $i = 1, 2$. By Lemma 3.9 we may assume that $\mathcal{X}_i^\natural \subset \text{Hom}(\mathbb{S}_{\mathbb{C}}, P_{i, \mathbb{C}})$ for $i = 1, 2$. By Pink [33, 3.8, 9.24] the proposition is true for

$$\text{M}_{K_1}(P_1, \varphi_1(\mathcal{X}_1)) \rightarrow \text{M}_{K_2}(P_2, \varphi_2(\mathcal{X}_2)) \text{ and } \text{M}_{K_{1, P/U}}(P_1/U_1, \varphi(\mathcal{X})_{1, P/U}) \rightarrow \text{M}_{K_{2, P/U}}(P_2/U_2, \varphi(\mathcal{X})_{2, P/U})$$

Hence by (4.1) it suffices to prove the proposition for

$$[f_{P/U}]: \text{EM}_{K_{1, P/U}}(P_1/U_1, \mathcal{X}_{1, P/U}^\natural)_{\mathbb{C}} \rightarrow \text{EM}_{K_{2, P/U}}(P_2/U_2, \mathcal{X}_{2, P/U}^\natural)_{\mathbb{C}}.$$

This is true because $\text{EM}_{K_{i, P/U}}(P_i/U_i, \mathcal{X}_{i, P/U}^\natural)$ is the universal vector extension of the abelian scheme $\text{M}_{K_{i, P/U}}(P_i/U_i, \varphi(\mathcal{X})_{i, P/U})$ over $\text{Sh}_{K_G}(G, \varphi(\mathcal{X})_G)$. $\square$

5. BI-ALGEBRAIC SETTING FOR ENLARGED MIXED SHIMURA VARIETIES

From this section, we will only consider connected enlarged mixed Shimura data and connected enlarged mixed Shimura varieties.

Definition 5.1. (1) *A **connected enlarged mixed Shimura datum** is a triple $(P, \mathcal{X}^{\natural+}, h)$ such that $(P, \mathcal{X}^{\natural+}, h) \subset (P, \mathcal{X}^\natural, h)$ for some enlarged mixed Shimura datum $(P, \mathcal{X}^\natural, h)$ with h injective and that $\mathcal{X}^{\natural+}$ is a left $P(\mathbb{R})^+W(\mathbb{C})$ -homogeneous space. It is said to **have generic Mumford-Tate group** if $(P, \mathcal{X}^\natural, h)$ has generic Mumford-Tate group.*

(2) *A **connected enlarged mixed Shimura variety** $S^\natural$ associated with the connected enlarged mixed Shimura datum $(P, \mathcal{X}^{\natural+}, h)$ is a quotient $\Gamma \backslash \mathcal{X}^{\natural+}$ for some congruence subgroup Γ of $P(\mathbb{Q})_+ := P(\mathbb{Q}) \cap P(\mathbb{R})_+$, where $P(\mathbb{R})_+$ is the stabilizer in $P(\mathbb{R})$ of $h(\mathcal{X}^{\natural+}) \subset \text{Hom}(\mathbb{S}_{\mathbb{C}}, P_{\mathbb{C}})$.*

Remark 5.2. (1) *For any enlarged mixed Shimura datum $(P, \mathcal{X}^\natural, h)$, let $\mathcal{X}^{\natural+}$ be a connected component of $\mathcal{X}^\natural$. Then $(P, \mathcal{X}^{\natural+}, h)$ is a connected enlarged mixed Shimura datum. Conversely every connected enlarged mixed Shimura datum arises in this way.*

(2) *By Remark 4.2, connected enlarged mixed Shimura varieties are precisely connected components of enlarged mixed Shimura varieties. So by Theorem 4.4 and Proposition 4.6, any connected enlarged mixed Shimura variety is algebraic and has a model over $\overline{\mathbb{Q}}$ and any Shimura morphism between connected enlarged mixed Shimura varieties is algebraic and descends to $\overline{\mathbb{Q}}$.*

(3) In practice we often consider the case $Z(P) = 1$. This can be achieved by replacing $(P, \mathcal{X}^{\natural+})$ by $(P, \mathcal{X}^{\natural+})/Z(P)$. Under this hypothesis every neat congruence subgroup is contained in $P^{\text{der}}(\mathbb{Q})$. Conversely fix a Levi decomposition $P = W \rtimes G$, then $P^{\text{der}} = W \rtimes G^{\text{der}}$. Hence any congruence subgroup $\Gamma < P^{\text{der}}(\mathbb{Q})$ is Zariski dense in P^{der} by Definition 3.1(4) and [35, Theorem 4.10].

We are interested in the uniformization $\text{unif}^{\natural}: \mathcal{X}^{\natural+} \rightarrow S^{\natural} = \Gamma \backslash \mathcal{X}^{\natural+}$. Use the notation of §3.1 (note that $\mathcal{D}^{\natural+} \simeq \mathcal{X}^{\natural+}$ and $\mathcal{D}^+ \simeq \mathcal{X}^+$ and so we do not distinguish $\mathcal{X}$ and $\mathcal{D}$). Let $\Gamma_{P/U} := \pi_{P/U}(\Gamma)$ and let $\Gamma_G := \pi(\Gamma)$. We will use indices to distinguish different uniformizations:

$$\begin{aligned} \mathcal{X}_{P/U}^{\natural+} &\xrightarrow{\text{unif}_{P/U}^{\natural}} S_{P/U}^{\natural} := \Gamma_{P/U} \backslash \mathcal{X}_{P/U}^{\natural+} & \varphi(\mathcal{X}^+) &\xrightarrow{\text{unif}} S := \Gamma \backslash \varphi(\mathcal{X}^+) \\ \varphi(\mathcal{X}^+)_{P/U} &\xrightarrow{\text{unif}_{P/U}} S_{P/U} := \Gamma_{P/U} \backslash \varphi(\mathcal{X}^+)_{P/U} & \varphi(\mathcal{X}^+)_G &\xrightarrow{\text{unif}_G} S_G := \Gamma_G \backslash \varphi(\mathcal{X}^+)_G \end{aligned}$$

Notation 5.3. For any subset $\tilde{Y}^{\natural}$ of $\mathcal{X}^{\natural+}$, the diagram in §3.4 gives rise to

$$\begin{array}{ccc} \mathcal{X}^{\natural+} & \xrightarrow{\pi^{\sharp}} & \varphi(\mathcal{X}^+) \\ \downarrow \pi_{P/U}^{\natural} & \lrcorner & \downarrow \pi_{P/U} \\ \mathcal{X}_{P/U}^{\natural+} & \xrightarrow{\pi_{P/U}^{\sharp}} & \varphi(\mathcal{X}^+)_{P/U} \\ \downarrow \pi_G^{\natural} & & \downarrow \pi_G \\ \varphi(\mathcal{X}^+)_G & \xrightarrow{\text{id}} & \varphi(\mathcal{X}^+)_G \end{array} \quad \begin{array}{ccc} \tilde{Y}^{\natural} & \hookrightarrow & \tilde{Y} \\ \downarrow & & \downarrow \\ \tilde{Y}_{P/U}^{\natural} & \hookrightarrow & \tilde{Y}_{P/U} \\ \downarrow & & \downarrow \\ \tilde{Y}_G & \xlongequal{\quad} & \tilde{Y}_G \end{array}$$

For any subset $Y^{\natural}$ of $S^{\natural}$, the diagram in §3.4 gives rise to

$$\begin{array}{ccc} S^{\natural} & \xrightarrow{[\pi^{\sharp}]} & S \\ \downarrow [\pi_{P/U}^{\natural}] & \lrcorner & \downarrow [\pi_{P/U}] \\ S_{P/U}^{\natural} & \xrightarrow{[\pi_{P/U}^{\sharp}]} & S_{P/U} \\ \downarrow [\pi_G^{\natural}] & & \downarrow [\pi_G] \\ S_G & \xrightarrow{\text{id}} & S_G \end{array} \quad \begin{array}{ccc} Y^{\natural} & \hookrightarrow & Y \\ \downarrow & & \downarrow \\ Y_{P/U}^{\natural} & \hookrightarrow & Y_{P/U} \\ \downarrow & & \downarrow \\ Y_G & \xlongequal{\quad} & Y_G \end{array}$$

5.1. Algebraic structure on $\mathcal{X}^{\natural+}$ and bi-algebraic subsets. Recall that the complex structure on $\mathcal{X}^{\natural+}$ is given by $\mathcal{X}^{\natural+} \simeq W(\mathbb{C}) \times \varphi(\mathcal{X}^+)_G$ and $\varphi(\mathcal{X}^+)_G \hookrightarrow \mathcal{X}_G^{\vee} = G(\mathbb{C})/F_{x_G}G_{\mathbb{C}}$ (for any $x_G \in \mathcal{X}_G$) is a complex algebraic variety and $\varphi(\mathcal{X}^+)_G$ is a semi-algebraic subset of $\mathcal{X}_G^{\vee}$, open in the usual topology.

Definition 5.4. Let $\tilde{Y}^{\natural}$ be a complex analytic subset of $\mathcal{X}^{\natural+}$, then

- (1) $\tilde{Y}^{\natural}$ is called an **irreducible algebraic subset** of $\mathcal{X}^{\natural+}$ if $\tilde{Y}^{\natural}$ is a complex analytic irreducible component of the intersection of its Zariski closure in $W(\mathbb{C}) \times \mathcal{X}_G^{\vee}$ with $\mathcal{X}^{\natural+}$.
- (2) $\tilde{Y}^{\natural}$ is called **algebraic** if it is a finite union of irreducible algebraic subsets of $\mathcal{X}^{\natural+}$.

The inclusion $\varphi(\mathcal{X}^+) \simeq \mathcal{X}^+ \hookrightarrow \mathcal{X}^\vee$ defines an algebraic structure on $\varphi(\mathcal{X}^+)$ (see [16, §6]). All maps in the diagrams in Notation 5.3 are algebraic. To see this, it suffices to prove the algebraicity of $\pi^\sharp: \mathcal{X}^{\natural+} \rightarrow \varphi(\mathcal{X}^+)$, which fits into the commutative diagram

$$\begin{array}{ccc} \mathcal{X}^{\natural+} = P(\mathbb{R})^+ W(\mathbb{C}) / \text{Cent}(h_x) \simeq W(\mathbb{C}) \times \varphi(\mathcal{X}^+)_G & \xhookrightarrow{\quad} & P(\mathbb{C}) / F_{x_G}^0 G_{\mathbb{C}} \simeq W(\mathbb{C}) \times \mathcal{X}_G^\vee \\ \downarrow \pi^\sharp & & \downarrow \\ \varphi(\mathcal{X}^+) = P(\mathbb{R})^+ W(\mathbb{C}) / \text{Stab}_{P(\mathbb{R})+W(\mathbb{C})}(\varphi(h_x)) & \xhookrightarrow{\quad} & \mathcal{X}^\vee = P(\mathbb{C}) / F_{x_G}^0 P_{\mathbb{C}} \end{array}$$

and hence is algebraic.

Now we are in the following situation: under $\text{unif}^\natural: \mathcal{X}^{\natural+} \rightarrow S^\natural$, the two spaces $\mathcal{X}^{\natural+}$ and $S^\natural$ are algebraic but the map $\text{unif}^\natural$ is transcendental. We want to understand the bi-algebraic objects in this context.

Definition 5.5. (1) A subset $\tilde{Y}^\natural$ is called **bi-algebraic** if $\tilde{Y}^\natural$ is an irreducible algebraic subset of $\mathcal{X}^{\natural+}$ and $Y^\natural := \text{unif}^\natural(\tilde{Y}^\natural)$ is a closed algebraic subvariety of $S^\natural$.
 (2) A subvariety $Y^\natural$ of $S^\natural$ is said to be **bi-algebraic** if it is the image of some bi-algebraic subset of $\mathcal{X}^{\natural+}$ under $\text{unif}^\natural$.

We will give a more concrete description of bi-algebraic subvarieties. In order to do this we introduce “quasi-linear subvarieties”.

5.2. Quasi-linear subvarieties and characterization of bi-algebraic subvarieties. Let $\tilde{Y}^\natural$ be a subset of $\mathcal{X}^{\natural+}$. Use Notation 5.3. Let $(Q, \mathcal{Y}^+)$ be the smallest connected mixed Shimura subdatum of $(P, \varphi(\mathcal{X}^+))$ such that $\tilde{Y}_{P/U} \subset \mathcal{Y}_{Q/U_Q}^+$. Assume that $\tilde{Y}_{P/U}$ is weakly special. Then $\tilde{Y}_{P/U} = (N/U_N)(\mathbb{R})^+ \tilde{y}_{P/U}$ for some normal subgroup N of Q (with $U_N := U \cap N$) and a point $\tilde{y}_{P/U} \in \tilde{Y}_{P/U}$. By (3.2) we have for $(P/U, \varphi(\mathcal{X}^+)_{P/U})$

$$(5.1) \quad 0 \rightarrow \mathcal{F}^0 \mathcal{V} \rightarrow \mathcal{X}_{P/U}^{\natural+} = \mathcal{V} \xrightarrow{\pi_{P/U}^\sharp} \varphi(\mathcal{X}^+)_{P/U} = \mathcal{V} / \mathcal{F}^0 \mathcal{V} \rightarrow 0.$$

Let $V^{(0)}$ be the largest $G_{Q,\mathbb{R}}$ -submodule of $V_{\mathbb{R}}$ on which $G_N(\mathbb{R})^+$ acts trivially. Then we have a unique inclusion

$$(5.2) \quad \tilde{Y}_{P/U}^{\text{univ}} \times_{\tilde{Y}_G} \tilde{\mathcal{C}}^\natural \subset \mathcal{X}_{P/U}^{\natural+}|_{\tilde{Y}_{P/U}} := \pi_{P/U}^{\sharp-1}(\tilde{Y}_{P/U})$$

where $\tilde{\mathcal{C}}^\natural = \mathcal{F}^0 \mathcal{V}^{(0)}|_{\tilde{Y}_G}$ and $\tilde{Y}_{P/U}^{\text{univ}}$ is defined by (compatible with (5.1))

$$0 \rightarrow \mathcal{F}^0 \mathcal{V}_N|_{\tilde{Y}_G} \rightarrow \tilde{Y}_{P/U}^{\text{univ}} \rightarrow \tilde{Y}_{P/U} \rightarrow 0.$$

Definition 5.6. (1) A subset $\tilde{Y}^\natural$ of $\mathcal{X}^{\natural+}$ is called **quasi-linear** if $\tilde{Y}$ is weakly special and

$$\tilde{Y}^\natural = \tilde{Y} \times_{\tilde{Y}_{P/U}} \left(\tilde{Y}_{P/U}^{\text{univ}} \times_{\tilde{Y}_G} (\tilde{L}^\natural \times \tilde{Y}_G) \times_{\tilde{Y}_G} \mathcal{F}^0 \mathcal{V}^\dagger|_{\tilde{Y}_G} \right)$$

where $V^\dagger$ is a $G_{Q,\mathbb{R}}$ -submodule of $V_{\mathbb{R}}$ complementary to $V_{N,\mathbb{R}}$ and $V^{(0)}$, and $\tilde{L}^\natural$ is an irreducible algebraic subvariety of some fiber of $\tilde{\mathcal{C}}^\natural \rightarrow \tilde{Y}_G$ (which is isomorphic to $\mathbb{C}^k$ for some k).

(2) A subset $Y^\natural$ of $S^\natural$ is said to be **quasi-linear** if it is the image of some quasi-linear subset of $\mathcal{X}^{\natural+}$ under $\text{unif}^\natural$.

It is clear that any quasi-linear subset of $\mathcal{X}^{\natural+}$ is complex analytic irreducible. Now we show that any quasi-linear subset $Y^{\natural}$ is a closed irreducible algebraic subvariety of $S^{\natural}$.

Up to replacing Γ by a finite subgroup we assume that $\Gamma = \Gamma_W \rtimes \Gamma_G$. Then $S_{P/U}^{\natural}$ is the universal vector extension of the abelian scheme $S_{P/U}$ over S_G . Let $Y^{\natural}$ be a bi-algebraic subset of $S^{\natural}$ and let $\tilde{Y}^{\natural}$ be the bi-algebraic subset of $\mathcal{X}^{\natural+}$ defining $Y^{\natural}$. Use Notation 5.3. Then Y is a weakly special subvariety of S . Then $Y_{P/U}$ is an abelian scheme over Y_G by [17, Proposition 1.1], and $S_{P/U}^{\natural}|_{Y_{P/U}} := [\pi_{P/U}^{\natural}]^{-1}(Y_{P/U})$ is a vector extension of the abelian scheme $Y_{P/U}$ over Y_G . Let $Y_{P/U}^{\text{univ}}$ be the universal vector extension of $Y_{P/U}$ over Y_G and consider the abelian scheme $[\pi_G]^{-1}(Y_G)$ over Y_G , then we have

$$S_{P/U}^{\natural}|_{Y_{P/U}} = Y_{P/U}^{\text{univ}} \times_{Y_G} \left(\omega_{[\pi_G]^{-1}(Y_G)^{\vee}/Y_G} / \omega_{Y_{P/U}^{\vee}/Y_G} \right),$$

where $\Gamma_{G_N} := \Gamma_G \cap N(\mathbb{Q})$, $Y_{P/U}^{\text{univ}}$ is the universal vector extension of the abelian scheme $Y_{P/U}/Y_G$, $\mathcal{C}^{\natural} := \Gamma_{G_N} \backslash (\mathcal{F}^0 \mathcal{V}^{(0)}|_{\tilde{Y}_G})$ is the largest trivial subbundle of $\omega_{[\pi_G]^{-1}(Y_G)^{\vee}/Y_G} / \omega_{Y_{P/U}^{\vee}/Y_G}$. Let $L^{\natural} := \text{unif}^{\natural}(\tilde{L}^{\natural})$. Then it is an irreducible algebraic subvariety of some fiber of $\mathcal{C}^{\natural} \rightarrow Y_G$, which is identified with the fiber of $\tilde{\mathcal{C}}^{\natural} \rightarrow \tilde{Y}_G$ containing $\tilde{L}^{\natural}$ under the map $\text{unif}^{\natural}$. Therefore $L_{Y_G}^{\natural} := \Gamma_{G_N} \backslash (\tilde{L}^{\natural} \times \tilde{Y}_G)$ is algebraic in $\mathcal{C}^{\natural}$. Then we have that

$$(5.3) \quad Y^{\natural} = Y \times_{Y_{P/U}} \left(Y_{P/U}^{\text{univ}} \times_{Y_G} L_{Y_G}^{\natural} \times_{Y_G} \mathbf{V}^{\dagger}|_{Y_G} \right),$$

where $\mathbf{V}^{\dagger}|_{Y_G} = \Gamma_{G_N} \backslash (\mathcal{F}^0 \mathcal{V}^{\dagger}|_{\tilde{Y}_G})$ is the subbundle of $\omega_{[\pi_G]^{-1}(Y_G)^{\vee}/Y_G} / \omega_{Y_{P/U}^{\vee}/Y_G}$ corresponding to the $G_{Q,\mathbb{R}}$ -submodule $V^{\dagger}$ of $V_{\mathbb{R}}$. Hence $Y^{\natural}$ is a closed algebraic irreducible subvariety of $S^{\natural}$.

Remark 5.7. *It is not hard to see that any subvariety $Y^{\natural}$ of $S^{\natural}$ of the form (5.3) is quasi-linear. Hence the discussion above proves that $Y^{\natural}$ is quasi-linear if and only if it is under the form (5.3).*

We are now ready to give the characterization of bi-algebraic subvarieties.

Theorem 5.8. *A subset $\tilde{Y}^{\natural}$ of $\mathcal{X}^{\natural+}$ (resp. a subset $Y^{\natural}$ of $S^{\natural}$) is bi-algebraic iff it is quasi-linear.*

Proof. We only need to prove that a bi-algebraic subset $\tilde{Y}^{\natural}$ of $\mathcal{X}^{\natural+}$ is quasi-linear. Up to replacing Γ by a finite subgroup we assume that $\Gamma = \Gamma_W \rtimes \Gamma_G$. Then $S_{P/U}^{\natural}$ is the universal vector extension of the abelian scheme $S_{P/U}$ over S_G . Use the notation of Notation 5.3.

Denote by $\Gamma_{Y^{\natural}} := \text{Im}(\pi_1(Y^{\natural, \text{sm}}) \rightarrow \pi_1(S^{\natural}) = \Gamma)$ and by $N := (\Gamma_{Y^{\natural}}^{\text{Zar}})^{\circ}$. Then for any point $\tilde{y}^{\natural} \in \tilde{Y}^{\natural}$, we have $\Gamma_{Y^{\natural}} \cdot \tilde{y}^{\natural} \subset \tilde{Y}^{\natural}$. So the bi-algebraicity of $Y^{\natural}$ implies

$$(5.4) \quad N(\mathbb{R})^+ W_N(\mathbb{C}) \tilde{y}^{\natural} \subset \tilde{Y}^{\natural} \text{ for any point } \tilde{y}^{\natural} \in \tilde{Y}^{\natural}.$$

By the characterization of bi-algebraic subsets for connected mixed Shimura varieties ([16, Corollary 8.3]), $\tilde{Y}$ is a weakly special subset of $\varphi(\mathcal{X}^+)$: $\tilde{Y} = N(\mathbb{R})^+ U_N(\mathbb{C}) \tilde{y}$ for some $\tilde{y} \in \tilde{Y}$ and N is a normal subgroup of Q , where $(Q, \mathcal{Y}^+)$ is the smallest connected mixed Shimura subdatum of $(P, \varphi(\mathcal{X}^+))$ such that $\tilde{Y} \subset \mathcal{Y}^+$. Then $Y_{P/U} \rightarrow Y_G$ is an abelian scheme and $S_{P/U}^{\natural}|_{Y_{P/U}} := [\pi_{P/U}^{\natural}]^{-1}(Y_{P/U})$ is a vector extension of $Y_{P/U}$ over Y_G .

Now (5.4) implies

$$\tilde{Y}^{\natural} = \tilde{Y} \times_{\tilde{Y}_{P/U}} \tilde{Y}_{P/U}^{\natural} \subset \varphi(\mathcal{X}^+) \times_{\varphi(\mathcal{X}^+)_{P/U}} \mathcal{X}_{P/U}^{\natural+}$$

and

$$(5.5) \quad \tilde{Y}_{P/U}^{\natural} = \tilde{Y}_{P/U}^{\text{univ}} \times_{\tilde{Y}_G} \tilde{Y}_{P/U}^{\natural, s}$$

where $\tilde{Y}_{P/U}^{\text{univ}}$ is as in (5.2).

Use the notation in (5.2). Let $V^{\text{nt}} := V_{\mathbb{R}}/(V_{N, \mathbb{R}} \oplus V^{(0)})$. Consider the vector bundle

$$\mathcal{F}^0 \mathcal{V}^{\text{nt}}|_{\tilde{Y}_G} = \mathcal{X}_{P/U}^{\natural+}|_{\tilde{Y}_{P/U}}/(\tilde{Y}_{P/U}^{\text{univ}} \times_{\tilde{Y}_G} \tilde{\mathcal{C}}^{\natural})$$

and the automorphic quotient-bundle

$$\mathbf{V}^{\text{nt}}|_{Y_G} := \Gamma_{G_N} \backslash (\mathcal{F}^0 \mathcal{V}^{\text{nt}}|_{\tilde{Y}_G})$$

of $\omega_{[\pi_G]^{-1}(Y_G)^{\vee}/Y_G}/\omega_{Y_{P/U}^{\vee}/Y_G}$. The equation (5.4) tells us that

$$(5.6) \quad \tilde{Y}_{P/U}^{\natural, s} = (\tilde{L}^{\natural} \times \tilde{Y}_G) \times_{\tilde{Y}_G} \tilde{Y}_{P/U}^{\natural, \text{extr}}$$

for some splitting $V_{\mathbb{R}}/V_{N, \mathbb{R}} = V^{(0)} \oplus V^{\text{nt}}$ as G_Q -modules, where $\tilde{L}^{\natural}$ is an irreducible algebraic subvariety of some fiber of $\tilde{\mathcal{C}}^{\natural} \rightarrow \tilde{Y}_G$ (which is isomorphic to $\mathbb{C}^k$ for some k) and $\tilde{Y}_{P/U}^{\natural, \text{extr}} \subset \mathcal{F}^0 \mathcal{V}^{\text{nt}}|_{\tilde{Y}_G}$ which surjects to $\tilde{Y}_G$. We have furthermore

$$(5.7) \quad \tilde{Y}_{P/U}^{\natural, \text{extr}} = G_N(\mathbb{R})^+ (\tilde{Y}_{P/U}^{\natural, \text{extr}})_{\tilde{y}_{G,0}}$$

We divide the rest of the proof into two steps.

Step I Prove that under the identification $\mathcal{F}^0 \mathcal{V}^{\text{nt}}|_{\tilde{Y}_G} = V^{\text{nt}, \vee}(\mathbb{R}) \times \tilde{Y}_G$ we have $(\tilde{Y}_{P/U}^{\natural, \text{extr}})_{\tilde{y}_G} = (\tilde{Y}_{P/U}^{\natural, \text{extr}})_{\tilde{y}_{G,0}}$ as subsets of $V^{\text{nt}, \vee}(\mathbb{R})$ for any $\tilde{y}_G \in \tilde{Y}_G$.

The theory of automorphic bundles tells us that the map

$$(5.8) \quad g: \mathbb{C}^l = \left(\mathcal{F}^0 \mathcal{V}^{\text{nt}}|_{\tilde{Y}_G} \right)_{\tilde{y}_{G,0}} \rightarrow \mathbb{C}^l = \left(\mathcal{F}^0 \mathcal{V}^{\text{nt}}|_{\tilde{Y}_G} \right)_{g\tilde{y}_{G,0}}$$

is algebraic (and in fact linear) for any $g \in G_N(\mathbb{R})^+$. Consider the map $G_N(\mathbb{R})^+ \rightarrow \text{Chow}(\mathbb{C}^l)$, $g \mapsto g \cdot (\tilde{Y}_{P/U}^{\natural, \text{extr}})_{\tilde{y}_{G,0}}$. This map descends to $\tilde{Y}_G = G_N(\mathbb{R})^+/\text{Stab}(\tilde{y}_{G,0}) \rightarrow \text{Chow}(\mathbb{C}^l)$ by (5.7), which is holomorphic because $\tilde{Y}_{P/U}^{\natural, \text{extr}}$ is complex analytic (in fact it is algebraic). Again by (5.7) this map descends to a holomorphic map $Y_G \rightarrow \text{Chow}(\mathbb{C}^l)$.

Lemma 5.9. *The image of the map $Y_G \rightarrow \text{Chow}(\mathbb{C}^l)$ is contained $\mathbb{C}^n$ for some $n \in \mathbb{Z}_{>0}$.*

Assume Lemma 5.9. This holomorphic map is then constant unless the boundary of the minimal compactification of Y_G is of codimension 1. Now (5.7) concludes this step when the boundary of the minimal compactification of Y_G has codimension at least 2.

Now assume that the boundary of the minimal compactification of Y_G has codimension 1. But Y_G is a weakly special subvariety of a Shimura variety, and hence itself is a locally symmetric variety. Therefore the minimal compactification of X_G is of codimension 1 if and only if Y_G is isomorphic (as an algebraic variety) to a modular curve (and $G_N = \text{SL}_2$ or PGL_2). But then any abelian scheme over Y_G with trivial constant part is a fiber product of the universal elliptic curve $\mathcal{E}$ over Y_G . Now the projection of $Y_{P/U}^{\natural, \text{extr}}$ to each factor $\omega_{\mathcal{E}/Y_G}$ is either the whole space or a section of $\omega_{\mathcal{E}/Y_G} \rightarrow Y_G$ invariant under $G_N(\mathbb{R})^+$ (which must be $\text{SL}_2(\mathbb{R})$ or $\text{PGL}_2(\mathbb{R})$). But any section of $\omega_{\mathcal{E}/Y_G} \rightarrow Y_G$ invariant under $G_N(\mathbb{R})^+$ corresponds to a modular form of level 1

and weight 2, so it must be 0. So the projection of $Y_{P/U}^{\natural, \text{extr}}$ to each factor $\omega_{\mathcal{E}/Y_G}$ is either the whole space or the zero section of $\omega_{\mathcal{E}/Y_G} \rightarrow Y_G$. Now we can conclude this step.

Proof of Lemma 5.9. It suffices to prove the claim for the image of $\tilde{Y}_G \rightarrow \text{Chow}(\mathbb{C}^l)$. Consider the action of $G_N(\mathbb{R})^+$ on $\mathbb{C}^l$ given by (5.8). For simplicity denote by $F := (\tilde{Y}_{P/U}^{\natural, \text{extr}})_{\tilde{y}_{G,0}}$, then it suffices to prove the claim for $G_N(\mathbb{R})^+/\text{Stab}_{G_N(\mathbb{R})^+}(F)$. Note that (5.7) implies

$$\text{Stab}_{G_N(\mathbb{R})^+}(\tilde{y}_{G,0}) \subset \text{Stab}_{G_N(\mathbb{R})^+}(F).$$

Let $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$ be the Cartan decomposition of $\mathfrak{g} := \text{Lie}(G_{N,\mathbb{R}})$ induced by $\tilde{y}_{G,0}$, then

$$\text{Lie}(\text{Stab}_{G_N(\mathbb{R})^+}(F)) = \mathfrak{k} + \mathfrak{p}_F$$

for some subalgebra $\mathfrak{p}_F$ of $\mathfrak{p}$. Therefore $G_N(\mathbb{R})^+/\text{Stab}_{G_N(\mathbb{R})^+}(F)$ is a quotient of

$$\tilde{X}_G = G_N(\mathbb{R})^+/\text{Stab}_{G_N(\mathbb{R})^+}(\tilde{y}_{G,0}) \subset \mathfrak{p} \simeq \mathbb{C}^m$$

for some $m \in \mathbb{Z}_{>0}$. Hence $G_N(\mathbb{R})^+/\text{Stab}_{G_N(\mathbb{R})^+}(F) \subset \mathfrak{p}/\mathfrak{p}_F \simeq \mathbb{C}^n$ for some $n \in \mathbb{Z}_{>0}$. $\square$

Step II Prove that $\tilde{Y}_{P/U}^{\natural, \text{extr}} = \mathcal{F}^0 \mathcal{V}^\dagger|_{\tilde{Y}_G}$ for some $G_{Q,\mathbb{R}}$ -submodule $V^\dagger$ of V^{nt} . Hence the theorem follows from (5.5), (5.6) and (5.7).

By Step I, we have

$$\Gamma_{G_N}(\tilde{Y}_{P/U}^{\natural, \text{extr}})_{\tilde{y}_{G,0}} \subset (\tilde{Y}_{P/U}^{\natural, \text{extr}})_{\tilde{y}_{G,0}}.$$

But $(\tilde{Y}_{P/U}^{\natural, \text{extr}})_{\tilde{y}_{G,0}}$ is algebraic, so

$$G_N(\mathbb{R})^+(\tilde{Y}_{P/U}^{\natural, \text{extr}})_{\tilde{y}_{G,0}} = (\tilde{Y}_{P/U}^{\natural, \text{extr}})_{\tilde{y}_{G,0}}.$$

Take any simple $G_{Q,\mathbb{R}}$ -quotient module V' of V^{nt} and consider $p': \mathcal{F}^0 \mathcal{V}^{\text{nt}}|_{\tilde{Y}_G} \rightarrow \mathcal{F}^0 \mathcal{V}'|_{\tilde{Y}_G}$. Then

$$G_N(\mathbb{R})^+ p'(\tilde{Y}_{P/U}^{\natural, \text{extr}})_{\tilde{y}_{G,0}} = p'(\tilde{Y}_{P/U}^{\natural, \text{extr}})_{\tilde{y}_{G,0}}.$$

Therefore $p'(\tilde{Y}_{P/U}^{\natural, \text{extr}})_{\tilde{y}_{G,0}}$ contains a $G_N(\mathbb{R})^+$ -submodule of $V'^{\vee}$. However the complex structure on $V^{\text{nt}, \vee}$ is given by its $G_{Q,\mathbb{R}}$ -module structure, and hence $p'(\tilde{Y}_{P/U}^{\natural, \text{extr}})_{\tilde{y}_{G,0}}$ contains a $G_Q(\mathbb{R})$ -submodule of $V'^{\vee}$, which is either 0 or $V'^{\vee}$. To sum it up, there exists a $G_{Q,\mathbb{R}}$ -submodule $V^\dagger$ of V^{nt} such that $(\tilde{Y}_{P/U}^{\natural, \text{extr}})_{\tilde{y}_{G,0}} = V^{\dagger, \vee}(\mathbb{R})$. Now (5.7) yields the conclusion. $\square$

5.3. Weakly special subvarieties. We close this section by defining weakly special subvarieties of $S^\natural$ and explain its relation with quasi-linear subvarieties. Note that for mixed Shimura varieties they coincide ([16, Corollary 8.3]).

Definition 5.10. (1) A subset $\tilde{Y}^\natural$ of $\mathcal{X}^{\natural+}$ is said to be **weakly special** if there exists a diagram in the category $\underline{\text{EMSD}}$

$$(R, \mathcal{Z}^{\natural+}) \xleftarrow{\varphi} (Q, \mathcal{Y}^{\natural+}) \xrightarrow{i} (P, \mathcal{X}^{\natural+})$$

and a point $\tilde{z}^\natural \in \mathcal{Z}^{\natural+}$ such that $\tilde{Y}^\natural$ is a connected component of $i(\varphi^{-1}(\tilde{z}^\natural))$.

(2) A subvariety $Y^\natural$ of $S^\natural$ is said to be **weakly special** if it is the image of some weakly special subset of $\mathcal{X}^{\natural+}$ under $\text{unif}^\natural$.

The geometric meaning of weakly special subvarieties is very clear: a subvariety $Y^{\natural}$ is weakly special in $S^{\natural}$ if and only if:

- $Y^{\natural} = Y \times_{Y_{P/U}} Y_{P/U}^{\natural}$.
- Y is weakly special. In particular $Y_{P/U}/Y_G$ is an abelian scheme.
- $Y_{P/U}^{\natural} = Y_{P/U}^{\text{univ}}$ where $Y_{P/U}^{\text{univ}}$ is the universal vector extension of $Y_{P/U}/Y_G$.

Now it is clear that every weakly special subvariety of $S^{\natural}$ is quasi-linear. On the other hand for any quasi-linear

$$\tilde{Y}^{\natural} = \tilde{Y} \times_{\tilde{Y}_{P/U}} \left(\tilde{Y}_{P/U}^{\text{univ}} \times_{\tilde{Y}_G} (\tilde{L}^{\natural} \times \tilde{Y}_G) \times_{\tilde{Y}_G} \mathcal{F}^0 \mathcal{V}^{\dagger}|_{\tilde{Y}_G} \right), \quad Y^{\natural} = Y \times_{Y_{P/U}} \left(Y_{P/U}^{\text{univ}} \times_{Y_G} L_{Y_G}^{\natural} \times_{Y_G} \mathbf{V}^{\dagger}|_{Y_G} \right),$$

we denote by $(\tilde{Y}^{\natural})^{\text{ws}} = \tilde{Y} \times_{\tilde{Y}_{P/U}} (\tilde{Y}_{P/U}^{\text{univ}} \times_{\tilde{Y}_G} \underline{0})$, $(Y^{\natural})^{\text{ws}} := Y \times_{Y_{P/U}} (Y_{P/U}^{\text{univ}} \times_{Y_G} \underline{0})$ and by

$$(5.9) \quad pr_{\tilde{Y}^{\natural}}^{\text{ws}}: \tilde{Y}^{\natural} \rightarrow (\tilde{Y}^{\natural})^{\text{ws}} \quad \text{and} \quad [pr]_{Y^{\natural}}^{\text{ws}}: Y^{\natural} \rightarrow (Y^{\natural})^{\text{ws}}$$

the natural projections (here we identify $\tilde{Y}_{P/U}^{\text{univ}} \times_{\tilde{Y}_G} \underline{0}$ with $\tilde{Y}_{P/U}^{\text{univ}}$ and $Y_{P/U}^{\text{univ}} \times_{Y_G} \underline{0}$ with $Y_{P/U}^{\text{univ}}$).

We can also define linear parts for any quasi-linear $\tilde{Y}^{\natural}$ and $Y^{\natural}$ as above: denote by $(\tilde{Y}^{\natural})^{\text{lin}} := \tilde{Y} \times_{\tilde{Y}_{P/U}} (\tilde{Y}_{P/U}^{\text{univ}} \times_{\tilde{Y}_G} \underline{0} \times_{\tilde{Y}_G} \mathcal{F}^0 \mathcal{V}^{\dagger}|_{\tilde{Y}_G})$ and $(Y^{\natural})^{\text{lin}} := Y \times_{Y_{P/U}} (Y_{P/U}^{\text{univ}} \times_{Y_G} \underline{0} \times_{Y_G} \mathbf{V}^{\dagger}|_{Y_G})$ and by

$$(5.10) \quad pr_{\tilde{Y}^{\natural}}^{\text{lin}}: \tilde{Y}^{\natural} \rightarrow (\tilde{Y}^{\natural})^{\text{lin}} \quad \text{and} \quad [pr]_{Y^{\natural}}^{\text{lin}}: Y^{\natural} \rightarrow (Y^{\natural})^{\text{lin}}$$

the natural projections (here we identify $\tilde{Y}_{P/U}^{\text{univ}} \times_{\tilde{Y}_G} \underline{0} \times_{\tilde{Y}_G} \mathcal{F}^0 \mathcal{V}^{\dagger}|_{\tilde{Y}_G}$ with $\tilde{Y}_{P/U}^{\text{univ}} \times_{\tilde{Y}_G} \mathcal{F}^0 \mathcal{V}^{\dagger}|_{\tilde{Y}_G}$ and $Y_{P/U}^{\text{univ}} \times_{Y_G} \underline{0} \times_{Y_G} \mathbf{V}^{\dagger}|_{Y_G}$ with $Y_{P/U}^{\text{univ}} \times_{Y_G} \mathbf{V}^{\dagger}|_{Y_G}$). Note that (5.9) and (5.10) coincide if $Y_G := [\pi^{\natural}](Y^{\natural})$ is a point.

6. THE AX LOGARITHMIQUE THEOREM

We prove in this section a transcendence theorem. We start with a criterion of bi-algebraicity, which will also be used in the next section.

Let $S^{\natural}$ be a connected enlarged mixed Shimura variety associated with $(P, \mathcal{X}^{\natural+})$ and let $\text{unif}^{\natural}: \mathcal{X}^{\natural+} \rightarrow S^{\natural}$ be the uniformization. Use Notation 5.3. Let $\tilde{Y}^{\natural}$ be a subset of $\mathcal{X}^{\natural+}$ such that $\tilde{Y}$ is weakly special, and hence $\tilde{Y} = N(\mathbb{R})^+ U_N(\mathbb{C}) \tilde{y}$ for some subgroup N of P and some point $\tilde{y} \in \varphi(\mathcal{X}^+)$. Let $\tilde{Y}_{P/U}^{\text{univ}}$ be as in (5.2).

Lemma 6.1. *Let $\tilde{Y}^{\natural}$ be an irreducible algebraic subset of $\mathcal{X}^{\natural+}$ such that*

- (1) $\tilde{Y}^{\natural} = \tilde{Y} \times_{\tilde{Y}_{P/U}} \tilde{Y}_{P/U}^{\natural} \subset \varphi(\mathcal{X}^+) \times_{\varphi(\mathcal{X}^+)_{P/U}} \tilde{\mathcal{X}}_{P/U}^{\natural+}$;
- (2) $\tilde{Y}$ is weakly special (and we use the notations above);
- (3) $\tilde{Y}^{\text{univ}} \subset \tilde{\mathcal{X}}_{P/U}^{\natural}$;
- (4) $\tilde{Y}_{P/U}^{\natural} = G_N(\mathbb{R})^+ (\tilde{Y}_{P/U}^{\natural})_{\tilde{y}_G}$ for one (and hence any) point $\tilde{y}_G \in \tilde{Y}_G$.

Then $\tilde{Y}^{\natural}$ is bi-algebraic.

Proof. It suffices to prove that $\tilde{Y}_{P/U}^{\natural}$ is bi-algebraic by condition (1) and (2). Rewrite (??)

$$(6.1) \quad 0 \rightarrow \mathcal{F}^0 \mathcal{V} \rightarrow \mathcal{X}_{P/U}^{\natural+} = \mathcal{V} \xrightarrow{\pi_{P/U}^{\natural}} \varphi(\mathcal{X}^+)_{P/U} = \mathcal{V}/\mathcal{F}^0 \mathcal{V} \rightarrow 0.$$

We have

$$(6.2) \quad S_{P/U}^{\natural}|_{Y_{P/U}} = Y_{P/U}^{\text{univ}} \times_{Y_G} \left(\omega_{[\pi_G]^{-1}(X_Y)^{\vee}/Y_G} / \omega_{Y_{P/U}^{\vee}/Y_G} \right),$$

where $Y_{P/U}^{\text{univ}}$ is the universal vector extension of the abelian scheme $Y_{P/U}/Y_G$. Let $\tilde{Y}_{P/U}^{\text{univ}}$ be defined by (compatible with (6.1))

$$0 \rightarrow \mathcal{F}^0 \mathcal{V}_N|_{\tilde{Y}_G} \rightarrow \tilde{Y}_{P/U}^{\text{univ}} \rightarrow \tilde{Y}_{P/U} \rightarrow 0.$$

Then the uniformization of (6.2) is

$$(6.3) \quad \tilde{\mathcal{X}}_{P/U}^{\natural+}|_{\tilde{Y}_{P/U}} (= \pi_{P/U}^{\natural-1}(\tilde{Y}_{P/U})) = \tilde{Y}_{P/U}^{\text{univ}} \times_{\tilde{Y}_G} \mathcal{F}^0 \mathcal{V}^s|_{\tilde{Y}_G}$$

for some G_Q -submodule V^s of V such that $V = V_N \oplus V^s$.

Now condition (3) implies

$$\tilde{Y}_{P/U}^{\natural} = \tilde{Y}_{P/U}^{\text{univ}} \times_{\tilde{Y}_G} \tilde{Y}_{P/U}^{\natural,s}$$

and hence

$$Y_{P/U}^{\natural} = Y_{P/U}^{\text{univ}} \times_{Y_G} \text{unif}_{P/U}^{\natural,s}(\tilde{Y}_{P/U}^{\natural,s}).$$

Hence it suffices to prove that $\text{unif}_{P/U}^{\natural,s}(\tilde{Y}_{P/U}^{\natural,s})$ is a closed algebraic subvariety of $\omega_{[\pi_G]^{-1}(Y_G)^{\vee}/Y_G} / \omega_{Y_{P/U}^{\vee}/Y_G}$.

Consider the diagram

$$\begin{array}{ccc} \mathcal{F}^0 \mathcal{V}^s|_{\tilde{Y}_G} & \xrightarrow{\text{unif}_{P/U}^{\natural,s}} & \omega_{[\pi_G]^{-1}(Y_G)^{\vee}/Y_G} / \omega_{Y_{P/U}^{\vee}/Y_G} \\ \downarrow & & \downarrow \\ \tilde{Y}_G & \xrightarrow{\text{unif}_G} & Y_G \end{array}$$

The restriction of $\text{unif}_{P/U}^{\natural,s}$ to any fiber

$$\left(\mathcal{F}^0 \mathcal{V}^s|_{\tilde{Y}_G} \right)_{\tilde{y}_G} \rightarrow \left(\omega_{[\pi_G]^{-1}(Y_G)^{\vee}/Y_G} / \omega_{Y_{P/U}^{\vee}/Y_G} \right)_{y_G},$$

where $\tilde{y}_G \in \tilde{Y}_G$ and $y_G = \text{unif}_G(\tilde{y}_G)$, is the identity map. So by [21, Theorem 1.9], there exists a fundamental domain $\mathfrak{F}$ of $\text{unif}_G: \tilde{Y}_G \rightarrow Y_G$ such that

$$\text{unif}_{P/U}^{\natural,s}|_{\mathfrak{F}} \text{ is definable in the o-minimal theory } \mathbb{R}_{\text{an,exp}}.$$

Therefore Condition (4) implies that $\text{unif}_{P/U}^{\natural,s}(\tilde{Y}_{P/U}^s)$ is closed in the usual topology (and hence a complex analytic subvariety of $\omega_{[\pi_G]^{-1}(Y_G)^{\vee}/Y_G} / \omega_{Y_{P/U}^{\vee}/Y_G}$) and is definable in the o-minimal theory $\mathbb{R}_{\text{an,exp}}$ (because $\text{unif}_{P/U}^{\natural,s}(\tilde{Y}_{P/U}^s) = \text{unif}_{P/U}^{\natural,s}(\tilde{Y}_{P/U}^s \cap \mathcal{F}^0 \mathcal{V}^s|_{\mathfrak{F}})$ and $\tilde{Y}_{P/U}^s$ is algebraic). Hence $Y_{P/U}^{\natural,s} := \text{unif}_{P/U}^{\natural,s}(\tilde{Y}_{P/U}^s)$ is a closed algebraic subvariety of $\omega_{[\pi_G]^{-1}(Y_G)^{\vee}/Y_G} / \omega_{Y_{P/U}^{\vee}/Y_G}$ by the Peterzil-Starchenko o-minimal GAGA theorem [28, Corollary 4.5]. $\square$

Corollary 6.2. *Let $\tilde{Y}^{\natural}$ be an irreducible algebraic subset of $\mathcal{X}^{\natural+}$ such that $\tilde{Y}$ is weakly special (and hence we can use the notation above Lemma 6.1). Assume $\tilde{Y}^{\natural} = N(\mathbb{R})^+ W_N(\mathbb{C})(\tilde{Y}^{\natural})_{\tilde{y}_G}$ for one (and hence any) point $\tilde{y}_G \in \tilde{Y}_G$. Then $\tilde{Y}^{\natural}$ is bi-algebraic.*

Proof. It is not hard to see that all the conditions of Lemma 3.6 are satisfied. $\square$

Theorem 6.3 (Ax logarithmique). *Let $S^{\natural}$ be a connected enlarged mixed Shimura variety associated with $(P, \mathcal{X}^{\natural+})$ with $\text{unif}^{\natural}: \mathcal{X}^{\natural+} \rightarrow S^{\natural}$. Let $Z^{\natural}$ be an irreducible subvariety of $S^{\natural}$. Let $\tilde{Z}^{\natural}$ be a complex analytic irreducible component of $\text{unif}^{\natural-1}(Z^{\natural})$ and let $\tilde{Z}^{\natural, \text{Zar}}$ be the Zariski closure of $\tilde{Z}^{\natural}$ in $\mathcal{X}^{\natural+}$. Then $\tilde{Z}^{\natural, \text{Zar}}$ is quasi-linear.*

Proof. Up to replacing Γ by a finite subgroup we assume that $\Gamma = \Gamma_W \rtimes \Gamma_G$. Then $S_{P/U}^{\natural}$ is the universal vector extension of the abelian scheme $S_{P/U}$ over S_G . Use the notation of Notation 5.3.

Denote by $\Gamma_{Z^{\natural}} := \text{Im}(\pi_1(Z^{\natural, \text{sm}}) \rightarrow \pi_1(S^{\natural}) = \Gamma)$ and by $N := (\Gamma_{Z^{\natural}}^{\text{Zar}})^{\circ}$. Then for any point $\tilde{y}^{\natural} \in \tilde{Z}^{\natural}$, we have $\Gamma_{Z^{\natural}} \cdot \tilde{y}^{\natural} \subset \tilde{Z}^{\natural}$. So

$$(6.4) \quad N(\mathbb{R})^+ W_N(\mathbb{C}) \tilde{z}^{\natural} \subset \tilde{Y}^{\natural, \text{Zar}} \text{ for any point } \tilde{z}^{\natural} \in \tilde{Z}^{\natural, \text{Zar}}.$$

Denote for simplicity $\tilde{Y}^{\natural} := \tilde{Z}^{\natural, \text{Zar}}$ and use the notation of Notation 5.3. By Ax logarithmique for connected mixed Shimura varieties ([16, Theorem 8.1]), $\tilde{Y}$ is a weakly special subset of $\varphi(\mathcal{X}^+)$: $\tilde{Y} = N(\mathbb{R})^+ U_N(\mathbb{C}) \tilde{z}$ for some $\tilde{z} \in \tilde{Y}$ and N is a normal subgroup of Q , where $(Q, \mathcal{Y}^+)$ is the smallest connected mixed Shimura subdatum of $(P, \varphi(\mathcal{X}^+))$ such that $\tilde{Z} \subset \mathcal{Y}^+$. But we have $\tilde{Y}^{\natural} = N(\mathbb{R})^+ W_N(\mathbb{C}) (\tilde{Y}^{\natural})_{\tilde{y}_G}$ for one (and hence any) point $\tilde{y}_G \in \tilde{Y}_G$. Now it suffices to apply Corollary 6.2. $\square$

7. THE AX-LINDEMANN THEOREM

Let $S^{\natural} = \Gamma \backslash \mathcal{X}^{\natural+}$ be a connected enlarged mixed Shimura variety associated with $(P, \mathcal{X}^{\natural+})$ with $\text{unif}^{\natural}: \mathcal{X}^{\natural+} \rightarrow S^{\natural}$. We hereby give four equivalent statements for the Ax-Lindemann(-Weierstraß) theorem and explain their equivalence. The proof will be given in the Appendix.

Theorem 7.1. *Let $Y^{\natural}$ be an irreducible subvariety of $S^{\natural}$ and let $\tilde{Z}^{\natural}$ be an irreducible algebraic subset of $\mathcal{X}^{\natural+}$ contained in $\text{unif}^{\natural-1}(Y^{\natural})$, maximal for these properties. Then $\tilde{Z}^{\natural}$ is quasi-linear.*

The next statement we give shall be called the *semi-algebraic form of Ax-Lindemann*. Recall that a connected semi-algebraic subset of $\mathcal{X}^{\natural+}$ is called **irreducible** if its $\mathbb{R}$ -Zariski closure in $W(\mathbb{C}) \times \mathcal{X}_G^{\vee}$ is an irreducible real algebraic variety. Note that any connected semi-algebraic subset of $\mathcal{X}^{\natural+}$ has only finitely many irreducible components.

Theorem 7.2. *Let $Y^{\natural}$ be an irreducible subvariety of $S^{\natural}$ and let $\tilde{Z}^{\natural}$ be a connected irreducible semi-algebraic subset of $\mathcal{X}^{\natural+}$ contained in $\text{unif}^{\natural-1}(Y^{\natural})$, maximal for these properties. Then $\tilde{Z}^{\natural}$ is complex analytic and each complex analytic irreducible component of $\tilde{Z}^{\natural}$ is quasi-linear.*

The equivalence of Theorem 7.1 and Theorem 7.2 follows easily from [30, Lemma 4.1], which claims that **maximal connected irreducible semi-algebraic subsets of $\mathcal{X}^{\natural+}$ which are contained in $\text{unif}^{\natural-1}(Y^{\natural})$ are all algebraic in the sense of Definition 5.4** (there is a typo in the proof of [30, Lemma 4.1]: $\mathbb{C}^{2n}$ should be $\mathbb{C}^n$).

The next two forms of Ax-Lindemann have more “equidistributional” taste.

Theorem 7.3. *Let $\tilde{Z}^{\natural}$ be an irreducible algebraic subset of $\mathcal{X}^{\natural+}$. Then $\text{unif}^{\natural}(\tilde{Z}^{\natural})^{\text{Zar}}$ is quasi-linear.*

Theorem 7.4. *Let $\tilde{Z}^{\natural}$ be a semi-algebraic subset of $\mathcal{X}^{\natural+}$. Then every irreducible component of $\text{unif}^{\natural}(\tilde{Z}^{\natural})^{\text{Zar}}$ is quasi-linear.*

Let us explain now why Theorem 7.1 implies Theorem 7.3. Let $\tilde{Z}^\natural$ be as in Theorem 7.3. Let $Y^\natural := \text{unif}^\natural(\tilde{Z}^\natural)^{\text{Zar}}$ and let $\tilde{W}^\natural$ be an irreducible algebraic subset of $\mathcal{X}^{\natural+}$ which contains $\tilde{Z}^\natural$ and is contained in $\text{unif}^{\natural-1}(Y^\natural)$, maximal for these properties. Such a $\tilde{W}^\natural$ exists by, for example, dimension reason. Then $Y^\natural = \text{unif}^\natural(\tilde{W}^\natural)$ and $\tilde{W}^\natural$ is an irreducible algebraic subset of $\mathcal{X}^{\natural+}$ which is contained in $\text{unif}^{\natural-1}(Y^\natural)$, maximal for these properties. Theorem 7.1 then implies that $\tilde{W}^\natural$ is irreducible bi-algebraic. So $Y = \text{unif}^\natural(\tilde{W}^\natural)^{\text{Zar}} = \text{unif}^\natural(\tilde{W}^\natural)$ is quasi-linear. Theorem 7.2 implies Theorem 7.4 by a similar argument because any semi-algebraic subset of $\mathcal{X}^{\natural+}$ has only finitely many connected irreducible components.

Let us explain now why Theorem 7.3 implies Theorem 7.1. Let $Y^\natural$ and $\tilde{Z}^\natural$ be as in Theorem 7.1. Then Theorem 7.3 shows that $\text{unif}^\natural(\tilde{Z}^\natural)^{\text{Zar}}$ is an irreducible bi-algebraic subvariety of S , which we shall call $Y_0^\natural$. It is clear that $Y_0^\natural$ is a subvariety of $Y^\natural$. Let $\tilde{Y}_0^\natural$ be the complex analytic irreducible component of $\text{unif}^{\natural-1}(Y_0^\natural)$ containing $\tilde{Z}^\natural$. Then $\tilde{Y}_0^\natural$ is irreducible algebraic. But then the maximality assumption on $\tilde{Z}^\natural$ implies $\tilde{Z}^\natural = \tilde{Y}_0^\natural$. Hence $\tilde{Z}^\natural$ is quasi-linear. Theorem 7.4 implies Theorem 7.2 by a similar argument.

8. THE AX-SCHANUEL CONJECTURE

8.1. Formulation of the conjecture. For mixed Shimura varieties we have the following Ax-Schanuel conjecture: let S be a connected mixed Shimura variety associated with $(P, \mathcal{X}^+)$ and let $\text{unif}: \mathcal{X}^+ \rightarrow S$ be the uniformization. Let $pr_{\mathcal{X}^+}: \mathcal{X}^+ \times S \rightarrow \mathcal{X}^+$ and $pr_S: \mathcal{X}^+ \times S \rightarrow S$ be the natural projections.

Conjecture 8.1 (Ax-Schanuel for mixed Shimura varieties). *Let $\Delta \subset \mathcal{X}^+ \times S$ be the graph of unif . Let $\mathfrak{Z} = \text{graph}(\tilde{Z} \xrightarrow{\text{unif}} Z)$ be a complex analytic irreducible subvariety of Δ and let $\mathcal{B}$ be its Zariski closure in $\mathcal{X}^+ \times S$. Let F be the smallest bi-algebraic (i.e. weakly special) subvariety of S which contains $Z = pr_S(\mathfrak{Z})$. Then*

$$\dim \mathcal{B} - \dim \mathfrak{Z} \geq \dim F.$$

We explain this conjecture. First replace $\mathfrak{Z}$ by a complex analytic irreducible component of $\mathcal{B} \cap \Delta$. Then denote by $\tilde{X} := pr_{\mathcal{X}^+}(\mathcal{B}) = \tilde{Z}^{\text{Zar}}$ and $Y := pr_S(\mathcal{B}) = Z^{\text{Zar}}$. We have $\mathcal{B} \subset \tilde{X} \times Y$. Therefore Conjecture 8.1 implies

$$(8.1) \quad \dim \tilde{X} + \dim Y - \dim \tilde{Z} \geq \dim \mathcal{B} - \dim \mathfrak{Z} \geq \dim F.$$

On the other hand let $\tilde{Y}$ (resp. $\tilde{F}$) be the complex analytic irreducible component of $\text{unif}^{-1}(Y)$ (resp. of $\text{unif}^{-1}(F)$) containing $\tilde{Z}$, then $\tilde{Z}$ is a complex analytic irreducible component of $\tilde{X} \cap \tilde{Y}$ since $\mathfrak{Z}$ is a complex analytic irreducible component of $\mathcal{B} \cap \Delta$. Hence we always have

$$(8.2) \quad \dim \tilde{Z} \geq \dim \tilde{X} + \dim \tilde{Y} - \dim \tilde{F}.$$

Now (8.1) and (8.2) together imply

$$\dim \mathcal{B} = \dim \tilde{X} + \dim Y, \quad \dim \tilde{Z} = \dim \tilde{X} + \dim \tilde{Y} - \dim \tilde{F},$$

so Conjecture 8.1 is equivalent to:

- $\mathcal{B} = \tilde{X} \times Y$;
- $\tilde{X}$ and $\tilde{Y}$ intersect properly in $\tilde{F}$.

As we shall see, Ax logarithmique and Ax-Lindemann for mixed Shimura varieties are special cases of this conjecture. We wish to make a similar conjecture for enlarged mixed Shimura varieties. However the naive guess is NOT always true: if $\tilde{Z}^\natural$ is contained in a fiber of $\mathcal{X}^{\natural+} \rightarrow \varphi(\mathcal{X}^+)$, then $\text{unif}^\natural|_{\tilde{Z}^\natural}$ is the identity map for some $\mathbb{C}^n$, and so $\mathcal{Z}^\natural := \text{graph}(\tilde{Z}^\natural \xrightarrow{\text{unif}^\natural} Z^\natural)$ is algebraic. So $(\mathcal{Z}^\natural)^{\text{Zar}} = \mathcal{Z}^\natural \neq \tilde{Z}^\natural \times Z^\natural$ and $\dim(\mathcal{Z}^\natural)^{\text{Zar}} - \dim \mathcal{Z}^\natural = 0$.

Counterexamples to the naive guess as above exist due to the existence of those $F^\natural$ such that $(\Delta^\natural \cap (\tilde{F}^\natural \times F^\natural))^{\text{Zar}} \neq \tilde{F}^\natural \times F^\natural$, and the existence of such $F^\natural$ is due to the non-linear part of $F^\natural$. However evidences suggest that this should be the only obstacle. So we make the following conjecture: let $S^\natural$ be a connected enlarged mixed Shimura variety associated with $(P, \mathcal{X}^{\natural+})$ and let $\text{unif}^\natural: \mathcal{X}^{\natural+} \rightarrow S^\natural$ be the uniformization.

Conjecture 8.2 (Ax-Schanuel for enlarged mixed Shimura varieties). *Let $\Delta^\natural \subset \mathcal{X}^{\natural+} \times S^\natural$ be the graph of $\text{unif}^\natural$. Let $\mathcal{Z}^\natural = \text{graph}(\tilde{Z}^\natural \xrightarrow{\text{unif}^\natural} Z^\natural)$ be a complex analytic irreducible subvariety of $\Delta^\natural$. Let $F^\natural$ (resp. $\tilde{F}^\natural$) be the smallest bi-algebraic, or equivalently quasi-linear, subvariety of $S^\natural$ (resp. subset of $\mathcal{X}^{\natural+}$) which contains $Z^\natural$ (resp. $\tilde{Z}^\natural$). In particular $F^\natural = \text{unif}^\natural(\tilde{F}^\natural)$.*

(1) *Let $\tilde{X}^\natural := (\tilde{Z}^\natural)^{\text{Zar}}$ and $Y^\natural := (Z^\natural)^{\text{Zar}}$. Then*

$$\dim \tilde{X}^\natural + \dim Y^\natural - \dim \tilde{Z}^\natural \geq \dim F^\natural.$$

(2) *Let $\mathcal{B}^\natural := (\mathcal{Z}^\natural)^{\text{Zar}} \subset \mathcal{X}^{\natural+} \times S^\natural$. Let $[pr]_{F^\natural}^{\text{lin}}$ and $pr_{\tilde{F}^\natural}^{\text{lin}}$ be as in (5.10). Denote for simplicity by $\mathbf{pr}_{F^\natural}^{\text{lin}} := (pr_{\tilde{F}^\natural}^{\text{lin}}, [pr]_{F^\natural}^{\text{lin}}): \tilde{F}^\natural \times F^\natural \rightarrow (\tilde{F}^\natural)^{\text{lin}} \times (F^\natural)^{\text{lin}}$. Then*

$$\dim \mathbf{pr}_{F^\natural}^{\text{lin}}(\mathcal{B}^\natural) - \dim \mathbf{pr}_{F^\natural}^{\text{lin}}(\mathcal{Z}^\natural) \geq \dim (F^\natural)^{\text{lin}}$$

Remark 8.3. *The two parts of Conjecture 8.2 do not imply each other: the first part implies Ax logarithmique (Theorem 6.3) and Ax-Lindemann (Theorem 7.1) while the second part does not, and the second part implies Conjecture 8.1 while the first part does not.*

8.2. Ax-Schanuel (Conjecture 8.2) implies Ax logarithmique (Theorem 6.3).

Theorem 8.4. (1) *The Ax-Schanuel conjecture implies the Ax logarithmique theorem.*

(2) *The Ax-Schanuel conjecture is true if $Z^\natural$ is algebraic: it is implied by Ax logarithmique.*

Proof. For (1): Let $Z^\natural$ be an irreducible subvariety of $S^\natural$ and let $\tilde{Z}^\natural$ be a complex analytic irreducible component of $\text{unif}^{\natural-1}(Z^\natural)$. We want to prove that $\tilde{X}^\natural = \tilde{Z}^{\natural, \text{Zar}}$ is bi-algebraic assuming Conjecture 8.2. But the first part of Conjecture 8.2 implies (since $Y^\natural = Z^\natural$)

$$\dim \tilde{X}^\natural = \dim \tilde{X}^\natural + \dim Y^\natural - \dim \tilde{Z}^\natural \geq \dim \tilde{F}^\natural.$$

Therefore $\tilde{X}^\natural = \tilde{F}^\natural$ is bi-algebraic since $\tilde{X}^\natural \subset \tilde{F}^\natural$.

For (2): Ax logarithmique implies $\tilde{X}^\natural = \tilde{F}^\natural$. Hence the first part is proven. Now it suffices to prove $\mathbf{pr}_{F^\natural}^{\text{lin}}(\mathcal{B}^\natural) = (\tilde{F}^\natural)^{\text{lin}} \times [pr]_{F^\natural}^{\text{lin}}(Z^\natural)$.

The group $P(\mathbb{R})W(\mathbb{C})$ acts on $\mathcal{X}^{\natural+} \times S^\natural$ by its action on the first factor. Let $\Gamma_{Z^\natural} := \text{Im}(\pi_1(Z^\natural) \rightarrow \pi_1(S^\natural) = \Gamma)$ be the monodromy group of $Z^\natural$ and let $N := (\Gamma_{Z^\natural}^{\text{Zar}})^\circ$. Then for any point $\tilde{z}^\natural \in \tilde{Z}^\natural$, we have $\Gamma_{Z^\natural} \cdot \tilde{z}^\natural \subset \mathcal{Z}^\natural$. So $N(\mathbb{R})^+ W_N(\mathbb{C}) \tilde{z}^\natural \subset \mathcal{B}^\natural$. So $\mathbf{pr}_{F^\natural}^{\text{lin}}(\mathcal{B}^\natural)_{[pr]_{F^\natural}^{\text{lin}}(Z^\natural)} := \mathbf{pr}_{F^\natural}^{\text{lin}}(\mathcal{B}^\natural \cap pr_{S^\natural}^{-1}(Z^\natural))$ is bi-algebraic, and hence equals $(\tilde{F}^\natural)^{\text{lin}}$, for any $z^\natural \in Z^\natural$ by (the last paragraph of the proof of) Ax logarithmique. Now we are done. $\square$

8.3. Ax-Schanuel (Conjecture 8.2) implies Ax-Lindemann (Theorem 7.1).

Theorem 8.5. (1) *The Ax-Schanuel conjecture implies the Ax-Lindemann theorem.*

(2) *The Ax-Schanuel conjecture is true if $\tilde{Z}^\natural$ is algebraic: it is implied by Ax-Lindemann.*

Proof. For (1): Let $\tilde{Z}^\natural$ be an irreducible algebraic subset of $\mathcal{X}^{\natural+}$ and let $Z^\natural := \text{unif}^\natural(\tilde{Z}^\natural)$. We want to prove that $Y^\natural = Z^{\natural, \text{Zar}}$ is bi-algebraic. But the first part of Conjecture 8.2 implies (since $\tilde{X}^\natural = \tilde{Z}^\natural$)

$$\dim Y^\natural = \dim \tilde{X}^\natural + \dim Y^\natural - \dim \tilde{Z}^\natural \geq \dim F^\natural,$$

and therefore $Y^\natural = F^\natural$ is bi-algebraic since $Y^\natural \subset F^\natural$.

For (2): Ax-Lindemann implies $Y^\natural = F^\natural$. Hence the first part is proven. Now it suffices to prove $\mathbf{pr}_{F^\natural}^{\text{lin}}(\mathcal{B}^\natural) = \mathbf{pr}_{\tilde{F}^\natural}^{\text{lin}}(\tilde{\mathcal{Z}}^\natural) \times (F^\natural)^{\text{lin}}$.

We start by the case where $\mathbf{pr}_{\tilde{F}^\natural}^{\text{lin}}(\tilde{\mathcal{Z}}^\natural)$ is a curve. The Ax-Lindemann theorem implies $Z^{\natural, \text{Zar}} = F^\natural$. So $\mathbf{pr}_{F^\natural}^{\text{lin}}(\mathcal{B}^\natural) \subset \mathbf{pr}_{\tilde{F}^\natural}^{\text{lin}}(\tilde{\mathcal{Z}}^\natural) \times (F^\natural)^{\text{lin}}$ and surjects to $\mathbf{pr}_{\tilde{F}^\natural}^{\text{ws}}(\tilde{\mathcal{Z}}^\natural)$ and $(F^\natural)^{\text{lin}}$. Therefore either $\mathbf{pr}_{F^\natural}^{\text{lin}}(\mathcal{B}^\natural) = \mathbf{pr}_{\tilde{F}^\natural}^{\text{lin}}(\tilde{\mathcal{Z}}^\natural) \times (F^\natural)^{\text{lin}}$ or $\mathbf{pr}_{F^\natural}^{\text{lin}}(\mathcal{B}^\natural) \rightarrow \dim(F^\natural)^{\text{lin}}$ is quasi-finite. Suppose that we are in the latter case. Up to replacing $S^\natural$ by a finite cover we may assume that $\mathbf{pr}_{F^\natural}^{\text{lin}}(\mathcal{B}^\natural) \rightarrow (F^\natural)^{\text{lin}}$ is a bijection. Let $U^\natural$ be a non-empty open subvariety of $(F^\natural)^{\text{lin}}$ such that $\mathbf{pr}_{\tilde{F}^\natural}^{\text{lin}}(\tilde{\mathcal{Z}}^\natural)|_{U^\natural} := \mathbf{pr}_{\tilde{F}^\natural}^{\text{lin}}(\tilde{\mathcal{Z}}^\natural) \cap \text{unif}^{\natural-1}(U^\natural)$ is smooth. Now we define $i: U^\natural \rightarrow \mathbf{pr}_{\tilde{F}^\natural}^{\text{lin}}(\tilde{\mathcal{Z}}^\natural)|_{U^\natural}$, $s^\natural \mapsto \mathbf{pr}_{F^\natural}^{\text{lin}}(\mathcal{B}^\natural)_{s^\natural}$. This map is algebraic and bijective, and hence is an isomorphism of algebraic varieties since $\mathbf{pr}_{\tilde{F}^\natural}^{\text{lin}}(\tilde{\mathcal{Z}}^\natural)|_{U^\natural}$ is smooth. But $\text{unif}^\natural|_{\mathbf{pr}_{\tilde{F}^\natural}^{\text{lin}}(\tilde{\mathcal{Z}}^\natural)|_{U^\natural}}$, which is not algebraic, is the inverse of i . This is a contradiction. Hence we must have $\mathbf{pr}_{F^\natural}^{\text{lin}}(\mathcal{B}^\natural) = \mathbf{pr}_{\tilde{F}^\natural}^{\text{lin}}(\tilde{\mathcal{Z}}^\natural) \times (F^\natural)^{\text{lin}}$ when $\mathbf{pr}_{\tilde{F}^\natural}^{\text{lin}}(\tilde{\mathcal{Z}}^\natural)$ is a curve.

Now we turn to $\mathbf{pr}_{\tilde{F}^\natural}^{\text{lin}}(\tilde{\mathcal{Z}}^\natural)$ of arbitrary dimension. Let $\tilde{z}^\natural \in \mathbf{pr}_{\tilde{F}^\natural}^{\text{lin}}(\tilde{\mathcal{Z}}^\natural)$ and denote by $z^\natural := \text{unif}^\natural(\tilde{z}^\natural)$. Let $\tilde{C}^\natural$ be any algebraic curve in $\mathbf{pr}_{\tilde{F}^\natural}^{\text{lin}}(\tilde{\mathcal{Z}}^\natural)$ passing through $\tilde{z}^\natural$. By the last paragraph we have $\tilde{C}^\natural \subset \mathbf{pr}_{F^\natural}^{\text{lin}}(\mathcal{B}^\natural)_{z^\natural} := \mathbf{pr}_{F^\natural}^{\text{lin}}(\mathcal{B}^\natural \cap \text{pr}_{S^\natural}^{-1}(z^\natural))$. By varying $\tilde{C}^\natural$ we get $\mathbf{pr}_{\tilde{F}^\natural}^{\text{lin}}(\tilde{\mathcal{Z}}^\natural) \subset \mathbf{pr}_{F^\natural}^{\text{lin}}(\mathcal{B}^\natural)_{z^\natural}$. By varying $\tilde{z}^\natural$ we get $\mathbf{pr}_{\tilde{F}^\natural}^{\text{lin}}(\tilde{\mathcal{Z}}^\natural) \times [\mathbf{pr}_{F^\natural}^{\text{lin}}(Z^\natural)]^{\text{Zar}} \subset \mathbf{pr}_{F^\natural}^{\text{lin}}(\mathcal{B}^\natural) \subset \mathbf{pr}_{\tilde{F}^\natural}^{\text{lin}}(\tilde{\mathcal{Z}}^\natural) \times (F^\natural)^{\text{lin}}$

and we are done. $\square$

9. PROOF OF AX-SCHANUEL WHEN RESTRICTED TO A FIBER OF $S^\natural \rightarrow S_G$

Let $S^\natural$ be a connected enlarged mixed Shimura variety associated with $(P, \mathcal{X}^{\natural+})$. The goal of this section is to prove the Ax-Schanuel conjecture (Conjecture 8.2) for a fiber of $[\pi^\natural]: S^\natural \rightarrow S_G$. More concretely let $E^\natural$ be the fiber of the natural projection $S^\natural \rightarrow S_G$ over $b \in S_G$. The uniformization of $E^\natural$ is given by $\text{unif}^\natural: W(\mathbb{C}) \rightarrow E^\natural$. We prove

Theorem 9.1. *Let $\Delta^\natural \subset W(\mathbb{C}) \times E^\natural$ be the graph of $\text{unif}^\natural$. Let $\mathcal{Z}^\natural = \text{graph}(\tilde{Z}^\natural \xrightarrow{\text{unif}^\natural} Z^\natural)$ be a complex analytic irreducible subvariety of $\Delta^\natural$. Let $F^\natural$ (resp. $\tilde{F}^\natural$) be the smallest bi-algebraic, or equivalently quasi-linear, subvariety of $E^\natural$ (resp. subset of $W(\mathbb{C})$) which contains $Z^\natural$ (resp. $\tilde{Z}^\natural$). In particular $F^\natural = \text{unif}^\natural(\tilde{F}^\natural)$.*

(1) Let $\tilde{X}^{\natural} := (\tilde{Z}^{\natural})^{\text{Zar}}$ and $Y^{\natural} := (Z^{\natural})^{\text{Zar}}$. Then

$$\dim \tilde{X}^{\natural} + \dim Y^{\natural} - \dim \tilde{Z}^{\natural} \geq \dim F^{\natural}.$$

(2) Let $\mathcal{B}^{\natural} := (\mathcal{Z}^{\natural})^{\text{Zar}} \subset \mathcal{X}^{\natural+} \times S^{\natural}$. Let $[pr]_{F^{\natural}}^{\text{ws}}$ and $pr_{\tilde{F}^{\natural}}^{\text{ws}}$ be as in (5.9) (note that (5.9) and (5.10) coincide in this case). Denote for simplicity by $\mathbf{pr}_{F^{\natural}}^{\text{ws}} := (pr_{\tilde{F}^{\natural}}^{\text{ws}}, [pr]_{F^{\natural}}^{\text{ws}}): \tilde{F}^{\natural} \times F^{\natural} \rightarrow (\tilde{F}^{\natural})^{\text{ws}} \times (F^{\natural})^{\text{ws}}$. Then

$$\dim \mathbf{pr}_{F^{\natural}}^{\text{ws}}(\mathcal{B}^{\natural}) - \dim \mathbf{pr}_{F^{\natural}}^{\text{ws}}(\mathcal{Z}^{\natural}) \geq \dim (F^{\natural})^{\text{ws}}$$

9.1. Ax-Schanuel for E . In this subsection we go back to mixed Shimura variety setting. Let S be a connected mixed Shimura variety associated with the mixed Shimura datum $(P, \mathcal{X}^+)$. Let E be the fiber of the natural projection $S \rightarrow S_G$ over $b \in S_G$. Then the uniformization of E is given by $\text{unif}: W(\mathbb{R})U(\mathbb{C}) \rightarrow E$, where the complex structure on $W(\mathbb{R})U(\mathbb{C})$ is given by $W(\mathbb{C})/F_b^0 W(\mathbb{C})$. We prove Conjecture 8.1 for E . More concretely we prove

Theorem 9.2. *Let $\Delta \subset W(\mathbb{R})U(\mathbb{C}) \times E$ be the graph of unif . Let $\mathcal{Z}$ be a complex analytic irreducible subvariety of Δ and let $\mathcal{B}$ be its Zariski closure in $W(\mathbb{R})U(\mathbb{C}) \times E$. Let F be the smallest weakly special subvariety of E containing $\text{pr}_E(\mathcal{Z})$. Then*

$$\dim \mathcal{B} - \dim \mathcal{Z} \geq \dim F.$$

Without loss of generality we assume that $\mathcal{Z}$ is a complex analytic irreducible component of $\mathcal{B} \cap \Delta$. The following notation will be used only in this subsection: let Γ_F be the monodromy group of F , let W_F be its Zariski closure in W and let $U_F := W_F \cap U$ and $V_F := W_F/U_F$ (hence $\tilde{F} = W_F(\mathbb{R})U_F(\mathbb{C})\tilde{x}$ for any $\tilde{x} \in \tilde{Z}$), $\Delta_F := \text{graph}(\tilde{F} \xrightarrow{\text{unif}} F)$ and

$$\begin{array}{ccc} W(\mathbb{R})U(\mathbb{C}) & \xrightarrow{\pi} & V(\mathbb{R}) \\ \text{unif} \downarrow & & \downarrow \text{unif}_V \\ E & \xrightarrow{[\pi]} & A \end{array} \quad \text{and} \quad \begin{array}{ccc} \pi: W(\mathbb{R})U(\mathbb{C}) \times E & \xrightarrow{(\pi, [\pi])} & V(\mathbb{R}) \times A \\ \uparrow & & \uparrow \\ \mathcal{Z} & \longrightarrow & \mathcal{Z}_V := \pi(\mathcal{Z}) \end{array}$$

The group $W(\mathbb{R})U(\mathbb{C})$ acts on $W(\mathbb{R})U(\mathbb{C}) \times E$ by its action on the first factor. Let $\mathfrak{F}$ be a fundamental set for unif such that $\text{unif}|_{\mathfrak{F}}$ is definable and let $\mathfrak{F}_V := \pi(\mathfrak{F})$, which is then a fundamental set for unif_V .

9.1.1. Basic setup.

Definition 9.3. • For any definable subset $\tilde{D}$ of $W(\mathbb{R})U(\mathbb{C})$, define $\tilde{D}^{\text{rl}}$ to be the smallest subset of $W(\mathbb{R})U(\mathbb{C})$ of the form $W'(\mathbb{R})w$, where W' is a subgroup of $W_{\mathbb{R}}$ and $w \in W(\mathbb{R})$.
• For any definable subset $\mathcal{D}$ of $W(\mathbb{R})U(\mathbb{C}) \times E$, a positive integer r and $M = W'(\mathbb{R})w$ ($W' < W_{\mathbb{R}}$, $w \in W(\mathbb{R})$), define $G_r(\mathcal{D}, M)$ to be the set of points $\mathbf{d} \in \mathcal{D}$ around which $\mathcal{D}$ is regular of dimension r and such that $\tilde{D}^{\text{rl}} = M$, where $\tilde{D}$ is the projection of the irreducible component of $\mathcal{D}$ containing $\mathbf{d}$ under $W(\mathbb{R})U(\mathbb{C}) \times E \rightarrow W(\mathbb{R})U(\mathbb{C})$.

Define

$$(9.1) \quad \Xi := \{w \in W(\mathbb{R}) \mid \dim G_{\dim_{\mathbb{R}} \tilde{Z}}(\mathcal{B} \cap w\Delta \cap w(\mathfrak{F} \times E), \tilde{Z}^{\text{rl}}) = \dim_{\mathbb{R}} \tilde{Z}\}$$

(resp. $\Xi_V := \{v \in V(\mathbb{R}) \mid \dim G_{\dim_{\mathbb{R}} \tilde{Z}_V}(\mathcal{B}_V \cap (v + \Delta_V) \cap (v + (\mathfrak{F}_V \times A)), \tilde{Z}_V^{\text{rl}}) = \dim_{\mathbb{R}} \tilde{Z}_V\}$)

Then Ξ and Ξ_V are definable sets. The request on $\tilde{Z}^{\text{rl}}$ ensures that under certain modifications of $\mathcal{B}$ and $\mathcal{Z}$, we still have that $\tilde{F}$ is the smallest weakly special subset containing $pr_{W(\mathbb{R})U(\mathbb{C})}(\mathcal{Z})$.

Note that we have (since $\mathcal{Z}$ is a complex analytic irreducible component of $\mathcal{B} \cap \Delta$)

$$(9.2) \quad \{\gamma \in \Gamma_W | \gamma \mathfrak{F} \cap \tilde{Z} \neq \emptyset\} = \Xi \cap \Gamma_W \text{ and } \{\gamma_V \in \Gamma_V | (\gamma_V + \mathfrak{F}_V) \cap \tilde{Z}_V \neq \emptyset\} = \Xi_V \cap \Gamma_V.$$

The inclusions $\supset$ are clear. The inclusions $\subset$ are true because $\Gamma_W \Delta = \Delta$ and $\Gamma_V + \Delta_V = \Delta_V$. We do the following reduction:

Lemma 9.4. *To prove Theorem 9.2, we may assume that every positive dimensional maximal connected semi-algebraic subset of Ξ is contained in a right coset of $\text{Stab}_{W(\mathbb{R})}(\mathcal{B})$.*

Proof. We do the lexicographic induction on $(\dim \mathcal{B} - \dim \mathcal{Z}, \dim \mathcal{Z})$ (upwards for the first factor and downwards for the second). Start with the case $\dim \mathcal{B} - \dim \mathcal{Z} = 0$. Then $\mathcal{B} = \mathcal{Z}$ and hence both $\tilde{Z}$ and Z are algebraic. Therefore Z is weakly special by [16, Corollary 8.3], and therefore $Z = F$. So $\mathcal{B} = \tilde{F} \times F$ and $\mathcal{Z} = \Delta_F$, contradicting to $\mathcal{B} = \mathcal{Z}$. So the starting point of the induction on the first factor is proven. Let us prove the starting point of the induction on the second factor. For a given $\dim \mathcal{B} - \dim \mathcal{Z}$, if $\dim \mathcal{Z} = \dim F$, then $\mathcal{Z} = \Delta_F$. But the action of $W(\mathbb{R})U(\mathbb{C})$ on $W(\mathbb{R})U(\mathbb{C}) \times E$ is algebraic, so $\tilde{F} \times F = W_F(\mathbb{R})U_{W_F}(\mathbb{C})\Delta_F \subset \mathcal{Z}^{\text{Zar}} = \mathcal{B}$, and therefore $\mathcal{B} = \tilde{F} \times F$. Hence we are done for this case.

Now we do the induction. Let C be a connected semi-algebraic curve in Ξ . For each c in an open neighborhood of C , let $\mathcal{Z}_c$ be a complex analytic irreducible component of $c\mathcal{B} \cap \Delta$ such that $\dim \mathcal{Z}_c = \dim \mathcal{Z} = \dim \tilde{Z}$ and $(pr_{W(\mathbb{R})U(\mathbb{C})}(\mathcal{Z}_c))^{\text{rl}} = \tilde{Z}^{\text{rl}}$. In particular, the smallest weakly special subset of $W(\mathbb{R})U(\mathbb{C})$ containing $pr_{W(\mathbb{R})U(\mathbb{C})}(\mathcal{Z}_c)$ is $\tilde{F}$. Let $c_0 \in C$, then there are 3 possibilities:

- (1) $\mathcal{B}$ is independent of $c \in C$;
- (2) $\mathcal{B}$ is not independent of $c \in C$ but $\mathcal{Z}_{c_0} \subset c\mathcal{B}$ for all $c \in C$;
- (3) $\mathcal{B}$ is not independent of $c \in C$ and $\mathcal{Z}_{c_0} \not\subset c'\mathcal{B}$ for some $c' \in C$.

If we are in case (2), then let $\mathcal{B}_1 := \mathcal{B} \cap c\mathcal{B}$ for a generic $c \in C$. Then $\dim \mathcal{B}_1 < \dim \mathcal{B}$. But $\dim \mathcal{Z}_{c_0} = \dim \mathcal{Z}$. Apply the inductive hypothesis on $\dim \mathcal{B} - \dim \mathcal{Z}$ to $(\mathcal{B}_1, \mathcal{Z}_{c_0})$, we have $\dim \mathcal{B}_1 - \dim \mathcal{Z}_{c_0} \geq \dim F$. Hence the Theorem 9.2 is true. If we are in case (3), then let $\mathcal{B}_2 := (C\mathcal{B})^{\text{Zar}}$ and let $\mathcal{Z}_2$ be a complex analytic irreducible component of $\mathcal{B}_2 \cap \Delta$. Then $\dim \mathcal{B}_2 = \dim \mathcal{B} + 1$ and $\dim \mathcal{Z}_2 = \dim \mathcal{Z} + 1$ since $\mathcal{Z}_2$ contains $\cup_{c \in C} \mathcal{Z}_c$. Hence $\dim \mathcal{B}_2 - \dim \mathcal{Z}_2 = \dim \mathcal{B} - \dim \mathcal{Z}$, and we can apply the inductive hypothesis on $\dim \mathcal{Z}$ to $(\mathcal{B}_2, \mathcal{Z}_2)$ and get $\dim \mathcal{B}_2 - \dim \mathcal{Z}_2 \geq \dim F$. Hence Theorem 9.2 is true.

So we may assume that we are in Case (1) for any connected semi-algebraic C in Ξ or in Ξ_U . Then every such C is contained in a right coset of $\text{Stab}_{W(\mathbb{R})}(\mathcal{B})$. Hence the same holds for every positive dimensional maximal connected semi-algebraic subset of Ξ . $\square$

This Lemma suggests that it suffices to prove Theorem 9.2 under the following assumption:

Assumption 9.5. *There exists a definable subset Ξ' in $P(\mathbb{R})$ such that $\Xi' \cap \Gamma_W = \Xi \cap \Gamma_W$ and that every positive dimensional maximal connected semi-algebraic subset of Ξ' is contained in a right coset of $\text{Stab}_{P(\mathbb{R})}(\mathcal{B})$.*

The following proposition will be used in the next subsection.

Proposition 9.6. *Under Assumption 9.5. Let $(v, a) \in \mathfrak{Z}_V$. Then $\mathcal{B}_{(v,a)}(:= \mathcal{B} \cap \pi^{-1}((v, a))) = (U'(\mathbb{C}) + \tilde{z}) \times Y_a$ for some $U' < U$ and $\tilde{z} \in \tilde{Z}_v \subset \tilde{X}_v$.*

Proof. Define

$$\Xi_U := \{u \in U(\mathbb{R}) \mid \dim G_{\dim_{\mathbb{R}} \tilde{Z}_v}(\mathcal{B}_{(v,a)} \cap (u + \Delta_{(v,a)}) \cap (u + (\mathfrak{F}_v \times T)), \tilde{Z}_v^{\text{rl}}) = \dim_{\mathbb{R}} \tilde{Z}_v\}.$$

Then Ξ_U is definable and

$$(9.3) \quad \Xi_U \cap \Gamma_U = \Xi' \cap \Gamma_U = \{\gamma_U \in \Gamma_U \mid (\gamma_U + \mathfrak{F}_v) \cap \tilde{Z} \neq \emptyset\}$$

since $\Gamma_U + \Delta_{(v,a)} = \Delta_{(v,a)}$. For any integer $M > 0$, let

$$\Theta_U(\tilde{Z}_v, M) := \{\gamma_U \in \Gamma_U \cap \Xi' \mid H(\gamma_U) \leq M\}.$$

Claim 1. $|\Theta_U(\tilde{Z}_v, M)| \gg M$.

Proof. Define the norm of $x = (x_1, \dots, x_m) \in U(\mathbb{C})$ to be $\|x\| := \max(|x_1|, \dots, |x_m|)$. It is clear that there exists a constant $c_1 > 0$ such that

$$(9.4) \quad H(\gamma_U) \leq c_1 \|x\|$$

for any $x \in U(\mathbb{C})$ and any $\gamma_U \in \Gamma_U$ such that $\gamma_U + x \in \mathfrak{F}_v$. Let $\omega := \bigwedge^{\dim \tilde{Z}_v} (dx_1 \wedge d\bar{x}_1 + \dots + dx_m \wedge d\bar{x}_m)$ and let $\tilde{Z}_v(M) := \{\tilde{u} \in \tilde{Z}_v \mid \|\tilde{u}\| \leq M\}$. Then there exists a constant c'_2 such that

$$\begin{aligned} \int_{\tilde{Z}_v(M) \cap (\gamma_U + \mathfrak{F}_v)} \omega &= \int_{(\tilde{Z}_v(M) - \gamma_U) \cap \mathfrak{F}_v} \omega \leq \sum_{I \subset \{1, \dots, m\}, |I| = \dim \tilde{Z}_v} \deg(p_I|_{(\tilde{Z}_v - \gamma_U) \cap \mathfrak{F}_v}) \int_{p_I(\{x \in \mathfrak{F}_v \mid \text{Im}(x) \leq M\})} \omega \\ &= c'_2 M \sum_{I \subset \{1, \dots, m\}, |I| = \dim \tilde{Z}_v} \deg(p_I|_{(\tilde{Z}_v - \gamma_U) \cap \mathfrak{F}_v}). \end{aligned}$$

The function $U(\mathbb{R}) \rightarrow \mathbb{Z}$, $u \mapsto \deg(p_I|_{pr_{U(\mathbb{C})}((\mathcal{B}_{(v,a)} - u) \cap \Delta_{(v,a)} \cap (\mathfrak{F}_v \times T))})$ is a definable function with value in $\mathbb{Z}$, and hence is uniformly bounded. Therefore there exists a constant c_2 such that

$$(9.5) \quad \int_{\tilde{Z}_v(M) \cap (\gamma_U + \mathfrak{F}_v)} \omega \leq c_2 M.$$

By [20, Theorem 0.1] there exists a constant $c_3 > 0$ such that

$$(9.6) \quad \int_{\tilde{Z}_v(M)} \omega \geq c_3 M^{\dim \tilde{Z}_v} \geq c_3 M^2.$$

Now we have

$$\tilde{Z}_v(M) = \bigcup_{\substack{\gamma_U \in \Gamma_U \\ (\gamma_U + \mathfrak{F}_v) \cap \tilde{Z}_v \neq \emptyset}} (\gamma_U + \mathfrak{F}_v) \cap \tilde{Z}_v(M) \subset \bigcup_{\gamma_U \in \Theta_U(\tilde{Z}_v, 2M)} (\gamma_U + \mathfrak{F}_v) \cap \tilde{Z}_v(M)$$

by (9.4) and $H(\gamma_U) \leq M \Rightarrow \gamma_U + \tilde{Z}_v(M) \subset (\gamma_U + \tilde{Z}_v)(2M)$. Integrating both sides with respect to ω we can conclude by (9.5) and (9.6). $\square$

By Pila-Wilkie [29, Theorem 3.6], there exists a positive dimensional connected semi-algebraic subset $K \subset \Xi'$ containing arbitrarily many points $\gamma_U \in \Gamma_U$. Fix a $\sigma_U \in K \cap \Gamma_U$. We have $K \subset \text{Stab}_{P(\mathbb{R})}(\mathcal{B}_{(v,a)})\sigma_U$ by Assumption 9.5. So $(K \cap \Gamma_U)\sigma_U^{-1} \subset \text{Stab}_{P(\mathbb{R})}(\mathcal{B}_{(v,a)}) \cap \Gamma_U$. Therefore the $\mathbb{Q}$ -group

$$U' := (\text{Stab}_{P(\mathbb{R})}(\mathcal{B}_{(v,a)}) \cap \Gamma_U)^{\text{Zar}},$$

which is the largest subgroup of U such that $\mathcal{B}_{(v,a)}$ is stable under $U'(\mathbb{C})$, is of positive dimension. This group is normal in P by Proposition 3.12(2), so we can consider $\rho: (P, \mathcal{X}^+) \rightarrow (P, \mathcal{X}^+)/U'$.

If $\dim \rho(\tilde{Z}_v) > 0$, then we can repeat the process above and obtain a subgroup U_2 of U which is larger than U' such that $\mathcal{B}_{(v,a)}$ is stable under $U_2(\mathbb{C})$. But this contradicts the maximality of U' . Hence $\rho(\tilde{Z}_v)$ is a point. So we can conclude because $\mathcal{B}_{(v,a)} \subset U'(\mathbb{C}) + \tilde{z}_v \subset \mathcal{B}_{(v,a)}$. $\square$

Corollary 9.7. *Under Assumption 9.5. Let $U_{\mathcal{B}} = (\text{Stab}_{W(\mathbb{R})}(\mathcal{B}) \cap \Gamma_U)^{\text{Zar}, \circ}$. Denote by*

$$\lambda: (P, \mathcal{X}^+) \rightarrow (P, \mathcal{X}^+)/U_{\mathcal{B}} \quad \text{and} \quad (\lambda, 1): W(\mathbb{R})U(\mathbb{C}) \times E \rightarrow (W/U_{\mathcal{B}})(\mathbb{R})(U/U_{\mathcal{B}})(\mathbb{C}) \times E.$$

Then $\mathcal{B} = (\lambda, 1)^{-1}(\lambda, 1)(\mathcal{B})$ and the natural projection $\lambda(\tilde{X}) \rightarrow \tilde{X}_V$ is a bijection.

Proof. For any $(v, a) \in \mathcal{Z}_V$ (then $a = \text{unif}_V(v)$), apply Proposition 9.6 to $\mathcal{B}_{(v,a)}$ we have

$$\mathcal{B}_{(v,a)} = (U'_v(\mathbb{C}) + \tilde{z}_v) \times Y_a$$

for some $U'_v < U$ and $\tilde{z}_v \in \tilde{Z}_v \subset \tilde{X}_v$. Then $U_{\mathcal{B}} < U'_v$.

Now let $\tilde{C}_V$ be any complex analytic irreducible curve in $\tilde{Z}_V$. There are uncountably many points in $\tilde{C}$ and there are only countably many subgroups of U , hence there exists a subgroup U' of U such that $U'_v = U'$ for uncountably many $v \in \tilde{C}_V$. Replacing $(P, \mathcal{X}^+)$ by $(P, \mathcal{X}^+)/U'$ (this can be done since $U' \triangleleft P$ by Proposition 3.12(2)) and every other object accordingly, we see that $U'_v \subset U'$ except for finitely many $v \in \tilde{C}_V$ by generic flatness (in the complex analytic geometry sense) and dimension reasons. But $\dim U'_v$ is constant except for finitely many $v \in \tilde{C}_V$ by generic flatness (in the complex analytic geometry sense) and dimension reasons. Hence $U'_v = U'$ except for finitely many $v \in \tilde{C}_V$. But then we must have $U'_v = U'$ for all $v \in \tilde{C}_V$ (this can be seen by, for example, considering the smallest complex analytic irreducible subvariety containing $\bigcup_{v \in \tilde{C}_V, U'_v = U'} \mathcal{B}_{(v, \text{unif}_V(v))}$). By varying $\tilde{C}_V$ we get $U'_v = U'$ for all $v \in \tilde{Z}_V$, i.e.

$$(9.7) \quad \mathcal{B}_{(v,a)} = (U'(\mathbb{C}) + \tilde{z}_v) \times Y_a, \quad \forall (v, a) \in \mathcal{Z}_V$$

for some $U' < U$ independent of v containing $U_{\mathcal{B}}$. So $\mathcal{B}|_{\mathcal{Z}_V} := \mathcal{B} \cap \pi^{-1}(\mathcal{Z}_V)$ is stable under U' . But $\mathcal{B}|_{\mathcal{Z}_V}$, which contains $\mathcal{Z}$, is Zariski dense in $\mathcal{B}$. So $\mathcal{B}$ is stable under U' , and hence $U' < U_{\mathcal{B}}$. So $U' = U_{\mathcal{B}}$ and therefore $\mathcal{B} = (\lambda, 1)^{-1}(\lambda, 1)(\mathcal{B})$. Now the natural projection $\lambda(\tilde{X}) \rightarrow \tilde{X}_V$ is a bijection by (9.7) and generic flatness and $\tilde{Z}_V^{\text{Zar}} = \tilde{X}_V$. $\square$

9.1.2. Proof of Theorem 9.2. The variety E is a T -torsor over A , where $T = \Gamma_U \backslash U(\mathbb{C})$ is an algebraic torus. Let T' be any subgroup of T , then we denote by $\underline{T}'$ the subvariety of E which is a T' -torsor over A , compatible with the T -torsor structure of E .

Theorem 9.8. *Under Assumption 9.5, we have $\mathcal{B} = \tilde{X} \times Y$ and*

$$(1) \quad \tilde{X} = W_{\mathcal{B}}(\mathbb{R})U_{\mathcal{B}}(\mathbb{C})\tilde{z} \text{ for some } \tilde{z} \in \tilde{Z}, \text{ where } W_{\mathcal{B}} := (\text{Stab}_{W(\mathbb{R})}(\mathcal{B}) \cap \Gamma_W)^{\text{Zar}};$$

(2) there exists a subgroup $U^\dagger$ of U such that $U^\dagger \oplus U_{\mathbf{B}} = U_F$ satisfying the following properties: for the projection $\rho: (P, \mathcal{X}^+) \rightarrow (P, \mathcal{X}^+)/U^\dagger$ (this can be done since $U^\dagger \triangleleft P$ by Proposition 3.12(2)) we have

- $\rho(\tilde{X}) = \rho(\tilde{F})$;
- $Y = [\rho]^{-1}[\rho](Y)$, where $[\rho]$ is the projection on the mixed Shimura varieties associated with ρ .

Remark 9.9. In the case $E = T$ or $E = A$, the conclusion is simpler: under the assumption of Lemma 9.4 we have $\mathbf{B} = \tilde{F} \times Y$. The case $E = T$ is Proposition 9.6, and the case $E = A$ will be proven later.

Before proving this theorem, let us see how it implies Theorem 9.2.

Theorem 9.8 implies Theorem 9.2. By Lemma 9.4, it is enough to prove Theorem 9.2 under Assumption 9.5. So we are in the situation of Theorem 9.8. Then

$$\begin{aligned} \dim \mathbf{B} - \dim \mathbf{Z} - \dim F &= \dim \tilde{X} + \dim Y - \dim \tilde{Z} - \dim \tilde{F} \\ &= \dim \rho(\tilde{X}) + \dim T^\dagger + \dim [\rho](Y) - \dim \tilde{Z} - (\dim \rho(\tilde{F}) + \dim U^\dagger(\mathbb{C})) \\ &= \dim [\rho](Y) - \dim \tilde{Z} \\ &= \dim [\rho](Y) - \dim \rho(\tilde{Z}) \geq 0 \end{aligned}$$

where the last equality is true because $U_{\mathbf{B}} \cap U^\dagger = 0$. □

Proof of Theorem 9.8.

Step I Prove $\dim \pi(W_{\mathbf{B}}) > 0$.

Let us point out that the proof of this step does NOT require any complex structure except for the proof of (9.10): we view $V(\mathbb{R})$, A and $\mathbf{B}$ as real algebraic varieties and $\tilde{Z}$ as real analytic. This fact is important for the next step.

For any integer $M > 0$, let

$$\Theta_V(\tilde{Z}_V, M) = \{\gamma_V \in \Gamma_V \cap \Xi_V \mid H(\gamma_V) \leq M\}.$$

Claim 2. $|\Theta_V(\tilde{Z}_V, M)| \gg M^{\dim_{\mathbb{R}} \tilde{Z}_V}$.

Proof. Define the norm of any $v = (v_1, v_2, \dots, v_{2n}) \in V(\mathbb{R}) \simeq \mathbb{R}^{2n}$ to be $\|v\| := \max(|v_1|, |v_2|, \dots, |v_{2n}|)$. It is clear that there exists a constant $c_1 > 0$ such that

$$(9.8) \quad H(\gamma_V) \leq c_1 \|\tilde{z}_V\|$$

for any $\tilde{z}_V \in V(\mathbb{R})$ and any $\gamma_V \in \Gamma_V$ such that $\gamma_V + \tilde{z}_V \in \mathfrak{F}_V$. Let $\omega_V := \bigwedge^{\dim_{\mathbb{R}} \tilde{Z}} (dv_1 + \dots + dv_{2n})$. Then there exists a constant c'_2 such that

$$\begin{aligned} \int_{\tilde{Z}_V \cap (\gamma_V + \mathfrak{F}_V)} \omega_V &= \int_{(\tilde{Z}_V - \gamma_V) \cap \mathfrak{F}_V} \omega_V \leq \sum_{I \subset \{1, \dots, 2n\}, |I| = \dim_{\mathbb{R}} \tilde{Z}} \deg(p_I|_{(\tilde{Z}_V - \gamma_V) \cap \mathfrak{F}_V}) \int_{p_I(\mathfrak{F}_V)} \omega_V \\ &= c'_2 \sum_{I \subset \{1, \dots, 2n\}, |I| = \dim_{\mathbb{R}} \tilde{Z}} \deg(p_I|_{(\tilde{Z}_V - \gamma_V)}) \end{aligned}$$

The function $V(\mathbb{R}) \rightarrow \mathbb{Z}$, $v \mapsto \deg(p_I|_{pr_{V(\mathbb{R})}}((\mathcal{B}_V - v) \cap \Delta_V \cap (\mathfrak{F}_V \times A)))$ is a definable function with values in $\mathbb{Z}$, and hence is uniformly bounded. Therefore there exists a constant c_2 such that

$$(9.9) \quad \int_{\tilde{Z}_V \cap (\gamma_V + \mathfrak{F}_V)} \omega_V \leq c_2.$$

Let $\tilde{Z}_V(M) := \{\tilde{z}_V \in \tilde{Z}_V \mid \|\tilde{z}_V\| \leq M\}$. Then by [20, Theorem 0.1] (note that $\tilde{Z}_V$ is complex analytic) there exists a constant $c_3 > 0$ such that

$$(9.10) \quad \int_{\tilde{Z}_V(M)} \omega_V \geq c_3 M^{\dim_{\mathbb{R}} \tilde{Z}_V}.$$

Now we have

$$\tilde{Z}_V(M) = \bigcup_{\substack{\gamma_V \in \Gamma_V \\ (\gamma_V + \mathfrak{F}_V) \cap \tilde{Z}_V \neq \emptyset}} (\gamma_V + \mathfrak{F}_V) \cap \tilde{Z}_V(M) \subset \bigcup_{\gamma_V \in \Theta_V(\tilde{Z}_V, M)} (\gamma_V + \mathfrak{F}_V) \cap \tilde{Z}_V \text{ by (9.8).}$$

Integrating both sides with respect to ω_V we can conclude by (9.9) and (9.10). $\square$

Consider the definable set Ξ_V . By the theorem of Pila-Wilkie ([29, Theorem 3.6]), there exist J definable block families

$$K^j \subset \Xi_V \times \mathbb{R}^l, \quad j = 1, \dots, J$$

and $c > 0$ such that for all $M \gg 0$, $\Theta_V(\tilde{Z}_V, M^{1/4})$ is contained in the union of at most $cM^{\delta/8}$ definable blocks of the form K_y^j ($y \in \mathbb{R}^l$). By Claim 2, there exist a $j \in \{1, \dots, J\}$ and a block $K_V := K_{y_0}^j$ of Ξ_V containing at least $M^{\delta/8}$ elements of $\Theta_V(\Xi_V, M^{1/4})$.

Consider $X^j := (\pi \times 1_{\mathbb{R}^l})^{-1}(K^j) \cap (\Xi \times \mathbb{R}^l)$, which is a definable family since π is $\mathbb{R}$ -algebraic. By [44, Ch. 3, 3.6], there exists a number $n_0 > 0$ such that each fibre X_y^j has at most n_0 connected components. So the definable set $\pi^{-1}(K_V) \cap \Xi'$ has at most n_0 connected components. Now

$$\pi(\pi^{-1}(K_V) \cap \Xi' \cap \Gamma_W) = K_V \cap \pi(\Xi' \cap \Gamma_W) = K_V \cap (\Xi_V \cap \Gamma_V)$$

by (9.2) and Assumption 9.5. So there exists a connected component K of $\pi^{-1}(K_V) \cap \Xi'$ such that $\pi(K \cap \Gamma_W)$ contains at least $M^{\delta/8}/n_0$ elements of $\Theta_V(\tilde{Z}_V, M^{1/4})$.

Fix a $\sigma_W \in K \cap \Gamma_W$ such that $\pi(\sigma_W) \in \Theta_V(\tilde{Z}_V, M^{1/4})$. By Assumption 9.5 we have $K \subset \text{Stab}_{W(\mathbb{R})}(\mathcal{B})\sigma_W$. We have shown that $\pi(K \cap \Gamma_W)$ contains at least $M^{\delta/8}/n_0$ elements of $\Theta_V(\tilde{Z}_V, M^{1/4})$. For any $\sigma_V \in \pi(K \cap \Gamma_W) \cap \Theta_V(\tilde{Z}_V, M^{1/4})$, let σ'_W be one of its pre-images in $K \cap \Gamma_W$. Then $\gamma_W := \sigma_W^{-1}\sigma'_W$ is an element of $\Gamma_W \cap \text{Stab}_{W(\mathbb{R})}(\mathcal{B})$ and $H(\pi(\gamma_W)) \ll M^{1/2}$. Therefore for $M \gg 0$, $\pi(\Gamma_W \cap \text{Stab}_{W(\mathbb{R})}(\mathcal{B}))$ contains at least $M^{\delta/8}/n_0$ elements γ_V such that $H(\gamma_V) \leq M$. Therefore $\dim \pi(W_{\mathcal{B}}) > 0$ since it is an infinite set.

Step II Prove $\pi(W_{\mathcal{B}}) = V_F$. Hence $\pi(\mathcal{B}) = \tilde{F}_V \times Y_V$. (Note that this proves Remark 9.9 for the case $E = A$).

Consider the projection $p: W \rightarrow V \rightarrow V/\pi(W_{\mathcal{B}})$ (we use the same notation for the projection on the underlying spaces $W(\mathbb{R})U(\mathbb{C}) \rightarrow (V/\pi(W_{\mathcal{B}}))(\mathbb{R})$). Then the image of $p(\tilde{Z})$ is a point: otherwise $\Gamma_{V/\pi(W_{\mathcal{B}})} \backslash (V/\pi(W_{\mathcal{B}}))(\mathbb{R})$ is a real algebraic variety, and hence we can repeat Step I (recall that the complex structure is not used in Step I except for proving (9.10), which we can get for $p(\tilde{Z})$ using directly (9.10)) and get $\dim p(W_{\mathcal{B}}) > 0$, which is absurd. Therefore

$$\tilde{Z}_V \subset \pi(W_{\mathcal{B}})(\mathbb{R}) + \tilde{z}_V \subset \tilde{X}_V.$$

The first inclusion implies that $\tilde{Z}_V$ is contained in $V_c + \tilde{z}_V$ (which is irreducible algebraic in $\tilde{F}_V$), where V_c is the largest complex analytic subspace of $\pi(W_{\mathcal{B}})(\mathbb{R})$. But then $\tilde{X}_V = \tilde{Z}_V^{\text{Zar}} \subset V_c + \tilde{z}_V$. So

$$\pi(W_{\mathcal{B}})(\mathbb{R}) + \tilde{z}_V \subset \tilde{X}_V \subset V_c + \tilde{z}_V \subset \pi(W_{\mathcal{B}})(\mathbb{R}) + \tilde{z}_V,$$

and therefore $\tilde{X}_V = \pi(W_{\mathcal{B}})(\mathbb{R}) + \tilde{z}_V$. But $\tilde{X}_V$ is complex analytic, so $\pi(W_{\mathcal{B}})$ is a sub-Hodge structure of V , and hence $\pi(W_{\mathcal{B}})(\mathbb{R}) + \tilde{z}_V$ is a weakly special subset of $V(\mathbb{R})$. But the smallest weakly special subset of $V(\mathbb{R})$ containing $\tilde{X}_V$ is $\tilde{F}_V$, so $\pi(W_{\mathcal{B}})(\mathbb{R}) + \tilde{z}_V = \tilde{F}_V$. Therefore $\pi(W_{\mathcal{B}}) = V_F$.

Step III Prove $\mathcal{B} = (W_{\mathcal{B}}(\mathbb{R})U_{\mathcal{B}}(\mathbb{C})\tilde{z}) \times Y$. In particular $\tilde{X} = W_{\mathcal{B}}(\mathbb{R})U_{\mathcal{B}}(\mathbb{C})\tilde{z}$.

This follows directly from Step II and Corollary 9.7.

For the rest we fix some extra notation. Let $\tilde{z} = (\tilde{z}_U, \tilde{z}_V)$ under the identification $W(\mathbb{R})U(\mathbb{C}) \simeq U(\mathbb{C}) \times V(\mathbb{R})$ in [16, §4]. We also use the notation of §3.3.

Step IV Let $U^\dagger$ be the smallest subgroup of U such that $\Psi(\Gamma_{V_F}, \tilde{z}_V) \subset U(\mathbb{Q}) + U_{\mathcal{B}}(\mathbb{R}) + U^\dagger(\mathbb{R})$. Consider $(P, \mathcal{X}^+) \xrightarrow{\rho} (P, \mathcal{X}^+)/U^\dagger$ (this can be done since $U^\dagger \triangleleft P$ by Proposition 3.12(2)). Then $Y = [\rho]^{-1}[\rho](Y)$, where $[\rho]$ is the projection on the mixed Shimura varieties associated with ρ .

First of all, $U_{\mathcal{B}} \triangleleft P$ by Proposition 3.12(2). Consider $(P, \mathcal{X}^+) \xrightarrow{\lambda} (P, \mathcal{X}^+)/U_{\mathcal{B}}$. Replace $(P, \mathcal{X}^+)$ (resp. $W, \tilde{X}, W_{\mathcal{B}}, \tilde{z}, Y$) by $(P, \mathcal{X}^+)/U_{\mathcal{B}}$ (resp. $W/U_{\mathcal{B}}, \lambda(\tilde{X}), V_{\mathcal{B}}, \lambda(\tilde{z})$ and $[\lambda](Y)$), then the assumption and the conclusion of this step do not change. Now $V_{\mathcal{B}}$ is a subgroup of W and $\tilde{X} = V_{\mathcal{B}}(\mathbb{R})\tilde{z}$.

Every fiber of $\tilde{X} \rightarrow \tilde{X}_V$ is a point, so

$$(9.11) \quad \tilde{Z} = \tilde{X}|_{\tilde{Y}_V} = \tilde{X} \cap \pi^{-1}(\tilde{Y}_V).$$

The monodromy group Γ_{Y_V} of Y_V , which stabilizes $\tilde{Y}_V$, is Zariski dense in V_F , so Γ_{Y_V} is of finite index in Γ_{V_F} because V_F is a vector group. Up to replacing Γ_V by a subgroup of finite index (which does not change the assumption or the conclusion of this step), we may assume $\Gamma_{Y_V} = \Gamma_{V_F}$. Let $\Gamma'_W := \pi^{-1}(\Gamma_{V_F}) \cap \Gamma_W$. Then (9.11) implies

$$(9.12) \quad Z_{z_V} = \left(\text{unif}(\Gamma'_W \tilde{Z}) \right)_{z_V} = \tilde{z}_U + \frac{1}{2} \Psi(\Gamma_{V_F}, \tilde{z}_V) + \Gamma_U \pmod{\Gamma_U}.$$

We have $\Psi(V(\mathbb{R}), V(\mathbb{R})) \subset U(\mathbb{R})$ since Ψ is defined over $\mathbb{Q}$. Write $\Gamma_U = \Gamma_{U^\dagger} \oplus \Gamma_{U_2}$, which induces a decomposition $U = U^\dagger \oplus U_2$ for a $\mathbb{Q}$ -subgroup U_2 of U . The assumption on $U^\dagger$ implies that for any coordinate (i.e. any 1-dimensional quotient U_0 of $U^\dagger(\mathbb{R})$), there exists a $u \in \Psi(\Gamma_{V_F}, \tilde{z}_V)$ such that the chosen coordinate of u , which we call u_0 , is irrational. But then $\text{unif}(\tilde{z}_U + U_0(\mathbb{R})) \subset Z_{z_V}^{\text{Zar}}$ since $\{lu_0 \bmod \mathbb{Z} | l \in \mathbb{Z}\}$ is dense in $[0, 1)$. So $Z_{z_V}^{\text{Zar}}$, which is complex analytic, contains $\text{unif}(\tilde{z}_U + U_0(\mathbb{C}))$, and so does Y_{z_V} since $Y = Z^{\text{Zar}}$. Hence the conclusion of this step follows by varying U_0 (which is any coordinate of $U^\dagger$).

Step V $\rho(\tilde{X})$ is weakly special. In particular $\rho(\tilde{X}) = \rho(\tilde{F})$ since $\tilde{F}$ is the smallest weakly special subset containing $\tilde{X}$.

Let $U_1 := U_{\mathcal{B}} + U^\dagger$ and let $(P, \mathcal{X}^+) \xrightarrow{\rho'} (P, \mathcal{X}^+)/U_1$. Then it suffices to prove that $\rho'(\tilde{X})$ is weakly special. For simplicity denote by $V_{\mathcal{B}} := \pi(W_{\mathcal{B}})$. By [16, Proposition 2.7], $\rho'(\tilde{X})$ is weakly special iff $\tilde{z}_V \in (N_{\rho'(W)}(\rho'(V_{\mathcal{B}}))/\rho'(U))(\mathbb{R})$ iff $\rho'(\Psi(V_{\mathcal{B}}(\mathbb{R}), \tilde{z}_V)) = 0$. We prove the last claim.

We have $\tilde{X}_V = \tilde{F}_V = V_{\mathcal{B}}(\mathbb{R}) + \tilde{z}_V$ and $X_V = A_0 + z_V$ where $A_0 = \Gamma_{V_{\mathcal{B}}} \setminus V_{\mathcal{B}}(\mathbb{R})$ is an abelian subvariety of A . By Step IV, we have $\rho'(\Psi(\Gamma_{V_F}, \tilde{z}_V)) \subset \rho'(U)(\mathbb{Q})$, and therefore $(1/2)\rho'(\Psi(N\Gamma_{V_F}, \tilde{z}_V)) \subset \Gamma_{\rho'(U)}$ for some $N \gg 0$ (since $\text{rank} \Gamma_{V_F} < \infty$). Now we can construct a new lattice Γ' with $N\Gamma_V$ and $\Gamma_{\rho'(U)}$, which is of finite index in $\Gamma_{\rho'(W)}$. Replacing $\Gamma_{\rho'(W)}$ by Γ' does not change the assumption or the conclusion of this step, so we may assume $(1/2)\rho'(\Psi(N\Gamma_{V_F}, \tilde{z}_V)) \subset \Gamma_{\rho'(U)}$. Now we can define C^∞ -morphisms

$$f: A_0 + z_V \rightarrow [\rho'](T), \quad a_0 + z_V \mapsto \rho'(\tilde{z}_U + (1/2)\Psi(\Gamma_{V_F}, \tilde{z}_V)) \bmod \Gamma_{\rho'(U)}$$

and

$$s: A_0 + z_V \rightarrow [\rho'](E)|_{A_0 + z_V}, \quad a_0 + z_V \mapsto (\rho'(\tilde{z}_U + (1/2)\Psi(\Gamma_{V_F}, \tilde{z}_V)), a_0 + z_V) \bmod \Gamma_{\rho'(W)}$$

where v_0 is any point of $V_{\mathcal{B}}(\mathbb{R})$ such that $\text{unif}_V(v_0) = a_0$. But Z_a is a single point for all $a \in A_0 + z_V$ by (9.21), so s is the inverse of $[\rho'\pi]|_Z$, and therefore s is a holomorphic section of $[\rho'](E)|_{A_0 + z_V} \rightarrow A_0 + z_V$. Locally on $U_i \subset A_0 + z_V$, s is represented by a holomorphic morphism $U_i \rightarrow [\rho'](T)$, which must equal to $f|_{U_i}$ by definition. Hence f is holomorphic since being holomorphic is a local condition. So f is constant.

But $\Psi(0, \tilde{z}_V) = 0$, and therefore $(1/2)\rho'(\Psi(V_{\mathcal{B}}(\mathbb{R}), \tilde{z}_V)) \subset \Gamma_{\rho'(U)}$. But $\Psi(V_{\mathcal{B}}(\mathbb{R}), \tilde{z}_V)$ is continuous and $\Psi(0, \tilde{z}_V) = 0$, so $\rho'(\Psi(V_{\mathcal{B}}(\mathbb{R}), \tilde{z}_V)) = 0$. Hence we are done.

Conclusion Now the conclusion follows from Step IV, V and VI. $\square$

9.2. Ax-Schanuel for $E^\natural$. In this subsection we prove the Ax-Schanuel conjecture for $E^\natural$, i.e. Theorem 9.1. Recall the statement of the theorem: $\Delta^\natural = \text{graph}(W(\mathbb{C}) \xrightarrow{\text{unif}^\natural} E^\natural)$, $\mathcal{Z}^\natural = \text{graph}(\tilde{Z}^\natural \xrightarrow{\text{unif}^\natural} Z^\natural)$ and $\mathcal{B}^\natural := (\mathcal{Z}^\natural)^{\text{Zar}} \subset W(\mathbb{C}) \times E^\natural$. Let $F^\natural$ (resp. $\tilde{F}^\natural$) be the smallest bi-algebraic, i.e. quasi-linear, subvariety of $E^\natural$ (resp. of $W(\mathbb{C})$) containing $Z^\natural$ (resp. $\tilde{Z}^\natural$). In particular $\text{unif}^\natural(\tilde{F}^\natural) = F^\natural$. Let $[pr]_{F^\natural}^{\text{ws}}$ and $pr_{\tilde{F}^\natural}^{\text{ws}}$ be as in (5.9). Denote for simplicity by $\mathbf{pr}_{F^\natural}^{\text{ws}} := (pr_{\tilde{F}^\natural}^{\text{ws}}, [pr]_{F^\natural}^{\text{ws}}): \tilde{F}^\natural \times F^\natural \rightarrow (\tilde{F}^\natural)^{\text{ws}} \times (F^\natural)^{\text{ws}}$. Then we want to prove

$$(9.13) \quad \dim \mathbf{pr}_{F^\natural}^{\text{ws}}(\mathcal{B}^\natural) - \dim \mathbf{pr}_{F^\natural}^{\text{ws}}(\mathcal{Z}^\natural) \geq \dim(F^\natural)^{\text{ws}}$$

and (for $\tilde{X}^\natural = pr_{W(\mathbb{C})}(\mathcal{B}^\natural) = \tilde{Z}^{\natural, \text{Zar}}$ and $Y^\natural = pr_{E^\natural}(\mathcal{B}^\natural) = Y^{\natural, \text{Zar}}$)

$$(9.14) \quad \dim \tilde{X}^\natural + \dim Y^\natural - \dim \tilde{Z}^\natural \geq \dim F^\natural.$$

Let $\mathfrak{F}^\natural$ be a fundamental set for $\text{unif}^\natural$ such that $\text{unif}^\natural|_{\mathfrak{F}^\natural}$ is definable.

Definition 9.10. • For any definable subset $\tilde{D}^\natural$ of $W(\mathbb{C})$, define $\tilde{D}^{\natural, \text{rl}}$ to be the smallest subset of $W(\mathbb{C})$ of the form $W'(\mathbb{C})w$, where W' is a subgroup of $W_{\mathbb{C}}$ and $w \in W(\mathbb{C})$.

- For a definable subset $\mathcal{D}^\natural$ of $W(\mathbb{C}) \times E^\natural$, a positive integer r and $M = W'(\mathbb{C})w$ ($W' < W_{\mathbb{C}}$, $w \in W(\mathbb{C})$), define $G_r(\mathcal{D}^\natural, M)$ to be the set of points $\mathbf{d}^\natural \in \mathcal{D}^\natural$ around which $\mathcal{D}^\natural$ is regular of dimension r and such that $\tilde{D}^{\natural, \text{rl}} = M$, where $\tilde{D}^\natural$ is the projection of the irreducible component of $\mathcal{D}^\natural$ containing $\mathbf{d}^\natural$ under $W(\mathbb{C}) \times E^\natural \rightarrow W(\mathbb{C})$.

Define

$$\begin{aligned} \Xi^\natural &:= \{w \in W(\mathbb{R}) \mid \dim G_{\dim_{\mathbb{R}} \tilde{Z}^\natural}(\mathcal{B}^\natural \cap w\Delta^\natural \cap w(\mathfrak{F}^\natural \times E^\natural), (\tilde{Z}^\natural)^{\text{rl}}) = \dim_{\mathbb{R}} \tilde{Z}^\natural\} \\ &= \{w \in W(\mathbb{R}) \mid \dim G_{\dim_{\mathbb{R}} (pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{Z}^\natural))}(\mathbf{pr}_{F^\natural}^{\text{ws}}(\mathcal{B}^\natural) \cap w\Delta^\natural \cap w(\mathfrak{F}^\natural \times E^\natural), pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{Z}^\natural)^{\text{rl}}) = \dim_{\mathbb{R}} pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{Z}^\natural)\} \end{aligned}$$

Then

$$(9.15) \quad \{\gamma \in \Gamma_W | \gamma \mathfrak{F}^\natural \cap \tilde{Z}^\natural \neq \emptyset\} = \{\gamma \in \Gamma_W | \gamma \mathfrak{F}^\natural \cap \text{pr}_{F^\natural}^{\text{ws}}(\tilde{Z}^\natural) \neq \emptyset\} = \Xi^\natural \cap \Gamma_W$$

since $\Gamma_W \Delta^\natural = \Delta^\natural$. We have

Lemma 9.11. *To prove Theorem 9.1, we may assume that every positive dimensional maximal connected semi-algebraic subset of $\Xi^\natural$ is contained in a right coset of $\text{Stab}_{W(\mathbb{R})}(\mathcal{B}^\natural) = \text{Stab}_{W(\mathbb{R})}(\text{pr}_{F^\natural}^{\text{ws}}(\mathcal{B}^\natural))$.*

The proof is a copy of the proof of Lemma 9.4, replacing $\mathcal{B}$ by either $\mathcal{B}^\natural$ or $\text{pr}_{F^\natural}^{\text{ws}}(\mathcal{B}^\natural)$ etc. Hence we work under the following assumption:

Assumption 9.12. *There exists a definable subset $\Xi^{\natural, \prime}$ in $P(\mathbb{R})$ such that $\Xi^{\natural, \prime} \cap \Gamma_W = \Xi^\natural \cap \Gamma_W$ and that every positive dimensional maximal connected semi-algebraic subset of $\Xi^{\natural, \prime}$ is contained in a right coset of $\text{Stab}_{P(\mathbb{R})}(\mathcal{B}^\natural) = \text{Stab}_{P(\mathbb{R})}(\text{pr}_{F^\natural}^{\text{ws}}(\mathcal{B}^\natural))$.*

Copying the proof of Proposition 9.6 and Corollary 9.7 we have

Proposition 9.13. *Under Assumption 9.12. Let $(v^\natural, a^\natural) \in \mathcal{Z}_V^\natural$. Then $\mathcal{B}_{(v^\natural, a^\natural)}^\natural (:= \mathcal{B}^\natural \cap \pi^{\natural-1}((v^\natural, a^\natural))) = (U'(\mathbb{C}) + \tilde{z}^\natural) \times (Y^\natural)_{a^\natural}$ for some $U' < U$ and $\tilde{z}^\natural \in (\tilde{Z}^\natural)_{v^\natural} \subset (\tilde{X}^\natural)_{v^\natural}$.*

Corollary 9.14. *Under Assumption 9.12. Let $U_{\mathcal{B}^\natural} = (\text{Stab}_{W(\mathbb{R})}(\mathcal{B}^\natural) \cap \Gamma_U)^{\text{Zar}, \circ}$. Then $U_{\mathcal{B}^\natural} \triangleleft P$ by Proposition 3.12(2). Denote by*

$$\lambda^\natural: (P, \mathcal{X}^{\natural+}) \rightarrow (P, \mathcal{X}^{\natural+})/U_{\mathcal{B}^\natural} \quad \text{and} \quad (\lambda^\natural, 1): W(\mathbb{C}) \times E^\natural \rightarrow (W/U_{\mathcal{B}^\natural})(\mathbb{C}) \times E^\natural.$$

Then $\mathcal{B}^\natural = (\lambda^\natural, 1)^{-1}(\lambda^\natural, 1)(\mathcal{B}^\natural)$ and the natural projection $\lambda^\natural(\tilde{X}^\natural) \rightarrow \tilde{X}_V^\natural$ is a bijection.

Define $W_{\mathcal{B}^\natural} := (\text{Stab}_{W(\mathbb{R})}(\mathcal{B}^\natural) \cap \Gamma_W)^{\text{Zar}, \circ}$. We have a Cartesian diagram

$$\begin{array}{ccc} W(\mathbb{C}) \times E^\natural \xrightarrow{(\pi^\natural, [\pi^\natural])} W(\mathbb{C})/F^0 W(\mathbb{C}) \times E & & \text{pr}_{F^\natural}^{\text{ws}}(\mathcal{B}^\natural) \hookrightarrow \mathcal{B} \\ (\pi^\natural, [\pi^\natural]) \downarrow \lrcorner & \downarrow (\pi, [\pi]) & \downarrow \\ V(\mathbb{C}) \times A^\natural \xrightarrow{(\pi_V^\natural, [\pi_V^\natural])} V(\mathbb{C})/F^0 V(\mathbb{C}) \times A & & \text{pr}_{F^\natural}^{\text{ws}}(\mathcal{B}^\natural)_V \hookrightarrow \mathcal{B}_V \end{array}$$

Proposition 9.15. *Under Assumption 9.12. Then $\text{pr}_{F^\natural}^{\text{ws}}(\mathcal{B}^\natural) = \text{pr}_{F^\natural}^{\text{ws}}(\tilde{X}^\natural) \times [\text{pr}]_{F^\natural}^{\text{ws}}(Y^\natural)$ and $\text{pr}_{F^\natural}^{\text{ws}}(\tilde{X}^\natural) = W_{\mathcal{B}^\natural}(\mathbb{C})\tilde{x}^\natural$ for any $\tilde{x}^\natural \in \text{pr}_{F^\natural}^{\text{ws}}(\tilde{X}^\natural)$. In particular $U_{\mathcal{B}^\natural} = U_{\mathcal{B}}$ by Corollary 9.14.*

Proof. Let $\Delta_V^\natural$ be the graph of $\text{unif}_V^\natural$ and define

$$\Xi_V^\natural := \{v \in V(\mathbb{R}) | \dim G_{\dim_{\mathbb{R}} \pi(\tilde{Z}^\natural)}((\pi(\tilde{X}^\natural) - v) \cap \Delta_V^\natural \cap (\mathfrak{F}_V^\natural \times A^\natural), (\pi(\tilde{Z}^\natural))^{\text{rl}}) = \dim_{\mathbb{R}} \pi(\tilde{Z}^\natural)\}.$$

Then since $\Gamma_V + \Delta_V^\natural = \Delta_V^\natural$, we have $\{\gamma_V \in \Gamma_V | (\gamma_V + \pi^\natural(\mathfrak{F}^\natural)) \cap \pi^\natural(\tilde{Z}^\natural) \neq \emptyset\} = \Xi_V^\natural \cap \Gamma_V$. So by (9.15) and Assumption 9.12, we have

$$(9.16) \quad \Xi_V^{\natural, \prime} \cap \Gamma_V = \pi(\Xi_V^{\natural, \prime} \cap \Gamma_W) = \pi(\Xi_V^{\natural, \prime} \cap \Gamma_W).$$

For any integer $M > 0$, let

$$\Theta_V^{\natural, \prime}(\pi(\tilde{Z}^\natural), M) = \{\gamma_V \in \Gamma_V \cap \Xi_V^{\natural, \prime} | H(\gamma_V) \leq M\}.$$

Lemma 9.16. $|\Theta_V^{\natural, \prime}(\pi(\tilde{Z}^\natural), M)| \gg M^{\dim(\pi_V^\sharp \circ \pi^\natural)(\tilde{Z}^\natural)}.$

Proof. Define the norm of $v = (v_1, v_2, \dots, v_{2n}) \in V(\mathbb{C}) \simeq \mathbb{C}^{2n}$ to be

$$\|v\| := \max(\|v_1\|, \|v_2\|, \dots, \|v_{2n}\|).$$

It is clear that there exists a constant $c_1 > 0$ such that

$$(9.17) \quad H(\gamma_V) \leq c_1 \|\tilde{z}_V\|$$

for any $\tilde{z}_V \in V(\mathbb{C})$ and any $\gamma_V \in \Gamma_V$ such that $\gamma_V + \tilde{z}_V \in \pi'(\mathfrak{F})$. For any $\tilde{K}^\natural \subset W(\mathbb{C})$ define

$$\pi^\natural(\tilde{K}^\natural)(M) := \{\tilde{z}_V^\natural \in \pi^\natural(\tilde{K}^\natural) \mid \|\tilde{z}_V^\natural\| \leq M\}.$$

Let $\omega_V := \bigwedge^{\dim \pi^\natural(\tilde{Z}^\natural)}(dv_1 \wedge d\bar{v}_1 + \dots + dv_{2n} \wedge d\bar{v}_{2n})$. Then there exists a constant c'_2 such that

$$\begin{aligned} \int_{\pi^\natural(\tilde{Z}^\natural)(M) \cap (\gamma_V + \pi^\natural(\mathfrak{F}))} \omega_V &= \int_{(\pi^\natural(\tilde{Z}^\natural)(M) - \gamma_V) \cap \pi(\mathfrak{F})} \omega_V \\ &\leq \sum_{I \subset \{1, \dots, 2n\}, |I| = \dim \pi^\natural(\tilde{Z}^\natural)} \deg(p_I|_{\pi^\natural(\tilde{Z}^\natural) - \gamma_V}) \int_{p_I(\pi^\natural(\mathfrak{F}^\natural)(M) - \gamma_V)} \omega_V \\ &= c'_2 \sum_{I \subset \{1, \dots, 2n\}, |I| = \dim \pi^\natural(\tilde{Z}^\natural)} \deg(p_I|_{\pi^\natural(\tilde{Z}^\natural) - \gamma_V}) M^{\dim \pi^\natural(\tilde{Z}^\natural) - \dim(\pi_V^\sharp \circ \pi^\natural)(\tilde{Z}^\natural)} \end{aligned}$$

The function $\mathbb{R}^{4n} \simeq V(\mathbb{C}) \rightarrow \mathbb{Z}$, $v \mapsto \deg(p_I|_{(\pi^\natural(\tilde{X}^\natural) - v) \cap \Delta_V^\natural \cap (\pi^\natural(\mathfrak{F}^\natural) \times A^\natural)})$ is a definable function with values in $\mathbb{Z}$, and hence is uniformly bounded. Therefore there exists a constant c_2 such that

$$(9.18) \quad \int_{\pi^\natural(\tilde{Z}^\natural)(M) \cap (\gamma_V + \pi^\natural(\mathfrak{F}))} \omega_V \leq c_2 M^{\dim \pi^\natural(\tilde{Z}^\natural) - \dim(\pi_V^\sharp \circ \pi^\natural)(\tilde{Z}^\natural)}.$$

By [20, Theorem 0.1] there exists a constant $c_3 > 0$ such that

$$(9.19) \quad \int_{\pi^\natural(\tilde{Z}^\natural)(M)} \omega_V \geq c_3 M^{\dim \pi^\natural(\tilde{Z}^\natural)}.$$

Now we have

$$\begin{aligned} \pi^\natural(\tilde{Z}^\natural)(M) &= \bigcup_{\substack{\gamma_V \in \Gamma_V \\ (\gamma_V + \pi^\natural(\mathfrak{F}^\natural)) \cap \pi^\natural(\tilde{Z}^\natural) \neq \emptyset}} (\gamma_V + \pi^\natural(\mathfrak{F}^\natural)) \cap \pi^\natural(\tilde{Z}^\natural)(M) \\ &\subset \bigcup_{\gamma_V \in \Theta_V^{\natural, \prime}(\pi^\natural(\tilde{Z}^\natural), M)} (\gamma_V + \pi^\natural(\mathfrak{F}^\natural)) \cap \pi^\natural(\tilde{Z}^\natural) \quad \text{by (9.17)}. \end{aligned}$$

Integrating both sides with respect to ω_V we can conclude by (9.18) and (9.19). $\square$

Lemma 9.17. *If $\dim(\pi_V^\sharp \circ \pi^\natural)(\tilde{Z}^\natural) \geq 1$, then $\dim \pi^\natural(W_{\mathfrak{B}^\natural}) > 0$.*

Proof. Consider the definable set $\Xi_V^{\natural, \prime}$. By the theorem of Pila-Wilkie ([29, Theorem 3.6]), there exist J definable block families

$$K^j \subset \Xi_V^{\natural, \prime} \times \mathbb{R}^l, \quad j = 1, \dots, J$$

and $c > 0$ such that for all $M \gg 0$, $\Theta_V^{\natural, \prime}(\pi^{\natural}(\tilde{Z}^{\natural}), M^{1/4})$ is contained in the union of at most $cM^{\delta/8}$ definable blocks of the form K_y^j ($y \in \mathbb{R}^l$). By Lemma 9.16, there exist a $j \in \{1, \dots, J\}$ and a block $K_V := K_{y_0}^j$ of $\Xi_V^{\natural, \prime}$ containing at least $M^{\delta/8}$ elements of $\Theta_V^{\natural, \prime}(\pi^{\natural}(\tilde{Z}^{\natural}), M^{1/4})$.

Consider $X^j := (\pi^{\natural} \times 1_{\mathbb{R}^l})^{-1}(K^j) \cap (\Xi^{\natural, \prime} \times \mathbb{R}^l)$, which is a definable family since $\pi^{\natural}$ is $\mathbb{R}$ -algebraic. By [44, Ch. 3, 3.6], there exists a number $n_0 > 0$ such that each fibre X_y^j has at most n_0 connected components. So the definable set $\pi^{\natural, -1}(K_V) \cap \Xi^{\natural, \prime}$ has at most n_0 connected components. Now

$$\pi^{\natural}(\pi^{\natural, -1}(K_V) \cap \Xi^{\natural, \prime} \cap \Gamma_W) = K_V \cap \pi^{\natural}(\Xi^{\natural, \prime} \cap \Gamma_W) = K_V \cap (\Xi_V^{\natural, \prime} \cap \Gamma_V)$$

by (9.16) and Assumption 9.12. So there exists a connected component K of $\pi^{\natural, -1}(K_V) \cap \Xi^{\natural, \prime}$ such that $\pi^{\natural}(K \cap \Gamma_W)$ contains at least $M^{\delta/8}/n_0$ elements of $\Theta_V^{\natural, \prime}(\pi^{\natural}(\tilde{Z}^{\natural}), M^{1/4})$.

Fix a $\sigma_W \in K \cap \Gamma_W$ such that $\pi^{\natural}(\sigma_W) \in \Theta_V^{\natural, \prime}(\pi^{\natural}(\tilde{Z}^{\natural}), M^{1/4})$. By Assumption 9.12 we have $K \subset \text{Stab}_{W(\mathbb{R})}(\mathcal{B}^{\natural})\sigma_W$. We have shown that $\pi^{\natural}(K \cap \Gamma_W)$ contains at least $M^{\delta/8}/n_0$ elements of $\Theta_V^{\natural, \prime}(\pi^{\natural}(\tilde{Z}^{\natural}), M^{1/4})$. For any $\sigma_V \in \pi^{\natural}(K \cap \Gamma_W) \cap \Theta_V^{\natural, \prime}(\pi^{\natural}(\tilde{Z}^{\natural}), M^{1/4})$, let σ_W be one of its preimages in $K \cap \Gamma_W$. Then $\gamma_W := \sigma_W^{-1}\sigma'_W$ is an element of $\Gamma_W \cap \text{Stab}_{W(\mathbb{R})}(\mathcal{B}^{\natural})$ and $H(\pi(\gamma_W)) \ll M^{1/2}$. Therefore for $M \gg 0$, $\pi^{\natural}(\Gamma_W \cap \text{Stab}_{W(\mathbb{R})}(\mathcal{B}^{\natural}))$ contains at least $M^{\delta/8}/n_0$ elements γ_V such that $H(\gamma_V) \leq M$. Therefore $\dim \pi^{\natural}(W_{\mathcal{B}^{\natural}}) > 0$ since it is an infinite set. $\square$

Consider the diagram

$$\begin{array}{ccccc} W(\mathbb{C}) \times E^{\natural} & \xrightarrow{(\pi^{\natural}, 1)} & V(\mathbb{C}) \times E^{\natural} & \xrightarrow{(p^{\natural}, 1)} & V(\mathbb{C})/\pi^{\natural}(W_{\mathcal{B}^{\natural}})(\mathbb{C}) \times E^{\natural} \\ (\pi^{\natural}, 1) \downarrow & & (\pi_V^{\natural}, 1) \downarrow & & \downarrow \\ W(\mathbb{C})/F^0W(\mathbb{C}) \times E^{\natural} & \longrightarrow & V(\mathbb{C})/F^0V(\mathbb{C}) \times E^{\natural} & \xrightarrow{(p, 1)} & V(\mathbb{C})/(\pi^{\natural}(W_{\mathcal{B}^{\natural}})(\mathbb{C}) + F^0V(\mathbb{C})) \times E^{\natural} \end{array}$$

Then we can get $(p \circ \pi_V^{\natural} \circ \pi^{\natural}, 1)(\mathbf{pr}_{F^{\natural}}^{\text{ws}}(\mathcal{B}^{\natural})) \subset \text{pt} \times E^{\natural}$: otherwise we can repeat the proof of Lemma 9.16 and Lemma 9.17 and get that $\dim(p^{\natural} \circ \pi^{\natural})(W_{\mathcal{B}^{\natural}}) > 0$, which is absurd. Thus

$$\pi^{\natural}(pr_{\tilde{F}^{\natural}}^{\text{ws}}(\tilde{X}^{\natural})) = \pi^{\natural}(W_{\mathcal{B}^{\natural}})(\mathbb{C}) + \tilde{L}^{\natural}$$

with $\tilde{L}^{\natural} \subset F^0V(\mathbb{C})$. Recall that (9.13) for $A^{\natural}$ follows from a theorem of Ax [3, Corollary 1], i.e.

$$\dim \mathbf{pr}_{F^{\natural}}^{\text{ws}}(\mathcal{B}^{\natural})_V - \dim \mathbf{pr}_{F^{\natural}}^{\text{ws}}(\mathcal{Z}^{\natural})_V \geq \dim(F^{\natural})_V^{\text{ws}}.$$

So $\mathbf{pr}_{F^{\natural}}^{\text{ws}}(\mathcal{B}^{\natural})_V = pr_{\tilde{F}^{\natural}}^{\text{ws}}(\tilde{X}^{\natural})_V \times [pr]_{F^{\natural}}^{\text{ws}}(Y^{\natural})_V$ by the discussion below Conjecture 8.1. But the restriction of the uniformizing map $V(\mathbb{C}) \rightarrow A^{\natural}$ to $F^0V(\mathbb{C})$ is the identity map. So $\tilde{L}^{\natural}$ is a point. So $(\pi^{\natural}, 1)(\mathbf{pr}_{F^{\natural}}^{\text{ws}}(\mathcal{B}^{\natural}))$ is stable under $\pi^{\natural}(W_{\mathcal{B}^{\natural}})(\mathbb{C})$. Thus $(\pi^{\natural}, 1)(\mathbf{pr}_{F^{\natural}}^{\text{ws}}(\mathcal{B}^{\natural})) = pr_{\tilde{F}^{\natural}}^{\text{ws}}(\tilde{X}^{\natural})_V \times [pr]_{F^{\natural}}^{\text{ws}}(Y^{\natural})$, and $pr_{\tilde{F}^{\natural}}^{\text{ws}}(\tilde{X}^{\natural})_V = \pi^{\natural}(W_{\mathcal{B}^{\natural}})(\mathbb{C}) + \tilde{x}_V^{\natural}$ for some point $\tilde{x}_V^{\natural}$. Now the conclusion follows from Corollary 9.14 $\square$

By Theorem 9.8 (note that enlarging $\mathcal{B}$ as we did in Lemma 9.4 does not change Y), there exists a subgroup $U^{\dagger}$ satisfying $U^{\dagger} \oplus U_{\mathcal{B}} = U_F$ such that $Y = [\rho]^{-1}[\rho](Y)$, where $\rho: (P, \varphi(\mathcal{X}^+)) \rightarrow (P, \varphi(\mathcal{X}^+))/U^{\dagger}$ and $[\rho]$ is the projection on the mixed Shimura varieties associated with ρ . Let $\rho^{\natural}: (P, \mathcal{X}^{\natural+}) \rightarrow (P, \mathcal{X}^{\natural+})/U^{\dagger}$ and $[\rho^{\natural}]$ be the projection on the enlarged mixed Shimura varieties associated with $\rho^{\natural}$. Recall $U_{\mathcal{B}^{\natural}} = U_{\mathcal{B}}$ by Proposition 9.15. We prove:

Proposition 9.18. $[pr]_{F^\natural}^{\text{ws}}(Y^\natural) = [\rho^\natural]^{-1}[\rho^\natural]([pr]_{F^\natural}^{\text{ws}}(Y^\natural)).$

Proof. Recall $U_{\mathfrak{B}^\natural} \triangleleft P$ by Proposition 3.12(2). Consider $(P, \mathcal{X}^{\natural+}) \xrightarrow{\lambda^\natural} (P, \mathcal{X}^{\natural+})/U_{\mathfrak{B}^\natural}$. Since $U_{\mathfrak{B}^\natural} = U_{\mathfrak{B}}$, we can replace $(P, \mathcal{X}^{\natural+})$ (resp. $W, pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{X}^\natural), W_{\mathfrak{B}^\natural}$ and $Y^\natural$) by $(P, \mathcal{X}^{\natural+})/U_{\mathfrak{B}^\natural}$ (resp. $W/U_{\mathfrak{B}^\natural}, \lambda^\natural(pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{X}^\natural)), V_{\mathfrak{B}^\natural} := W_{\mathfrak{B}^\natural}/U_{\mathfrak{B}^\natural}$ and $[\lambda^\natural](Y^\natural)$). Now $V_{\mathfrak{B}^\natural}$ is a subgroup of W and $\lambda^\natural(pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{X}^\natural)) = V_{\mathfrak{B}^\natural}(\mathbb{C})\tilde{z}^\natural$ by Proposition 9.15.

We have $pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{X}^\natural)_V = V_{\mathfrak{B}^\natural}(\mathbb{C}) + \tilde{z}_V^\natural$. Every fiber of $\lambda^\natural(pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{X}^\natural)) \rightarrow pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{X}^\natural)_V$ is a point, so

$$(9.20) \quad pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{Z}^\natural) = pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{X}^\natural) \cap \pi^{\natural,-1}([pr]_{F^\natural}^{\text{ws}}(Y^\natural)_V).$$

The monodromy group $\Gamma_{Y_V^\natural}$ of $Y_V^\natural$, which stabilizes $[pr]_{F^\natural}^{\text{ws}}(Y^\natural)_V$, is Zariski dense in $V_{F^\natural}$, so $\Gamma_{Y_V^\natural}$ is of finite index in $\Gamma_{V_{F^\natural}}$ because $V_{F^\natural}$ is a vector group. Up to replacing Γ_V by a subgroup of finite index, we may assume $\Gamma_{Y_V^\natural} = \Gamma_{V_{F^\natural}}$. Let $\Gamma'_W := \pi^{\natural,-1}(\Gamma_{V_{F^\natural}}) \cap \Gamma_W$. Then (9.20) implies

$$(9.21) \quad [pr]_{F^\natural}^{\text{ws}}(Z)_{z_V^\natural} = \left(\text{unif}(\Gamma'_W \tilde{Z}^\natural) \right)_{z_V^\natural} = \tilde{z}_U^\natural + \frac{1}{2} \Psi(\Gamma_{V_{F^\natural}}, \tilde{z}_V^\natural) + \Gamma_U \pmod{\Gamma_U}.$$

Recall the construction of $U^\dagger$ (Step IV of the proof of Theorem 9.8, above (9.11)): identify $V(\mathbb{C})/F^0V(\mathbb{C})$ with $V(\mathbb{R})$ by $i: V(\mathbb{R}) \subset V(\mathbb{C}) \rightarrow V(\mathbb{C})/F^0V(\mathbb{C})$ and then $U^\dagger$ is the smallest subgroup of U such that $\Psi(\Gamma_{V_{F^\natural}}, i^{-1}(\pi_V^\natural(\tilde{z}_V^\natural))) \subset U(\mathbb{Q}) + U_{\mathfrak{B}^\natural}(\mathbb{R}) + U^\dagger(\mathbb{R})$ (here we used the facts $U_{\mathfrak{B}^\natural} = U_{\mathfrak{B}}$ by Proposition 9.15 and $V_{F^\natural} = V_F$). The assumption on $U^\dagger$ implies that for any coordinate (i.e. any 1-dimensional quotient U_0) of $U^\dagger(\mathbb{C})$, there exists a $u \in \Psi(\Gamma_{V_{F^\natural}}, \tilde{z}_V^\natural)$ such that the chosen coordinate of u , which we call u_0 , has irrational real part. But then $\text{unif}(\tilde{z}_U^\natural + U_0(\mathbb{R})) \subset [pr]_{F^\natural}^{\text{ws}}(Z)_{z_V^\natural}^{\text{Zar}}$ since $\{lu_0 \bmod \mathbb{Z} | l \in \mathbb{Z}\}$ is dense in $[0, 1)$. So $[pr]_{F^\natural}^{\text{ws}}(Z)_{z_V^\natural}^{\text{Zar}}$, which is complex analytic, contains $\text{unif}^\natural(\tilde{z}_U^\natural + U_0(\mathbb{C}))$, and so does $[pr]_{F^\natural}^{\text{ws}}(Y^\natural)_{z_V^\natural}$ since $Y^\natural = Z^{\natural, \text{Zar}}$. Hence the conclusion follows by varying U_0 (which is any coordinate of $U^\dagger$). $\square$

Now we are ready to prove Theorem 9.1, i.e. (9.13) and (9.14).

Proof of (9.13).

$$\begin{aligned} & \dim \mathbf{pr}_{F^\natural}^{\text{ws}}(\mathfrak{B}^\natural) - \dim \mathbf{pr}_{F^\natural}^{\text{ws}}(\mathfrak{Z}^\natural) - \dim(F^\natural)^{\text{ws}} \\ &= \dim U_{\mathfrak{B}^\natural}(\mathbb{C}) + \dim pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{X}^\natural)_V + \dim [pr]_{F^\natural}^{\text{ws}}(Y^\natural) - \dim \mathbf{pr}_{F^\natural}^{\text{ws}}(\mathfrak{Z}^\natural) - \dim(F^\natural)^{\text{ws}} \quad \text{by Proposition 9.15} \\ &= (\dim U_{\mathfrak{B}^\natural}(\mathbb{C}) + \dim pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{X}^\natural)_V) + (\dim T^\dagger + \dim[\rho^\natural]([pr]_{F^\natural}^{\text{ws}}(Y^\natural))) - \dim \mathbf{pr}_{F^\natural}^{\text{ws}}(\mathfrak{Z}^\natural) - \dim(F^\natural)^{\text{ws}} \\ & \hspace{15em} \text{by Proposition 9.18} \\ &\geq \dim([\rho^\natural]([pr]_{F^\natural}^{\text{ws}}(Y^\natural)) \rightarrow [pr]_{F^\natural}^{\text{ws}}(Y^\natural)_V) - \dim(pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{Z}^\natural) \rightarrow pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{Z}^\natural)_V) \\ & \hspace{10em} \text{since } U_{\mathfrak{B}^\natural} \oplus U^\dagger = U_F \text{ and (9.13) is true for } A^\natural \\ &= \dim([\rho^\natural]([pr]_{F^\natural}^{\text{ws}}(Y^\natural)) \rightarrow [pr]_{F^\natural}^{\text{ws}}(Y^\natural)_V) - \dim(\rho^\natural(pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{Z}^\natural)) \rightarrow pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{Z}^\natural)_V) \quad \text{since } U_{\mathfrak{B}^\natural} \cap U^\dagger = 0 \\ &\geq 0. \end{aligned}$$

$\square$

Proof of (9.14). Let L_0 be a fiber of $\tilde{X}^\natural \rightarrow pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{X}^\natural)$. Then L_0 is an algebraic subvariety of $\mathbb{C}^k$ for some k . We have $V_{\mathcal{B}^\natural}(\mathbb{C}) + L_0 \subset \tilde{X}_V^\natural \subset V_F(\mathbb{C}) + L_0$. By taking the Zariski closure of the images under $\text{unif}_V^\natural$, we get by Proposition 9.15 (recall that $\tilde{F}^\natural$ is the smallest bi-algebraic subset of $W(\mathbb{C})$ containing $\tilde{X}^\natural$) $(F^\natural)_V^{\text{ws}} \times L_0 \subset F_V^\natural \subset (F^\natural)_V^{\text{ws}} \times L_0$. So $F_V^\natural = (F^\natural)_V^{\text{ws}} \times L_0$. Therefore $F^\natural = (F^\natural)^{\text{ws}} \times L_0$. Now we have

$$\begin{aligned} \dim \tilde{X}^\natural + \dim Y^\natural - \dim \tilde{Z}^\natural &= (\dim pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{X}^\natural) + \dim L_0) + \dim Y^\natural - \dim \tilde{Z}^\natural \\ &= (\dim pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{X}^\natural) + \dim [pr]_{F^\natural}^{\text{ws}}(Y^\natural) - \dim pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{Z}^\natural)) + \dim L_0 \\ &\quad + \dim(Y^\natural \rightarrow [pr]_{F^\natural}^{\text{ws}}(Y^\natural)) - \dim(\tilde{Z}^\natural \rightarrow pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{Z}^\natural)) \\ &\geq \dim(F^\natural)^{\text{ws}} + \dim L_0 + \dim(Y^\natural \rightarrow [pr]_{F^\natural}^{\text{ws}}(Y^\natural)) - \dim(\tilde{Z}^\natural \rightarrow pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{Z}^\natural)) \\ &\geq \dim F^\natural \end{aligned}$$

where the last but second inequality follows from part (2) of Ax-Schanuel, i.e. (9.13), and the discussion below Conjecture 8.1. $\square$

10. SPECIAL SUBVARIETIES

Let $S^\natural$ be a connected enlarged mixed Shimura variety associated with $(P, \mathcal{X}^{\natural+})$ and let $\text{unif}^\natural: \mathcal{X}^{\natural+} \rightarrow S = \Gamma \backslash \mathcal{X}^{\natural+}$ be the uniformization. Use notation of §3.4.

Definition 10.1. (1) A subset $\tilde{Y}^\natural$ of $\mathcal{X}^{\natural+}$ is said to be **special** if it is the underlying space of some connected enlarged mixed Shimura subdatum $(Q, \mathcal{Y}^{\natural+}) \hookrightarrow (P, \mathcal{X}^{\natural+})$ (in the category $\mathcal{EMSD}$). **Special points** of $\mathcal{X}^{\natural+}$ are precisely the special subsets of dimension 0.

(2) A subset $Y^\natural$ of $S^\natural$ is called a **special subvariety** if it is the image of some subset of $\mathcal{X}^{\natural+}$ under $\text{unif}^\natural$ (Proposition 4.6(1) implies that any such defined subset of $S^\natural$ is a closed algebraic subvariety of $S^\natural$). **Special points** of $S^\natural$ are precisely the special subvarieties of dimension 0.

Remark 10.2. The definition of morphisms $\mathcal{EMSD}$ (Definition 3.4) yields the following direct corollary: under the geometric comparison $\pi^\natural: \mathcal{X}^{\natural+} \rightarrow \varphi(\mathcal{X}^+)$ (see §3.4), there is a bijection

$$\{\text{special subsets of } \mathcal{X}^{\natural+}\} \xrightarrow{\sim} \{\text{special subsets of } \varphi(\mathcal{X}^+)\}, \quad \tilde{Y}^\natural \mapsto \tilde{Y} := \pi^\natural(\tilde{Y}^\natural),$$

which induces $\{\text{special points of } \mathcal{X}^{\natural+}\} \xrightarrow{\sim} \{\text{special points of } \varphi(\mathcal{X}^+)\}$.

Theorem 10.3. Let $Y^\natural$ be a special subvariety of $S^\natural$. Then the set of special points contained in $Y^\natural$ is Zariski dense.

Proof. Use Notation 5.3. Then Y_G is a special subvariety of S_G and, by the theory of Shimura varieties, the set of special points of S_G contained in Y_G is Zariski dense. Because $Y^\natural$ is special, we have that $Y^\natural$ is a torsor over the universal vector extension of the abelian scheme $Y_{P/U}/Y_G$, and hence the fibers of $Y^\natural \rightarrow Y_G$ are of constant dimension, say n .

Let $Z^\natural \subset Y^\natural$ be the Zariski closure of the set of special points of $S^\natural$ contained in $Y^\natural$. Let $y_G \in Y_G$ be a special point of S_G , then the set of special points of $S^\natural$ contained in the fiber $Y_{y_G}^\natural$ is Zariski dense in $Y_{y_G}^\natural$: to see this it suffices to prove that the set of special points of $S_{P/U}^\natural$ contained in $Y_{P/U, y_G}^\natural$ is Zariski dense, which is true because $Y_{P/U, y_G}^\natural$ is the universal vector

extension of the abelian variety $Y_{P/U, y_G}$. Hence $\dim Z_{y_G}^\natural = \dim Y_{y_G}^\natural = n$ for any special point $y_G \in Y_G$. However the semi-continuity implies that the set

$$\Sigma := \{y_G \in Y_G \mid \dim Z_{y_G}^\natural \geq n\},$$

which contains all special points of Y_G by the argument above, is Zariski closed in Y_G . Hence $\Sigma = Y_G$. Therefore $Z^\natural = Y^\natural$. $\square$

The converse of Theorem 10.3 is an ‘‘André-Oort’’ type conjecture.

Conjecture 10.4 (André-Oort for enlarged mixed Shimura varieties). *Let $Y^\natural$ be an irreducible subvariety of a connected enlarged mixed Shimura variety $S^\natural$. If the set of special points of $S^\natural$ contained in $Y^\natural$ is Zariski dense in $Y^\natural$, then $Y^\natural$ is a special subvariety.*

Conjecture 10.4 implies directly the mixed André-Oort conjecture:

Conjecture 10.5 (mixed André-Oort). *Let Y be an irreducible subvariety of a connected mixed Shimura variety S . If the set of special points of S contained in Y is Zariski dense in Y , then Y is a special subvariety of S .*

But in fact, we have

Lemma 10.6. *Conjecture 10.4 and Conjecture 10.5 are equivalent.*

Proof. We only need to prove Conjecture 10.4 assuming Conjecture 10.5. Let $Y^\natural$ be as in Conjecture 10.4 and let $\Sigma^\natural$ be the set of special points of $S^\natural$ contained in $Y^\natural$. Use Notation 5.3. Then $\Sigma := [\pi^\natural](\Sigma^\natural)$ is a set of special points of S such that $\Sigma^{\text{Zar}} = Y$. Hence Y is a special subvariety of S by Conjecture 10.5.

Let $Z^\natural$ be the special subvariety of $S^\natural$ which maps to Y under the bijection

$$\{\text{special subvarieties of } S^\natural\} \xrightarrow{\sim} \{\text{special subvarieties of } S\}$$

given by Remark 10.2. But $[\pi^\natural]: \Sigma^\natural \rightarrow \Sigma$ is a bijection by Remark 10.2. So the set of special points of $S^\natural$ contained in $Z^\natural$ is $\Sigma^\natural$. Now Theorem 10.3 implies that $(\Sigma^\natural)^{\text{Zar}} = Z^\natural$. Hence $Y^\natural = Z^\natural$ is special. $\square$

Remark 10.7. *A key point to prove Lemma 10.6 is that the bijection in Remark 10.2 preserves 0-dimensional special subvarieties. In general this is not true: an r -dimensional special subvariety of $S^\natural$ is often mapped to a special subvariety of S of smaller dimension.*

Conjecture 10.5 has been intensively studied. It is proven unconditionally for any mixed Shimura variety of abelian type (i.e. its pure part is of abelian type) by Pila-Tsimerman [31], Tsimerman [38] and Gao [16]. It is also proven under the generalized Riemann Hypothesis (for CM fields) for all mixed Shimura varieties by Klingler-Ullmo-Yafaev [21], Ullmo [39], Ullmo-Yafaev [42], Daw-Orr [13] and Gao [15]. Then Lemma 10.6 implies the corresponding results for enlarged mixed Shimura varieties.

Taking into account of not only special points but also special subvarieties of higher dimensions, we have a ‘‘Zilber-Pink’’ type conjecture.

Conjecture 10.8 (Zilber-Pink for enlarged mixed Shimura varieties). *Let $Y^\natural$ be an irreducible subvariety of a connected enlarged mixed Shimura variety $S^\natural$. Assume that $S^\natural$ has no proper*

special subvariety which contains $Y^\natural$. Then

$$\bigcup_{\substack{S^{\natural'} \text{ special,} \\ \dim(S^{\natural'}) < \text{codim}(Y^\natural)}} S^{\natural'} \cap Y^\natural$$

is not Zariski dense in $Y^\natural$.

The Zilber-Pink conjecture is naturally a generalization of the André-Oort conjecture. Unlike the André-Oort conjecture, there is no immediate equivalence (as given by Lemma 10.6) between Conjecture 10.8 and the Zilber-Pink conjecture for mixed Shimura varieties (see Remark 10.7). There are some results about this conjecture in some special cases (Habegger-Pila [18, 19], Orr [26], Gao [17]), but in general little is known.

We finish this paper by the following lemma, which explains the relation between special and weakly special subvarieties (Definition 5.10).

Lemma 10.9. *A subset $\tilde{Y}^\natural$ of $\mathcal{X}^{\natural+}$ is special if and only if $\tilde{Y}^\natural$ is weakly special and contains a special point. Equivalently, a subvariety $Y^\natural$ of $S^\natural$ is special if and only if it is weakly special and contains a special point.*

Proof. The “only if” part is clear. We prove the “if” part. Let $(Q, \mathcal{Y}^{\natural+})$ be as in Definition 5.10(1) defining $\tilde{Y}^\natural$ and let $N := \text{Ker}\varphi$. Let $(T, \tilde{y}^\natural)$ be the connected enlarged mixed Shimura datum giving a special point contained in $\tilde{Y}^\natural$. Let Q' be the subgroup of P generated by $i(N)$ and T . Then $(Q', Q'(\mathbb{R})^+ W_{Q'}(\mathbb{C}) \tilde{y}^\natural)$ defines a connected enlarged mixed Shimura subdatum of $(P, \mathcal{X}^{\natural+})$ (in the category $\underline{\mathcal{EMSD}}$), and $\tilde{Y}^\natural = Q'(\mathbb{R})^+ W_{Q'}(\mathbb{C}) \tilde{y}^\natural$. Hence $\tilde{Y}^\natural$ is special. $\square$

11. APPENDIX: PROOF OF AX-LINDEMANN (THEOREM 7.1)

11.1. Outline. We prove the Ax-Lindemann theorem (in the form Theorem 7.1) in the rest of the section. The proof is a modification of the author’s previous work on the Ax-Lindemann theorem for mixed Shimura varieties [16, §9-11]. Consider the diagram

$$\begin{array}{ccccc} (P, \mathcal{X}^{\natural+}) & \xrightarrow{\pi^\natural} & (P, \varphi(\mathcal{X}^+)) & \xrightarrow{\pi} & (G, \varphi(\mathcal{X}^+)_G), \\ \downarrow \text{unif}^\natural & & \downarrow \text{unif} & & \downarrow \text{unif}_G \\ S^\natural & \xrightarrow{[\pi^\natural]} & S & \xrightarrow{[\pi]} & S_G, \\ & \xleftarrow{[\pi^\natural]} & & & \end{array} \quad \begin{array}{ccccc} \tilde{Z}^\natural & \longrightarrow & \tilde{Z} & \longrightarrow & \tilde{Z}_G \\ \downarrow & & \downarrow & & \downarrow \\ Z^\natural & \longrightarrow & Z & \longrightarrow & Z_G \end{array}$$

and use Notation 5.3. First of all we do the following reduction:

- The conclusion is obviously true if $\tilde{Z}$ is a point. Hence we may assume $\dim \tilde{Z} > 0$.
- Let $(P', \mathcal{X}'^{\natural+})$ be the smallest connected enlarged mixed Shimura subdatum of $(P, \mathcal{X}^{\natural+})$ such that $\tilde{Z}^\natural \subset \mathcal{X}'^{\natural+}$ and let $S'^\natural$ be the corresponding connected enlarged mixed Shimura subvariety of $S^\natural$. Then $(P', \mathcal{X}'^{\natural+})$ has generic Mumford-Tate group by Proposition 3.12. Replace $(P, \mathcal{X}^{\natural+})$ by $(P', \mathcal{X}'^{\natural+})$, $S^\natural$ by $S'^\natural$, $\text{unif}^\natural$ by $\text{unif}'^\natural: \mathcal{X}'^{\natural+} \rightarrow S'^\natural$ and $Y^\natural$ by the irreducible component $Y'^\natural$ of $Y^\natural \cap S'^\natural$ such that $\tilde{Z}^\natural \subset (\text{unif}'^\natural)^{-1}(Y'^\natural)$, then $\tilde{Z}^\natural$ is an irreducible algebraic subset of $\mathcal{X}'^{\natural+}$ contained in $(\text{unif}'^\natural)^{-1}(Y'^\natural)$ and is maximal

for these properties. By definition, $\tilde{Z}^\natural$ is bi-algebraic in $\mathcal{X}'^{\natural,+}$ if and only if it is bi-algebraic in $\mathcal{X}^{\natural,+}$. So making the replacement above does not change the assumption or the conclusion of the theorem.

- Replace $(P, \mathcal{X}^{\natural,+})$ by $(P, \mathcal{X}^{\natural,+})/Z(P) = (P/Z(P), \mathcal{X}^{\natural,+})$ and other objects accordingly.
- Let $Y_0^\natural := \text{unif}^\natural(\tilde{Z}^\natural)^{\text{Zar}}$, then $\tilde{Z}^\natural$ is an irreducible algebraic subset of $\mathcal{X}^{\natural,+}$ contained in $\text{unif}^{\natural-1}(Y_0^\natural)$ and is maximal for these properties. Hence we may assume $Y^\natural = Y_0^\natural$.
- After the previous reduction, there is a unique complex analytic irreducible component of $\text{unif}^{\natural-1}(Y^\natural)$ containing $\tilde{Z}^\natural$, which we denote by $\tilde{Y}^\natural$.
- Replace Γ by a neat subgroup contained in $P^{\text{der}}(\mathbb{Q})$ (Remark 5.2(3)).

Let $\tilde{F}^\natural$ be the smallest quasi-linear subset of $\mathcal{X}^{\natural,+}$ containing $\tilde{Y}^\natural$. Let N be the connected algebraic monodromy group associated with $Y^{\natural, \text{sm}}$ and let $W_N := W \cap N$ ($U_N := U \cap N$). Denote by $\tilde{F} := \pi^\natural(\tilde{F}^\natural)$ and $\tilde{F}_G := \pi(\tilde{F})$. Then by Theorem 6.3, we have

$$\tilde{F} = N(\mathbb{R})^+ U_N(\mathbb{C}) \tilde{z} \text{ and } \tilde{F}_G = G_N(\mathbb{R})^+ \tilde{z}_G$$

where $\tilde{z} \in \tilde{Z}$ and $\tilde{z}_G = \pi(\tilde{z})$.

Let $(Q, \mathcal{Y}^+)$ be the smallest connected mixed Shimura subdatum of $(P, \mathcal{X}^+)$ such that $\tilde{Z} \subset \mathcal{Y}^+$. Let $\Gamma_Q := \Gamma \cap Q(\mathbb{Q})$. Then $\tilde{F}$ is Hodge generic in $\mathcal{Y}^+$, $N \triangleleft Q$ and $N \triangleleft Q^{\text{der}}$. Define

$$\Gamma_{\tilde{Z}^\natural} := \{\gamma \in \Gamma_Q \mid \gamma \cdot \tilde{Z}^\natural = \tilde{Z}^\natural\} \text{ and } \Gamma_{G_Q, \tilde{Z}_G} := \{\gamma \in \Gamma_{G_Q} \mid \gamma \cdot \tilde{Z}_G = \tilde{Z}_G\}$$

and $H_{\tilde{Z}^\natural} := (\Gamma_{\tilde{Z}^\natural}^{\text{Zar}})^\circ$, $H_{\tilde{Z}_G} := (\Gamma_{G_Q, \tilde{Z}_G}^{\text{Zar}})^\circ$. Then $H_{\tilde{Z}^\natural}$ (resp. $H_{\tilde{Z}_G}$) is the largest connected subgroup of Q^{der} (resp. G^{der}) such that $H_{\tilde{Z}^\natural}(\mathbb{R})^+ W_{H_{\tilde{Z}^\natural}}(\mathbb{C})$ (resp. $H_{\tilde{Z}_G}(\mathbb{R})^+$) stabilizes $\tilde{Z}^\natural$ (resp. $\tilde{Z}_G$).

Define $U_{H_{\tilde{Z}^\natural}} := U \cap H_{\tilde{Z}^\natural}$ and $W_{H_{\tilde{Z}^\natural}} := W \cap H_{\tilde{Z}^\natural}$. Both of them are normal in $H_{\tilde{Z}^\natural}$. Define $V_{H_{\tilde{Z}^\natural}} := W_{H_{\tilde{Z}^\natural}}/U_{H_{\tilde{Z}^\natural}}$ and $G_{H_{\tilde{Z}^\natural}} := H_{\tilde{Z}^\natural}/W_{H_{\tilde{Z}^\natural}}$.

Remark 11.1. *We do NOT know whether $W_{H_{\tilde{Z}^\natural}}(\mathbb{R}) U_{H_{\tilde{Z}^\natural}}(\mathbb{C})$ stabilizes $\tilde{Z}$. This can be seen by restricting to a fiber: the morphism $\pi^\natural$ then becomes $W(\mathbb{C}) \rightarrow W(\mathbb{C})/F^0 W(\mathbb{C})$ and a $W_{H_{\tilde{Z}^\natural}}(\mathbb{C})$ -orbit becomes a $W_{H_{\tilde{Z}^\natural}}(\mathbb{C})/(W_{H_{\tilde{Z}^\natural}}(\mathbb{C}) \cap F^0 W(\mathbb{C}))$ -orbit, but NOT a $W_{H_{\tilde{Z}^\natural}}(\mathbb{R}) U_{H_{\tilde{Z}^\natural}}(\mathbb{C})$ -orbit!*

Lemma 11.2. *The set $\tilde{Y}^\natural$ is stable under $H_{\tilde{Z}^\natural}(\mathbb{R})^+ W_{H_{\tilde{Z}^\natural}}(\mathbb{C})$.*

Proof. Every fiber of $\mathcal{X}^{\natural,+} \rightarrow \varphi(\mathcal{X}^+)_G$ can be canonically identified with $W(\mathbb{C})$. So it is enough to prove that $\tilde{Y}$ is stable under $H_{\tilde{Z}}(\mathbb{R})^+$: If $W_{H_{\tilde{Z}^\natural}}(\mathbb{R}) \tilde{y} \subset \tilde{Y}^\natural$ for $\tilde{y}^\natural \in \tilde{Y}^\natural$, then $W_{H_{\tilde{Z}^\natural}}(\mathbb{C}) \tilde{y}^\natural \subset \tilde{Y}^\natural$ because $\tilde{Y}^\natural$ is complex analytic and $W_{H_{\tilde{Z}^\natural}}(\mathbb{C}) \tilde{y}^\natural$ is the smallest complex analytic subset of $\mathcal{X}^{\natural,+}$ containing $W_{H_{\tilde{Z}^\natural}}(\mathbb{R}) \tilde{y}^\natural$.

If not, then since $H_{\tilde{Z}^\natural}(\mathbb{Q})$ is dense (in the usual topology) in $H_{\tilde{Z}^\natural}(\mathbb{R})^+$, there exists $h \in H_{\tilde{Z}^\natural}(\mathbb{Q})$ such that $h \tilde{Y}^\natural \neq \tilde{Y}^\natural$. Then $\tilde{Z}^\natural \subset \tilde{Y}^\natural \cap h \tilde{Y}^\natural$, and hence contained in a complex analytic irreducible component $\tilde{Y}'^\natural$ of it. Consider the Hecke correspondence T_h

$$\Gamma \backslash \mathcal{X}^{\natural,+} \xleftarrow{[h]} (\Gamma \cap h^{-1} \Gamma h) \backslash \mathcal{X}^{\natural,+} \xrightarrow{[1]} \Gamma \backslash \mathcal{X}^{\natural,+}$$

Then $T_h(Y^\natural) = \text{unif}^\natural(h \cdot \text{unif}^{\natural-1}(Y^\natural))$. Hence

$$Y^\natural \cap T_h(Y^\natural) = \text{unif}^\natural(\text{unif}^{\natural-1}(Y^\natural) \cap (h \cdot \text{unif}^{\natural-1}(Y^\natural))).$$

On the other hand, $T_h(Y^\natural)$ is equidimensional of the same dimension as $Y^\natural$, hence by reason of dimension, $h\tilde{Y}^\natural$ is an irreducible component of $\text{unif}^{\natural-1}(T_h(Y^\natural)) = \Gamma h\Gamma\tilde{Y}^\natural$. So $\text{unif}(h\tilde{Y}^\natural)$ is an irreducible component of $T_h(Y^\natural)$.

Since $\tilde{Y}^{\natural, \prime}$ is a complex analytic irreducible component of $\tilde{Y}^\natural \cap h\tilde{Y}^\natural$, it is also a complex analytic irreducible component of $\text{unif}^{\natural-1}(Y^\natural) \cap (h\tilde{Y}^\natural) = \Gamma\tilde{Y}^\natural \cap h\tilde{Y}^\natural$. So $Y^{\natural, \prime} := \text{unif}^\natural(\tilde{Y}^{\natural, \prime})$ is a complex analytic irreducible component of $Y^\natural \cap \text{unif}(h\tilde{Y}^\natural)$. So $Y^{\natural, \prime}$ is a complex analytic irreducible component of $Y^\natural \cap T_h(Y^\natural)$, and hence is algebraic since $Y^\natural \cap T_h(Y^\natural)$ is.

Since $h\tilde{Y}^\natural \neq \tilde{Y}^\natural$ and $Y^\natural$ is irreducible, $\dim(Y^{\natural, \prime}) < \dim(Y^\natural)$. But $\tilde{Z}^\natural \subset \tilde{Y}^\natural \cap h\tilde{Y}^\natural \subset \text{unif}^{\natural-1}(Y^{\natural, \prime})$. This contradicts the minimality of $Y^\natural$. $\square$

Lemma 11.3. $H_{\tilde{Z}^\natural} \triangleleft N$.

Proof. We have $\tilde{Z}^\natural \subset \tilde{F}^\natural$. Consider $pr_{\tilde{F}^\natural}^{\text{ws}}$, the projection of $\tilde{F}^\natural$ to its weakly special part $(\tilde{F}^\natural)^{\text{ws}}$ (see (5.9)). Let $(Q, \mathcal{Y}^{\natural+})$ be the enlarged mixed Shimura subdatum of $(P, \mathcal{X}^{\natural+})$ associated to $(Q, \mathcal{Y}^+)$. Then the image of $pr_{\tilde{F}^\natural}^{\text{ws}}(\tilde{Z}^\natural)$ under $(Q, \mathcal{Y}^{\natural+}) \rightarrow (Q, \mathcal{Y}^{\natural+})/N$ is a Hodge generic point (since $\tilde{F}$ is Hodge generic in $\mathcal{Y}^+$), which is stable under $H_{\tilde{Z}^\natural}/(H_{\tilde{Z}^\natural} \cap N)$. So $H_{\tilde{Z}^\natural} < N$.

Let H' be the algebraic group generated by $\gamma^{-1}H_{\tilde{Z}^\natural}\gamma$ for all $\gamma \in \Gamma_{Y^{\natural, \text{sm}}}$, where $\Gamma_{Y^{\natural, \text{sm}}}$ is the monodromy group of $Y^{\natural, \text{sm}}$. Since H' is invariant under conjugation by $\Gamma_{Y^{\natural, \text{sm}}}$, it is invariant under $(\Gamma_{Y^{\natural, \text{sm}}})^{\text{Zar}}$, therefore invariant under conjugation by N .

By Lemma 11.2, $\tilde{Y}^\natural$ is invariant under $H_{\tilde{Z}^\natural}(\mathbb{R})^+W_{H_{\tilde{Z}^\natural}}(\mathbb{C})$. On the other hand, $\tilde{Y}^\natural$ is also invariant under $\Gamma_{Y^{\natural, \text{sm}}}$. So $\tilde{Y}^\natural$ is invariant under the action of $H'(\mathbb{R})^+W_{H'}(\mathbb{C})$ where $W_{H'} := W \cap H'$. Since $H'(\mathbb{R})^+W_{H'}(\mathbb{C})\tilde{Z}^\natural$ is semi-algebraic, there exists an irreducible algebraic subset of $\mathcal{X}^{\natural+}$, say $\tilde{E}^\natural$, which contains $H'(\mathbb{R})^+W_{H'}(\mathbb{C})\tilde{Z}^\natural$ and is contained in $\tilde{Y}^\natural$ by [30, Lemma 4.1]. Now $\tilde{Z}^\natural \subset \tilde{E}^\natural \subset \tilde{Y}^\natural$, so $\tilde{Z}^\natural = \tilde{E}^\natural = H'(\mathbb{R})^+W_{H'}(\mathbb{C})\tilde{Z}^\natural$ by maximality of $\tilde{Z}^\natural$, and therefore $H' = H_{\tilde{Z}^\natural}$ by the maximality of $H_{\tilde{Z}^\natural}$. So $H_{\tilde{Z}^\natural}$ is invariant under conjugation by N . But $H_{\tilde{Z}^\natural} < N$, so $H_{\tilde{Z}^\natural}$ is normal in N . $\square$

Corollary 11.4. $G_{H_{\tilde{Z}^\natural}} \triangleleft G_Q^{\text{der}}$.

Proof. We have $G_{H_{\tilde{Z}^\natural}} \triangleleft G_N \triangleleft G_Q^{\text{der}}$, and so $G_{H_{\tilde{Z}^\natural}} \triangleleft G_Q^{\text{der}}$ since all the three groups are reductive. $\square$

Lemma 11.5. (1) Y is weakly special. Hence $\tilde{Y} = \tilde{F} = N(\mathbb{R})^+\tilde{y}$ for any $\tilde{y} \in \tilde{Y}$.

(2) $\text{unif}(\tilde{Z})^{\text{Zar}} = Y$.

Proof. We have $\text{unif}(\tilde{Z})^{\text{Zar}} = Y$ because $Y^\natural = \text{unif}^\natural(\tilde{Z}^\natural)^{\text{Zar}}$. The Ax-Lindemann for mixed Shimura varieties (originally proven as [16, Theorem 1.2], but we use the equivalent statement [14, Theorem 3.1.3]) implies that $Y = Z^{\text{Zar}}$ is weakly special. $\square$

Proposition 11.6. $\tilde{Z}_G = G_{H_{\tilde{Z}^\natural}}(\mathbb{R})^+\tilde{z}_G$. Hence $\tilde{Z}_G$ is weakly special.

The proof of Proposition 11.6 will occupy §11.2. Here we show how to prove the Ax-Lindemann theorem assuming this proposition. The following theorem is a direct corollary of Theorem 9.1 and Theorem 8.5.

Theorem 11.7. The Ax-Lindemann theorem (Theorems 7.1~7.4) is true if $\tilde{Z}_G$ is a point.

Proof of Theorem 7.1. We divide the proof into the following steps.

Step I Prove $\tilde{Z}^\natural = H_{\tilde{Z}^\natural}(\mathbb{R})^+ W_{H_{\tilde{Z}^\natural}}(\mathbb{C})(\tilde{Z}^\natural)_{\tilde{z}_G}$ for some $\tilde{z}_G \in \tilde{Z}_G$.

Consider a fibre of $\tilde{Z}^\natural$ over a Hodge-generic point $\tilde{z}_G \in \tilde{Z}_G$ such that $\pi^\natural|_{\tilde{Z}^\natural}$ is flat at $\tilde{z}_G$ (such a point exists by [1, §4, Lemma 1.4] and generic flatness). Suppose that $\tilde{K}^\natural$ is an irreducible algebraic component of $\tilde{Z}_{\tilde{z}_G}^\natural$ such that $\dim(\tilde{Z}_{\tilde{z}_G}^\natural) = \dim(\tilde{K}^\natural)$, then the flatness of $\pi^\natural|_{\tilde{Z}^\natural}$ at $\tilde{z}_G$ implies $\dim(\tilde{Z}^\natural) = \dim(\tilde{Z}_G) + \dim(\tilde{Z}_{\tilde{z}_G}^\natural) = \dim(\tilde{Z}_G) + \dim(\tilde{K}^\natural)$.

Consider the set $\tilde{E}^\natural := H_{\tilde{Z}^\natural}(\mathbb{R})^+ W_{H_{\tilde{Z}^\natural}}(\mathbb{C})\tilde{K}^\natural$. It is semi-algebraic (since $\tilde{K}$ is algebraic and the action of $P(\mathbb{R})^+ W(\mathbb{C})$ on $\mathcal{X}^{\natural,+}$ is algebraic). The fact $\tilde{K}^\natural \subset \tilde{Z}^\natural$ implies that $\tilde{E}^\natural \subset \tilde{Z}^\natural$. Now Proposition 11.6 implies that $\pi^\natural(\tilde{E}^\natural) = G_{H_{\tilde{Z}^\natural}}(\mathbb{R})^+ \tilde{z}_G = \tilde{Z}_G$ and that the $\mathbb{R}$ -dimension of every fiber of $\pi^\natural|_{\tilde{E}^\natural}$ is at least $\dim_{\mathbb{R}}(\tilde{K}^\natural)$. So

$$\dim(\tilde{Z}^\natural) \geq \dim(\tilde{E}^\natural) \geq \dim(\pi^\natural(\tilde{E}^\natural)) + \dim(\tilde{K}^\natural) = \dim(\tilde{Z}_G) + \dim(\tilde{K}^\natural) = \dim(\tilde{Z}^\natural).$$

So $\tilde{E}^\natural = \tilde{Z}^\natural$ since $\tilde{Z}^\natural$ is irreducible.

Next let $\tilde{K}^{\natural,\prime}$ be an irreducible algebraic subset which contains $\tilde{Z}_{\tilde{z}_G}^\natural$ and is contained in $\text{unif}^{\natural,-1}(Y^\natural)_{\tilde{z}_G}$, maximal for these properties. Then $\tilde{K}^{\natural,\prime}$ is quasi-linear by Theorem 11.7. We have $\tilde{K}^{\natural,\prime} \subset \tilde{Y}^\natural$ since $\tilde{Y}^\natural$ is an irreducible component of $\pi^{\natural,-1}(Y^\natural)$. Consider $\tilde{E}^{\natural,\prime} := H_{\tilde{Z}^\natural}(\mathbb{R})^+ W_{H_{\tilde{Z}^\natural}}(\mathbb{C})\tilde{K}^{\natural,\prime}$. Then $\tilde{E}^{\natural,\prime} \subset \tilde{Y}^\natural$ by Lemma 11.2. But $\tilde{E}^{\natural,\prime}$ is semi-algebraic, so by [30, Lemma 4.1], there exists an irreducible algebraic subset of $\mathcal{X}^{\natural,+}$, say $\tilde{E}_{\text{alg}}^{\natural,\prime}$ which contains $\tilde{E}^{\natural,\prime}$ and is contained in $\tilde{Y}^\natural$. So $\tilde{Z}^\natural = \tilde{E}^\natural \subset \tilde{E}_{\text{alg}}^{\natural,\prime} \subset \tilde{Y}^\natural$. Now the maximality of $\tilde{Z}^\natural$ implies that

$$(11.1) \quad \tilde{Z}^\natural = \tilde{E}_{\text{alg}}^{\natural,\prime} = \tilde{E}^{\natural,\prime} = H_{\tilde{Z}^\natural}(\mathbb{R})^+ W_{H_{\tilde{Z}^\natural}}(\mathbb{C})\tilde{K}^{\natural,\prime} \text{ and } \tilde{K}^{\natural,\prime} = \tilde{Z}_{\tilde{z}_G}^\natural.$$

Step II Prove $\tilde{Z} = H_{\tilde{Z}^\natural}(\mathbb{R})^+ U_{H_{\tilde{Z}^\natural}}(\mathbb{C})\tilde{z}$.

Since $\tilde{K}^{\natural,\prime}$ is quasi-linear, we can write $\tilde{K}^{\natural,\prime} := \pi^\natural(\tilde{K}^{\natural,\prime}) = W'(\mathbb{R})U'(\mathbb{C})\tilde{z}$ with $W' < W$, $U' = U \cap W'$ and $\tilde{z} \in \tilde{Z}_{\tilde{z}_G}$. Then $W_{H_{\tilde{Z}^\natural}} < W'$ and $\tilde{K}^{\natural,\prime}$ is stable under $W'(\mathbb{C})$. The complex structure of $\pi^{-1}(\tilde{z}_G)$ comes from $W(\mathbb{R})U(\mathbb{C}) \simeq W(\mathbb{C})/F_{\tilde{z}_G}^0 W(\mathbb{C})$, where $F_{\tilde{z}_G}^0 W(\mathbb{C}) = \exp(F_{\tilde{z}_G}^0 \text{Lie} W_{\mathbb{C}})$. So the fact that $\tilde{Z}_{\tilde{z}_G}^\natural$ is a complex subspace of $\pi^{-1}(\tilde{z}_G)$ implies that W'/U' is a $\text{MT}(\tilde{z}_G) = G_Q$ -module. Hence W' is a G_Q -group.

Define $Q' := W'H_{\tilde{Z}^\natural}$, then Q' is a subgroup of Q since $W' > W_{H_{\tilde{Z}^\natural}}$ and $G_{H_{\tilde{Z}^\natural}}W' = W'$. Now we have that $\tilde{Z}^\natural = H_{\tilde{Z}^\natural}(\mathbb{R})^+ W_{H_{\tilde{Z}^\natural}}(\mathbb{C})\tilde{K}^{\natural,\prime}$ is stable under Q' . So $H_{\tilde{Z}^\natural} = Q'$ because $H_{\tilde{Z}^\natural}$ is the largest subgroup of Q^{der} such that $H_{\tilde{Z}^\natural}(\mathbb{R})^+ W_{H_{\tilde{Z}^\natural}}(\mathbb{C})$ stabilizes $\tilde{Z}^\natural$. So $W_{H_{\tilde{Z}^\natural}} = W'$ is a G_Q -group. Therefore $\tilde{Z} = H_{\tilde{Z}^\natural}(\mathbb{R})^+ U_{H_{\tilde{Z}^\natural}}(\mathbb{C})\tilde{z}$.

Step III Prove $H_{\tilde{Z}^\natural} \triangleleft Q$, and hence $\tilde{Z}$ is weakly special and $H_{\tilde{Z}^\natural} = N$.

First of all, $U_{H_{\tilde{Z}^\natural}} \triangleleft Q$ by Proposition 3.12(2).

Next consider the complex structure of $\pi^{-1}(\tilde{z}_G)$ which comes from $W(\mathbb{R})U(\mathbb{C}) \simeq W(\mathbb{C})/F_{\tilde{z}_G}^0 W(\mathbb{C})$. So the fact that $\tilde{Z}_{\tilde{z}_G}^\natural$ is a complex subspace of $\pi^{-1}(\tilde{z}_G)$ implies that $V_{H_{\tilde{Z}^\natural}}$ is a $\text{MT}(\tilde{z}_G) = G_Q$ -module. Hence $W_{H_{\tilde{Z}^\natural}}$ is a G_Q -group. Besides, $G_{H_{\tilde{Z}^\natural}} \triangleleft G_Q$ by Proposition 11.6. In particular, $G_{H_{\tilde{Z}^\natural}}$ is reductive.

Then let us prove $W_{H_{\tilde{Z}^\natural}} \triangleleft Q$. It suffices to prove $W_{H_{\tilde{Z}^\natural}} \triangleleft W_Q$. For any $\tilde{z} \in \tilde{Z}$, we have proved in Step I that $\tilde{Z}_{\tilde{z}_G} = W_{H_{\tilde{Z}^\natural}}(\mathbb{R})U_{H_{\tilde{Z}^\natural}}(\mathbb{C})\tilde{z}$ is weakly special. Hence there is a connected mixed Shimura subdatum $(R, \mathcal{Z}^+) \hookrightarrow (Q, \mathcal{Y}^+)$ such that $\tilde{z} \in \mathcal{Z}^+$ and $W_{H_{\tilde{Z}^\natural}} \triangleleft R$. Define W^* to be the G_Q -subgroup (of W_Q) generated by $W_R := \mathcal{R}_u(R)$, then $W_{H_{\tilde{Z}^\natural}} \triangleleft W^*$ since $W_{H_{\tilde{Z}^\natural}}$ is a G_Q -group.

Fix a Levi decomposition $H_{\tilde{Z}^\natural} = W_{H_{\tilde{Z}^\natural}} \rtimes G_{H_{\tilde{Z}^\natural}}$ and choose a compatible Levi decomposition $Q = W_Q \rtimes G_Q$ (as is shown in [16, Lemma 9.7]). Let Q^* be the group generated by $G_Q R$, then $\mathcal{R}_u(Q^*) = W^*$ and $Q^*/Q^* = G_Q$. The group Q^* defines a connected mixed Shimura datum $(Q^*, \mathcal{Y}^{*+})$ with $\mathcal{Y}^{*+} = Q^*(\mathbb{R})^+ U_Q^*(\mathbb{C})\tilde{z}$. Now $\tilde{Z} = H_{\tilde{Z}^\natural}(\mathbb{R})^+ U_{H_{\tilde{Z}^\natural}}(\mathbb{C})\tilde{z} \subset \mathcal{Y}^{*+}$. But $\tilde{Z}$ is Hodge generic in $\mathcal{Y}^+$ by assumption, hence $Q = Q^*$ and $W_Q = W^*$. So $W_{H_{\tilde{Z}^\natural}} \triangleleft W_Q$ and hence $W_{H_{\tilde{Z}^\natural}} \triangleleft Q$.

Use the notation in §3.3. We are done if we can prove:

$$(u, v, 1)(0, 0, g)(-u, -v, 1) \in H_{\tilde{Z}^\natural} \text{ for any } u \in U_Q, v \in V_Q \text{ and } g \in G_{H_{\tilde{Z}^\natural}}.$$

By [16, Corollary 2.14], there exist decompositions $U_Q = U_N \oplus U_N^\perp$ and $V_Q = V_N \oplus V_N^\perp$ as G_Q -modules such that G_N acts trivially on $U_N^\perp$ and $V_N^\perp$. Now

$$\begin{aligned} (u, v, 1)(0, 0, g)(-u, -v, 1) &= (u, v, g)(-u, -v, 1) \\ &= (u - g \cdot u, v - g \cdot v, g) \\ &= ((u_N + u_N^\perp) - g \cdot (u_N + u_N^\perp), (v_N + v_N^\perp) - g \cdot (v_N + v_N^\perp), g) \\ &= (u_N - g \cdot u_N, v_N - g \cdot v_N, g) \\ &= (u_N, v_N, 1)(0, 0, g)(-u_N, -v_N, 1) \in H_{\tilde{Z}^\natural}, \end{aligned}$$

where the last inclusion follows from Lemma 11.3.

Conclusion In view of Step I and Step III, we can apply Corollary 6.2 to $\tilde{Z}^\natural$. So $\tilde{Z}^\natural$ is bi-algebraic, and hence is quasi-linear by Theorem 5.8. $\square$

11.2. Estimate. Keep notation and assumptions as in the previous subsection. The group $G_Q^{\text{der}} = H_1 \dots H_r$ is an almost direct product, where H_i 's are non-trivial simple groups and are normal in G_Q^{der} . We have a decomposition

$$(G_Q^{\text{ad}}, \mathcal{Y}_{G_Q}^+) \simeq \prod_{i=1}^r (H_i^{\text{ad}}, \mathcal{Y}_{H_i}^+).$$

Let $S_{G_Q}^{\text{ad}} := \Gamma_{G_Q}^{\text{ad}} \backslash \mathcal{Y}_{G_Q}^+$. Shrinking $\Gamma_{G_Q}^{\text{ad}}$ if necessary, we may assume $S_{G_Q}^{\text{ad}} \simeq \prod_{i=1}^r S_{H_i}$, where S_{H_i} is a connected pure Shimura variety associated with $(H_i^{\text{ad}}, \mathcal{Y}_{H_i}^+)$.

Without loss of generality we may assume $G_N = H_1 \dots H_l$. It suffices to prove $H_i < G_{H_{\tilde{Z}^\natural}}$ for each $i = 1, \dots, l$. The case $l = 0$ is trivial, so we assume that $l \geq 1$. Define $Q_i := \pi^{\natural-1}(H_i)$.

Consider $\mathcal{X}^{\natural+}|_{\mathcal{Y}^+} := \pi^{\natural-1}(\mathcal{Y}^+)$. For the uniformization $\text{unif}^\natural: \mathcal{X}^{\natural+}|_{\mathcal{Y}^+} \rightarrow S^\natural|_{S_Q} := [\pi^\natural]^{-1}(S_Q)$, there exists a fundamental set $\mathfrak{F}^\natural$ such that $\text{unif}^\natural|_{\mathfrak{F}^\natural}$ is definable in the o-minimal theory $\mathbb{R}_{\text{an}, \text{exp}}$: by [16, §10.1] there exists a fundamental set $\mathfrak{F}$ for $\text{unif}_Q: \mathcal{Y}^+ \rightarrow S_Q$ such that $\text{unif}_Q|_{\mathfrak{F}}$ is definable in $\mathbb{R}_{\text{an}, \text{exp}}$. It suffices to take

$$\mathfrak{F}^\natural := \pi_{P/U}^{\natural-1}(\mathfrak{F}_{Q/U_Q}) \times_{\mathfrak{F}_{Q/U_Q}} \mathfrak{F} \subset \pi_{P/U}^{\natural-1}(\mathcal{Y}_{Q/U_Q}^+) \times_{\mathcal{Y}_{Q/U_Q}^+} \mathcal{Y}^+ = \mathcal{X}^{\natural+}|_{\mathcal{Y}^+}.$$

We have $\tilde{Y}_G = \prod_{i=1}^l H_i(\mathbb{R})^+ \tilde{Z}_G$ by Lemma 11.5(2). Define the following Shimura morphisms for each $i = 1, \dots, l$

$$\begin{array}{ccc} (G_Q, \mathcal{Y}_{G_Q}^+) & \xrightarrow{p_i} & (G_i, \mathcal{Y}_{G,i}^+) := (G_Q^{\text{ad}}, \mathcal{Y}_{G_Q}^+) / \prod_{j \neq i} H_j^{\text{ad}} \\ \downarrow \text{unif}_{G_Q} & & \downarrow \text{unif}_{G,i} \\ S_{G_Q} & \xrightarrow{[p_i]} & S_{G,i} \end{array}$$

Recall that the $\mathfrak{F}$ we take as in [16, §10.1] satisfies $\mathfrak{F}_G = \prod_{i=1}^r \mathfrak{F}_{G,i}$ where $\mathfrak{F}_{G,i} = p_i(\mathfrak{F}_G)$. Fix $i \in \{1, \dots, l\}$. Define $\tilde{Y}_{G,i} := p_i(\tilde{Y}_G) = H_i^{\text{ad}}(\mathbb{R})^+ \pi_i(\tilde{Z}_G)$, $\tilde{Z}_{G,i} := p_i(\tilde{Z}_G)$ and $Y_{G,i} := [p_i](Y_G)$, then $\text{unif}_{G,i}(\tilde{Z}_{G,i})$ is Zariski dense in $Y_{G,i}$ by Lemma 11.5. If $\dim(\tilde{Z}_{G,i}) = 0$, then $\tilde{Z}_{G,i}$ is a finite set of points since it is algebraic. But then $\text{unif}_{G,i}(\tilde{Z}_{G,i})$, and hence $Y_{G,i} = \text{unif}_{G,i}(\tilde{Z}_{G,i})^{\text{Zar}}$ is also a finite set of points. So $\dim(Y_{G,i}) = 0$, which contradicts $\tilde{Y}_{G,i} = H_i^{\text{ad}}(\mathbb{R})^+ p_i(\tilde{Z}_G)$. To sum it up, we have $\dim(\tilde{Z}_{G,i}) > 0$. For convenience (and a slight abuse of notation), we will denote by

$$\pi_i^{\natural} (:= p_i \circ \pi_i^{\natural}|_{\mathcal{X}^{\natural+}|_{\mathcal{Y}^+}}) : \mathcal{X}^{\natural+}|_{\mathcal{Y}^+} = \pi^{\natural-1}(\mathcal{Y}^+) \rightarrow \mathcal{Y}_{G,i}^+ \text{ and } \pi_i^{\natural} : Q \rightarrow G_i$$

Fix a point $\tilde{z}^{\natural} \in \mathfrak{F}^{\natural} \cap \tilde{Z}^{\natural}$. Take an algebraic curve $C_{G,i} \subset \tilde{Z}_{G,i}$ passing through $\pi_i^{\natural}(\tilde{z}^{\natural})$. Now $\pi_i^{\natural}(\tilde{Z}^{\natural} \cap \pi_i^{\natural-1}(C_{G,i})) = \tilde{Z}_{G,i} \cap C_{G,i} = C_{G,i}$, and hence there exists an algebraic curve $C^{\natural} \subset \tilde{Z}^{\natural} \cap \pi_i^{\natural-1}(C_{G,i})$ passing through $\tilde{z}^{\natural}$ such that $\dim(\pi_i(C^{\natural})) = 1$.

The set $\mathfrak{F}_{G,i} = p_i(\mathfrak{F}_G)$ is a fundamental set of $\text{unif}_{G,i}$ and $\text{unif}_{G,i}|_{\mathfrak{F}_{G,i}}$ is definable. We define for any irreducible semi-algebraic subvariety $A^{\natural}$ (resp. $A_{G,i}$) of $\text{unif}^{\natural-1}(\mathcal{Y}^{\natural})$ (resp. $\text{unif}_{G,i}^{-1}(Y_{G,i})$) the following sets: define

$$\begin{aligned} \Sigma^{(i)}(A^{\natural}) &:= \{g \in Q_i(\mathbb{R}) \mid \dim(gA^{\natural} \cap \text{unif}^{\natural-1}(\mathcal{Y}^{\natural}) \cap \mathfrak{F}^{\natural}) = \dim(A^{\natural})\} \\ (\text{resp. } \Sigma_G^{(i)}(A_{G,i}) &:= \{g \in H_i^{\text{ad}}(\mathbb{R}) \mid \dim(gA_{G,i} \cap \text{unif}_{G,i}^{-1}(\overline{Y_{G,i}}) \cap \mathfrak{F}_{G,i}) = \dim(A_{G,i})\}) \end{aligned}$$

and

$$\begin{aligned} \Sigma'^{(i)}(A^{\natural}) &:= \{g \in Q_i(\mathbb{R}) \mid g^{-1}\mathfrak{F}^{\natural} \cap A^{\natural} \neq \emptyset\} \\ (\text{resp. } \Sigma_G'^{(i)}(A_{G,i}) &:= \{g \in H_i^{\text{ad}}(\mathbb{R}) \mid g^{-1}\mathfrak{F}_{G,i} \cap A_{G,i} \neq \emptyset\}) \end{aligned}$$

Then $\Sigma^{(i)}(A^{\natural})$ and $\Sigma_G^{(i)}(A_{G,i})$ are by definition definable. Let $\Gamma_{G,i}^{\text{ad}} := p_i(\Gamma_Q^{\text{ad}})$.

Lemma 11.8. $\Sigma'^{(i)}(A^{\natural}) \cap \Gamma_Q = \Sigma^{(i)}(A^{\natural}) \cap \Gamma_Q$ (resp. $\Sigma_G'^{(i)}(A_{G,i}) \cap \Gamma_{G,i}^{\text{ad}} = \Sigma_G^{(i)}(A_{G,i}) \cap \Gamma_{G,i}^{\text{ad}}$).

Proof. First $\Sigma^{(i)}(A^{\natural}) \cap \Gamma_Q \subset \Sigma'^{(i)}(A^{\natural}) \cap \Gamma_Q$ by definition. Conversely for any $\gamma \in \Sigma'^{(i)}(A^{\natural}) \cap \Gamma_Q$, the set $\gamma^{-1}\mathfrak{F}^{\natural} \cap A^{\natural}$ contains an open subspace of $A^{\natural}$ since $\mathfrak{F}^{\natural}$ is by choice open in $\mathcal{X}^{\natural+}|_{\mathcal{Y}^+}$. Hence $\gamma A^{\natural} \cap \text{unif}^{\natural-1}(\mathcal{Y}^{\natural}) \cap \mathfrak{F}^{\natural} = \gamma A^{\natural} \cap \mathfrak{F}^{\natural}$ contains an open subspace of $\gamma A^{\natural}$ which must be of dimension $\dim(A^{\natural})$. Hence $\gamma \in \Sigma^{(i)}(A^{\natural}) \cap \Gamma_Q$. The proof for $A_{G,i}$ is the same. $\square$

This lemma implies

$$(11.2) \quad \begin{aligned} \Sigma^{(i)}(C^{\natural}) \cap \Gamma_Q &= \Sigma'^{(i)}(C^{\natural}) \cap \Gamma_Q \subset \Sigma'^{(i)}(\tilde{Z}^{\natural}) \cap \Gamma_Q = \Sigma^{(i)}(\tilde{Z}^{\natural}) \cap \Gamma_Q \\ (\text{resp. } \Sigma_G^{(i)}(C_{G,i}) \cap \Gamma_{G,i}^{\text{ad}} &= \Sigma_G'^{(i)}(C_{G,i}) \cap \Gamma_{G,i}^{\text{ad}} \subset \Sigma_G'^{(i)}(\tilde{Z}_{G,i}) \cap \Gamma_{G,i}^{\text{ad}} = \Sigma^{(i)}(\tilde{Z}_{G,i}) \cap \Gamma_{G,i}^{\text{ad}}) \end{aligned}$$

Lemma 11.9. $\pi_i^{\natural}(\Gamma_Q \cap \Sigma^{(i)}(C^{\natural})) = \Gamma_{G,i}^{\text{ad}} \cap \Sigma_G^{(i)}(C_{G,i})$.

Proof. By Lemma 11.8, it suffices to prove $\pi_i^{\natural}(\Gamma \cap \Sigma'^{(i)}(C^{\natural})) = \Gamma_{G,i}^{\text{ad}} \cap \Sigma_G'^{(i)}(C_{G,i})$. The inclusion $\subset$ is clear by definition. For the other inclusion, $\forall \gamma_{G,i} \in \Gamma_{G,i}^{\text{ad}} \cap \Sigma_G'^{(i)}(C_{G,i})$, $\exists c_{G,i} \in C_{G,i}$ such that $\gamma_{G,i} \cdot c_{G,i} \in \mathfrak{F}_{G,i}$.

Take a point $c^{\natural} \in C^{\natural}$ such that $\pi_i^{\natural}(c^{\natural}) = c_{G,i}$ and define $c_G := \pi^{\natural}(c^{\natural}) \in \mathcal{Y}_{G_Q}^+$. Suppose $c_G = (c_{G,1}, \dots, c_{G,r})$ under the decomposition $(G_Q^{\text{ad}}, \mathcal{Y}_{G_Q}^+) \simeq \prod_{i=1}^r (H_i^{\text{ad}}, \mathcal{X}_{H,i}^+)$. Then by choice of $\mathfrak{F}_G$, there exists $\gamma'_G \in \Gamma_{G_Q}^{\text{ad}}$ whose i -th coordinate is precisely the $\gamma_{G,i}$ in the last paragraph such that $\gamma'_G \cdot c_G \in \mathfrak{F}_G$.

Let $\gamma_G \in \Gamma_{G_Q}$ be such that its image under $\Gamma_{G_Q} \rightarrow \Gamma_{G_Q}^{\text{ad}}$ is γ'_G , then $\gamma_G \cdot c^{\natural} \in \pi^{\natural-1}(\mathfrak{F}_G)$. Therefore there exist $\gamma_V \in \Gamma_{V_Q}$, $\gamma_U \in \Gamma_{U_Q}$ such that $(\gamma_U, \gamma_V, \gamma_G)c^{\natural} \in \mathfrak{F}^{\natural}$. Denote by $\gamma = (\gamma_U, \gamma_V, \gamma_G)$, then $\gamma \in \Gamma_Q \cap \Sigma'^{(i)}(C^{\natural})$ and $\pi_i^{\natural}(\gamma) = \gamma_{G,i}$. $\square$

For $T > 0$, define

$$\Theta_G^{(i)}(C_{G,i}, T) := \{\gamma_G \in \Gamma_{G,i}^{\text{ad}} \cap \Sigma_G^{(i)}(C_{G,i}) \mid H(\gamma_G) \leq T\}.$$

Proposition 11.10. *There exists a constant $\delta > 0$ s.t. for all $T \gg 0$, $|\Theta_G^{(i)}(C_{G,i}, T)| \geq T^{\delta}$.*

Proof. This follows directly from [21, Theorem 1.3] applied to $((G_i, \mathcal{X}_{G,i}^+, S_{G,i}, \tilde{Z}_{G,i}))$. $\square$

Let us prove how these facts imply $H_i < G_{H_{\tilde{Z}^{\natural}}}$. Take a faithful representation $G_Q^{\text{ad}} \hookrightarrow \text{GL}_n$ which sends $\Gamma_{G_Q}^{\text{ad}}$ to $\text{GL}_n(\mathbb{Z})$. Consider the definable set $\Sigma_G^{(i)}(C_{G,i})$. By the theorem of Pila-Wilkie ([29, Theorem 3.6]), there exist $J = J(\delta)$ definable block families

$$B^j \subset \Sigma_G^{(i)}(C_{G,i}) \times \mathbb{R}^l, \quad j = 1, \dots, J$$

and $c = c(\delta) > 0$ such that for all $T \gg 0$, $\Theta_G^{(i)}(C_{G,i}, T^{1/2n})$ is contained in the union of at most $cT^{\delta/4n}$ definable blocks of the form B_y^j ($y \in \mathbb{R}^l$). By Proposition 11.10, there exist a $j \in \{1, \dots, J\}$ and a block $B_{G,i} := B_{y_0}^j$ of $\Sigma_G^{(i)}(C_{G,i})$ containing at least $T^{\delta/4n}$ elements of $\Theta_G^{(i)}(C_{G,i}, T^{1/2n})$.

Let $\Sigma^{(i)} := \Sigma^{(i)}(C^{\natural}) \cap \Sigma^{(i)}(\tilde{Z}^{\natural})$, which is by definition a definable set. Consider $X^j := (\pi_i^{\natural} \times 1_{\mathbb{R}^l})^{-1}(B^j) \cap (\Sigma^{(i)} \times \mathbb{R}^l)$, which is a definable family since $\pi_i^{\natural}$ is algebraic. By [44, Ch. 3, 3.6], there exists a number $n_0 > 0$ such that each fibre X_y^j has at most n_0 connected components. So the definable set $\pi_i^{\natural-1}(B_{G,i}) \cap \Sigma^{(i)}$ has at most n_0 connected components. Now

$$\pi_i^{\natural}(\pi_i^{\natural-1}(B_{G,i}) \cap \Sigma^{(i)} \cap \Gamma_Q) = B_{G,i} \cap \pi_i^{\natural}(\Sigma^{(i)}(C^{\natural}) \cap \Gamma_Q) = B_{G,i} \cap \Sigma_G^{(i)}(C_{G,i}) \cap \Gamma_{G,i}^{\text{ad}} = B_{G,i} \cap \Gamma_{G,i}^{\text{ad}}$$

by (11.2) and Lemma 11.9. So there exists a connected component B of $\pi_i^{\natural-1}(B_{G,i}) \cap \Sigma^{(i)}$ such that $\pi_i^{\natural}(B \cap \Gamma_Q)$ contains at least $T^{\delta/4n}/n_0$ elements of $\Theta_G^{(i)}(C_{G,i}, T^{1/2n})$.

We have $B\tilde{Z}^{\natural} \subset \text{unif}^{\natural-1}(Y^{\natural})$ since $\Sigma^{(i)}(\tilde{Z}^{\natural})\tilde{Z}^{\natural} \subset \text{unif}^{\natural-1}(Y^{\natural})$ by analytic continuation, and $\tilde{Z}^{\natural} \subset \sigma^{-1}B\tilde{Z}^{\natural}$ for any $\sigma \in B \cap \Gamma_Q$. But B is connected, and therefore $\sigma^{-1}B\tilde{Z}^{\natural} = \tilde{Z}^{\natural}$ by maximality of $\tilde{Z}^{\natural}$ and [30, Lemma 4.1]. So $B \subset \sigma \text{Stab}_{Q_i(\mathbb{R})}(\tilde{Z}^{\natural})$ for any $\sigma \in B \cap \Gamma_Q$.

Fix a $\gamma_0 \in B \cap \Gamma_Q$ such that $\pi_i^{\natural}(\gamma_0) \in \Theta_G^{(i)}(C_{G,i}, T^{1/2n})$. We have already shown that $\pi_i^{\natural}(B \cap \Gamma_Q)$ contains at least $T^{\delta/4n}/n_0$ elements of $\Theta_G^{(i)}(C_{G,i}, T^{1/2n})$. For any $\gamma'_{G,i} \in \pi_i^{\natural}(B \cap \Gamma_Q)$, let γ' be one of its pre-images in $B \cap \Gamma_Q$. Then $\gamma := \gamma'^{-1}\gamma_0$ is an element of $\Gamma_Q \cap \text{Stab}_{Q_i(\mathbb{R})}(\tilde{Z}^{\natural}) = \Gamma_Q \cap Q_i(\mathbb{R})$ such that $H(\pi_i^{\natural}(\gamma)) \ll T^{1/2}$. Therefore for $T \gg 0$,

$\pi_i^{\natural}(\Gamma_{\tilde{Z}^{\natural}}) \cap H_i^{\text{ad}}(\mathbb{R})$ contains at least $T^{\delta/4n}/n_0$ elements $\gamma_{G,i}$ such that $H(\gamma_{G,i}) \leq T$. Hence $\dim(\pi_i^{\natural}(H_{\tilde{Z}^{\natural}}) \cap H_i^{\text{ad}}) > 0$ since $\pi_i^{\natural}(H_{\tilde{Z}^{\natural}}) \cap H_i^{\text{ad}}$ contains infinitely many rational points. But $\pi_i^{\natural}(H_{\tilde{Z}^{\natural}}) = p_i \pi_i^{\natural}(H_{\tilde{Z}^{\natural}}) = p_i(G_{H_{\tilde{Z}^{\natural}}})$ by definition. So $H_i^{\text{ad}} < p_i(G_{H_{\tilde{Z}^{\natural}}})$ since H_i^{ad} is simple and $p_i(G_{H_{\tilde{Z}^{\natural}}}) \cap H_i^{\text{ad}} \triangleleft H_i^{\text{ad}}$ by Corollary 11.4.

As a normal subgroup of G_N , $G_{H_{\tilde{Z}^{\natural}}}$ is the almost direct product of some H_j 's ($j = 1, \dots, l$). So $H_i^{\text{ad}} < p_i(G_{H_{\tilde{Z}^{\natural}}})$ implies $H_i < G_{H_{\tilde{Z}^{\natural}}}$. Now we are done.

REFERENCES

- [1] Y. André. Mumford-Tate groups of mixed Hodge structures and the theorem of the fixed part. *Compos. Math.*, 82(1):1–24, 1992.
- [2] J. Ax. On Schanuel's conjectures. *Annals Math.*, 93:252–268, 1971.
- [3] J. Ax. Some topics in differential algebraic geometry I: Analytic subgroups of algebraic groups. *American Journal of Mathematics*, 94:1195–1204, 1972.
- [4] B. Barbieri-Viale and A. Bertapelle. Sharp de Rham realization. *Advances in Math.*, 222:1308–1338, 2009.
- [5] D. Bertrand, D. Masser, A. Pillay, and U. Zannier. Relative Manin-Mumford for semi-abelian surfaces. *Proceedings of the Edinburgh Mathematical Society*, page online, 2016.
- [6] D. Bertrand and A. Pillay. A Lindemann-Weierstrass theorem for semi-abelian varieties over function fields. *J. Amer. Math. Soc.*, 23(2):491–533, 2010.
- [7] D. Bertrand and A. Pillay. Galois theory, function Lindemann-Weierstrass, and Manin maps. *Pacific J. of Math.*, 281:51–82, 2016.
- [8] A. Buium. *Differential Algebraic Groups of Finite Dimension*. Number 1506 in LNM. Springer-Verlag, 1992.
- [9] A. Buium. Effective bound for the geometric Lang conjecture. *Duke Math. J.*, 71:475–499, 1993.
- [10] A. Buium. Geometry of differential polynomial functions I: algebraic groups. *American Journal of Mathematics*, 115:1385–1444, 1993.
- [11] P. Cohen. Humbert surfaces and transcendence properties of automorphic functions. *Rocky Mountain J. Math.*, 26:987–1001, 1996.
- [12] R. Coleman. The universal vectorial bi-extension and p -adic heights. *Inv. Math.*, 103:631–650, 1991.
- [13] C. Daw and M. Orr. Heights of pre-special points of Shimura varieties. *Math. Annalen*, online.
- [14] Z. Gao. *Le théorème d'Ax-Lindemann et ses applications à la conjecture de Zilber-Pink (The mixed Ax-Lindemann theorem and its applications to the Zilber-Pink conjecture)*. PhD thesis, Leiden University and Université Paris-Sud, 2014.
- [15] Z. Gao. About the mixed André-Oort conjecture: reduction to a lower bound for the pure case. *Comptes rendus Mathématiques*, 354:659–663, 2016.
- [16] Z. Gao. Towards the André-Oort conjecture for mixed Shimura varieties: the Ax-Lindemann-weierstrass theorem and lower bounds for Galois orbits of special points. *J. Reine Angew. Math (Crelle)*, online, 2015.
- [17] Z. Gao. A special point problem of André-Pink-Zannier in the universal family of abelian varieties. *Annali della Scuola Normale Superiore di Pisa, Classe di Scienze*, to appear.
- [18] P. Habegger and J. Pila. Some unlikely intersections beyond André-Oort. *Compos. Math.*, 148(01):1–27, January 2012.
- [19] P. Habegger and J. Pila. O-minimality and certain atypical intersections. *Annales scientifiques de l'Ecole Normale Supérieure*, to appear.
- [20] J. Hwang and W. To. Volumes of complex analytic subvarieties of Hermitian symmetric spaces. *American Journal of Mathematics*, 124(6):1221–1246, 2002.
- [21] B. Klingler, E. Ullmo, and A. Yafaev. The hyperbolic Ax-Lindemann-Weierstrass conjecture. *Publ. math. IHES*, to appear.
- [22] G. Laumon. Transformation de Fourier généralisée. <http://arxiv.org/abs/alg-geom/9603004>, Preprint IHES.
- [23] Y. Manin. Algebraic curves over fields with differentiation. *Izv. Akad. Nauk SSSR. Ser. Mat.*, 22:737–756, 1958.
- [24] Y. Manin. Rational points on algebraic curves over function fields. *Izv. Akad. Nauk SSSR. Ser. Mat.*, 27:1395–1440, 1963.

- [25] B. Moonen. Linearity properties of Shimura varieties, I. *Journal of Algebraic Geometry*, 7(3):539–467, 1988.
- [26] M. Orr. Families of abelian varieties with many isogenous fibres. *J.Reine Angew. Math (Crelle)*, to appear.
- [27] C. Peters and J. Steenbrink. *Mixed Hodge Structures*, volume 52 of *A Series of Modern Surveys in Mathematics*. Springer, 2008.
- [28] Y. Peterzil and S. Starchenko. Complex analytic geometry and analytic-geometric categories. *J.Reine Angew. Math (Crelle)*, 626:39–74, 2009.
- [29] J. Pila. O-minimality and the André-Oort conjecture for $\mathbb{C}^n$. *Annals Math.*, 173:1779–1840, 2011.
- [30] J. Pila and J. Tsimerman. The André-Oort conjecture for the moduli space of Abelian surfaces. *Compos. Math.*, 149:204–216, February 2013.
- [31] J. Pila and J. Tsimerman. Ax-Lindemann for $\mathcal{A}_g$. *Annals Math.*, 179:659–681, 2014.
- [32] J. Pila and J. Tsimerman. Ax-Schanuel for the j -function. *Duke Math. J.*, To appear.
- [33] R. Pink. *Arithmetical compactification of mixed Shimura varieties*. PhD thesis, Bonner Mathematische Schriften, 1989.
- [34] R. Pink. A combination of the conjectures of Mordell-Lang and André-Oort. In *Geometric Methods in Algebra and Number Theory*, volume 253 of *Progress in Mathematics*, pages 251–282. Birkhäuser, 2005.
- [35] V. Platonov and A. Rapinchuk. *Algebraic Groups and Number Theory*. Academic Press, INC., 1994.
- [36] H. Shiga and J. Wolfart. Criteria for complex multiplication and transcendence properties of automorphic functions. *J.Reine Angew. Math (Crelle)*, 463:1–25, 1995.
- [37] J. Tsimerman. Ax-Schanuel and o-minimality. In *O-Minimality and Diophantine Geometry*, number 421 in London Math. Soc. Lecture Note Series. Camb. Univ. Press, 2015.
- [38] J. Tsimerman. A proof of the André-Oort conjecture for $\mathcal{A}_g$. *Preprint, available on the author's page*, 2015.
- [39] E. Ullmo. Applications du théorème de Ax-Lindemann hyperbolique. *Compos. Math.*, 150:175–190, 2014.
- [40] E. Ullmo. Structures spéciales et problème de Pink-Zilber. *Astérisque*. to appear.
- [41] E. Ullmo and A. Yafaev. A characterisation of special subvarieties. *Mathematika*, 57(2):263–273, 2011.
- [42] E. Ullmo and A. Yafaev. Nombre de classes des tores de multiplication complexe et bornes inférieures pour orbites Galoisiennes de points spéciaux. *Bull. de la SMF*, 143:197–228, 2015.
- [43] E. Ullmo and A. Yafaev. The hyperbolic Ax-Lindemann in the compact case. *Duke Math. J.*, to appear.
- [44] L. van der Dries. *Tame topology and o-minimal structures*, volume 248 of *London Math. Soc. Lecture Note Series*. Camb. Univ. Press, 1998.
- [45] G. Wüstholz. Algebraic groups, Hodge theory, and transcendence. In *Proceedings of the International Congress of Mathematicians*, volume 1,2, pages 476–483, 1986.

CNRS AND INSTITUT MATHÉMATIQUES DE JUSSIEU-PARIS RIVE GAUCHE, 4 PLACE DE JUSSIEU, 75005 PARIS, FRANCE

E-mail address: ziyang.gao@imj-prg.fr